



ELSEVIER

www.elsevier.nl/locate/jorgchem

Journal of Organometallic Chemistry 602 (2000) 217–221

Journal  
of Organo  
metallic  
Chemistry

## Subject Index of Volume 602

### Ab initio calculations

Dimethylaluminium enolates and alkoxides derived from trimethylaluminium and aromatic ketones: a synthetic, structural and theoretical investigation, 15

### Acetylacetone

Rh(acac)(CO)(PR<sub>3</sub>) and Rh(oxinate)(CO)(PR<sub>3</sub>) complexes—substitution chemistry and structural aspects, 59

### Alkene

Copper(I) complexes with conjugated dienes, 5

### Alkylations

Dimethylaluminium enolates and alkoxides derived from trimethylaluminium and aromatic ketones: a synthetic, structural and theoretical investigation, 15

### Alkyne

A mononuclear alkyne-stabilized Mo<sup>(IV)</sup>O<sub>2</sub> moiety, 72

Copper(I) complexes with conjugated dienes, 5

### Alkynes

Heterobimetallische Komplexe von Titan(IV)-Quecksilber(II) und Platin(II)-Quecksilber(II), 91

### Alkynols

Organotransition metal-modified sugars: Part 16. Diastereoselective synthesis of carbohydrate-derived spirocyclic 2-oxacyclohexylidene chromium and tungsten complexes, 37

### Alkynyl complex

π-Allyliridium(I) complexes, Ir(η<sup>3</sup>-CH<sub>2</sub>CHCHAR)(CO)(PPh<sub>3</sub>)<sub>2</sub> (Ar = Ph, C<sub>6</sub>H<sub>4</sub>Me-*p*, C<sub>6</sub>H<sub>4</sub>Br-*p*). Comparison of their structures and chemical properties with analogous Rh complexes, 144

### Allene

π-Allyliridium(I) complexes, Ir(η<sup>3</sup>-CH<sub>2</sub>CHCHAR)(CO)(PPh<sub>3</sub>)<sub>2</sub> (Ar = Ph, C<sub>6</sub>H<sub>4</sub>Me-*p*, C<sub>6</sub>H<sub>4</sub>Br-*p*). Comparison of their structures and chemical properties with analogous Rh complexes, 144

### π-Allyl complex

π-Allyliridium(I) complexes, Ir(η<sup>3</sup>-CH<sub>2</sub>CHCHAR)(CO)(PPh<sub>3</sub>)<sub>2</sub> (Ar = Ph, C<sub>6</sub>H<sub>4</sub>Me-*p*, C<sub>6</sub>H<sub>4</sub>Br-*p*). Comparison of their structures and chemical properties with analogous Rh complexes, 144

### Aluminium

Dimethylaluminium enolates and alkoxides derived from trimethylaluminium and aromatic ketones: a synthetic, structural and theoretical investigation, 15

### Aluminum

Precursor chemistry of Group III nitrides: Part XVI. Synthesis and structure of monomeric penta-coordinated intramolecularly adduct-stabilized amidobisazides of aluminum, gallium and indium with an all-nitrogen coordination sphere: OMCVD of GaN using (N<sub>3</sub>)<sub>2</sub>Ga{N[CH<sub>2</sub>CH<sub>2</sub>(NEt<sub>2</sub>)<sub>2</sub>]}, 29

### Amide

Precursor chemistry of Group III nitrides: Part XVI. Synthesis and structure of monomeric penta-coordinated intramolecularly adduct-stabilized amidobisazides of aluminum, gallium and indium with an all-nitrogen coordination sphere: OMCVD of GaN using (N<sub>3</sub>)<sub>2</sub>Ga{N[CH<sub>2</sub>CH<sub>2</sub>(NEt<sub>2</sub>)<sub>2</sub>]}, 29

### Aminoethyl ferrocene

Synthesis and reactions of the ferrocene derived hydroxymethyl phosphine FeCH(CH<sub>3</sub>)P(CH<sub>2</sub>OH)<sub>2</sub> and its sulfide: crystal structures of [FeCH(CH<sub>3</sub>)P(S)R<sub>2</sub>] R = CH<sub>2</sub>OH, CH<sub>2</sub>CH<sub>2</sub>CN and FeCH(CH<sub>3</sub>)P(S)(CH<sub>2</sub>O)<sub>2</sub>PPh (Fc = ferrocenyl), 125

### Arene complexes

Reactivity of dinuclear arene ruthenium complexes: reactions of the hydrido complex [(*p*-Me-C<sub>6</sub>H<sub>4</sub>-Pr')<sub>2</sub>Ru<sub>2</sub>Cl<sub>2</sub>(μ-Cl)(μ-H)] with NaX and HX (X = F, Cl, Br, I), 188

### Aurophilic

Aurophilic association in *endo*-dicyclopentadienechlorogold(I), 133

### Azide

Precursor chemistry of Group III nitrides: Part XVI. Synthesis and structure of monomeric penta-coordinated intramolecularly adduct-stabilized amidobisazides of aluminum, gallium and indium with an all-nitrogen coordination sphere: OMCVD of GaN using (N<sub>3</sub>)<sub>2</sub>Ga{N[CH<sub>2</sub>CH<sub>2</sub>(NEt<sub>2</sub>)<sub>2</sub>]}, 29

### bis(Diphenylphosphino)amine complexes

Iridium carbonyl cluster complexes with bidentate phosphine ligands; the X-ray crystal structure of [Ir<sub>4</sub>(CO)<sub>8</sub>(dppa)<sub>2</sub>]<sub>3</sub>(THF), 137

### bis(Diphenylphosphino)ethene complexes

Iridium carbonyl cluster complexes with bidentate phosphine ligands; the X-ray crystal structure of [Ir<sub>4</sub>(CO)<sub>8</sub>(dppa)<sub>2</sub>]<sub>3</sub>(THF), 137

### Bis(trimethylsilyl)methyl ligand

Zur Elektronenstruktur metallorganischer Komplexe der f-Elemente IL. Erstmalige Aufklärung der Elektronenstruktur eines metallorganischen σ-Komplexes der f-Elemente: Tris(bis(trimethylsilyl)methyl)erbium(III), 65

### Boron

New zwitterionic heterocycles derived from 1-chloro-(dimethyl)silyl-1-borylalkenes and 2-lithio-1-methylimidazole and 1-lithio-indazole, 45

### Butadiene

Copper(I) complexes with conjugated dienes, 5

### Carbene complexes

Organotransition metal-modified sugars: Part 16. Diastereoselective synthesis of carbohydrate-derived spirocyclic 2-oxacyclohexylidene chromium and tungsten complexes, 37

### Carbocation

The behavior of 3-ferrocenyl-1-methyl-1,2-pentamethyleneallyl and 1,5-diferrocenyl-3-methyl-2,4-tetramethylene-1,4-dienyl carbocations in the cationic dimerization of 1,3-dienes, 105

### Carbohydrates

Organotransition metal-modified sugars: Part 16. Diastereoselective synthesis of carbohydrate-derived spirocyclic 2-oxacyclohexylidene chromium and tungsten complexes, 37

### Carbonyl

Iridium carbonyl cluster complexes with bidentate phosphine ligands; the X-ray crystal structure of [Ir<sub>4</sub>(CO)<sub>8</sub>(dppa)<sub>2</sub>]<sub>3</sub>(THF), 137

## Catalysis

A mononuclear alkyne-stabilized Mo<sup>(IV)</sup>O<sub>2</sub> moiety, 72  
*Cis*- and *trans*-bis(diphenylphosphino)ethene  
*Cis*- and *trans*-bis(diphenylphosphino)ethene seven-coordinate complexes of molybdenum(II) and tungsten(II). Crystal structures of [MI<sub>2</sub>(CO)<sub>3</sub>(*cis*-dppen)] {M = Mo or W; dppen = bis(diphenylphosphino)ethene}, [WI(CO)<sub>2</sub>(*cis*-dppen)<sub>2</sub>]I, [MI<sub>2</sub>(CO)<sub>2</sub>{P(OR)<sub>3</sub>}(*cis*-dppen)] (M = Mo, R = Me; M = W, R = Et or Ph), 115

## Cluster complexes

Iridium carbonyl cluster complexes with bidentate phosphine ligands; the X-ray crystal structure of [Ir<sub>4</sub>(CO)<sub>8</sub>(dppta)<sub>2</sub>]<sup>3</sup>(THF), 137

## Cobaloxime

Organocobaloximes with mixed-dioxime equatorial ligands: a one-pot synthesis, 1

## Cobalt

Organocobaloximes with mixed-dioxime equatorial ligands: a one-pot synthesis, 1

## Copolymerization

Catalytic activity of cationic diphas palladium(II) complexes in the alkene/CO copolymerization in organic solvents and water in dependence on the length of the alkyl chain at the phosphine ligands, 173

## Copper(I)

Copper(I) complexes with conjugated dienes, 5

## Crystal field analysis

Zur Elektronenstruktur metallorganischer Komplexe der f-Elemente IL. Erstmalige Aufklärung der Elektronenstruktur eines metallorganischen σ-Komplexes der f-Elemente: Tris(trimethylsilyl)methyl)erbium(III), 65

## Crystal structure

Aurophilic association in *endo*-dicyclopentadienechlorogold(I), 133

## Copper(I) complexes with conjugated dienes, 5

Dimethylaluminium enolates and alkoxides derived from trimethylaluminium and aromatic ketones: a synthetic, structural and theoretical investigation, 15

## Crystal structures

*Cis*- and *trans*-bis(diphenylphosphino)ethene seven-coordinate complexes of molybdenum(II) and tungsten(II). Crystal structures of [MI<sub>2</sub>(CO)<sub>3</sub>(*cis*-dppen)] {M = Mo or W; dppen = bis(diphenylphosphino)ethene}, [WI(CO)<sub>2</sub>(*cis*-dppen)<sub>2</sub>]I, [MI<sub>2</sub>(CO)<sub>2</sub>{P(OR)<sub>3</sub>}(*cis*-dppen)] (M = Mo, R = Me; M = W, R = Et or Ph), 115

Use of the aqua complex [*fac*-Mn(CO)<sub>3</sub>(dppe)OH<sub>2</sub>]BF<sub>4</sub> for the preparation of a series of *fac*-Mn(CO)<sub>3</sub>(dppe)Z and [*fac*-Mn(CO)<sub>3</sub>(dppe)Z]BF<sub>4</sub> complexes. X-ray crystal structures of three representative examples, 97

## Cyclic voltammetry

Electrochemistry of ruthenium metallocenes: Part 3. Synthesis and properties of ruthenium [2<sub>2</sub>]paracyclophane complexes with methacrylic acid and methacrylate ester substituents, 75

## Cycloaddition

The behavior of 3-ferrocenyl-1-methyl-1,2-pentamethyleneallyl and 1,5-diferrocenyl-3-methyl-2,4-tetramethylene-1,4-dienyl carbocations in the cationic dimerization of 1,3-dienes, 105

## Cyclodimerization

The behavior of 3-ferrocenyl-1-methyl-1,2-pentamethyleneallyl and 1,5-diferrocenyl-3-methyl-2,4-tetramethylene-1,4-dienyl carbocations in the cationic dimerization of 1,3-dienes, 105

## Cyclopentadienyl complexes

Silane dendrimers containing titanium complexes on their periphery, 208

## Cyclophanes

Electrochemistry of ruthenium metallocenes: Part 3. Synthesis and properties of ruthenium [2<sub>2</sub>]paracyclophane complexes with

methacrylic acid and methacrylate ester substituents, 75

## Deprotonation

Reactivity of Pd(II) complexes containing the orthometallated C,C-chelating ligand C<sub>6</sub>H<sub>4</sub>-2-PPh<sub>2</sub>C(H)COCH<sub>2</sub>PPh<sub>3</sub> towards deprotonating reagents. Part 2, 151

The behavior of 3-ferrocenyl-1-methyl-1,2-pentamethyleneallyl and 1,5-diferrocenyl-3-methyl-2,4-tetramethylene-1,4-dienyl carbocations in the cationic dimerization of 1,3-dienes, 105

## Dimethylbutadiene

Copper(I) complexes with conjugated dienes, 5

## Dimethyl zirconium complexes

Monocyclopentadienyl and mono(2-methoxyethyl)cyclopentadienyl zirconium methyl complexes, 51

## Dioxime

Organocobaloximes with mixed-dioxime equatorial ligands: a one-pot synthesis, 1

## Electrode modification

Electrochemistry of ruthenium metallocenes: Part 3. Synthesis and properties of ruthenium [2<sub>2</sub>]paracyclophane complexes with methacrylic acid and methacrylate ester substituents, 75

## Electronic factor

The behavior of 3-ferrocenyl-1-methyl-1,2-pentamethyleneallyl and 1,5-diferrocenyl-3-methyl-2,4-tetramethylene-1,4-dienyl carbocations in the cationic dimerization of 1,3-dienes, 105

*endo*-Dicyclopentadiene

Aurophilic association in *endo*-dicyclopentadienechlorogold(I), 133

## Enolates

Dimethylaluminium enolates and alkoxides derived from trimethylaluminium and aromatic ketones: a synthetic, structural and theoretical investigation, 15

## Erbium

Zur Elektronenstruktur metallorganischer Komplexe der f-Elemente IL. Erstmalige Aufklärung der Elektronenstruktur eines metallorganischen σ-Komplexes der f-Elemente: Tris(trimethylsilyl)methyl)erbium(III), 65

## Ferrocene

The behavior of 3-ferrocenyl-1-methyl-1,2-pentamethyleneallyl and 1,5-diferrocenyl-3-methyl-2,4-tetramethylene-1,4-dienyl carbocations in the cationic dimerization of 1,3-dienes, 105

## Gallium

Precursor chemistry of Group III nitrides: Part XVI. Synthesis and structure of monomeric penta-coordinated intramolecularly adduct-stabilized amidobisazides of aluminum, gallium and indium with an all-nitrogen coordination sphere: OMCVD of GaN using (N<sub>3</sub>)<sub>2</sub>Ga{N[CH<sub>2</sub>CH<sub>2</sub>(NET<sub>2</sub>)<sub>2</sub>]}, 29

## Gallium nitride

Precursor chemistry of Group III nitrides: Part XVI. Synthesis and structure of monomeric penta-coordinated intramolecularly adduct-stabilized amidobisazides of aluminum, gallium and indium with an all-nitrogen coordination sphere: OMCVD of GaN using (N<sub>3</sub>)<sub>2</sub>Ga{N[CH<sub>2</sub>CH<sub>2</sub>(NET<sub>2</sub>)<sub>2</sub>]}, 29

## Gold(I)

Aurophilic association in *endo*-dicyclopentadienechlorogold(I), 133

## Halogenation

Halogenation and oxygenation of the metal and oxidation of phosphorus in complexes of the type M(dppe)<sub>2</sub>(N<sub>2</sub>)<sub>2</sub> (M = Mo, W) by reaction with thionylhalides, 24

## Heterocycles

New zwitterionic heterocycles derived from 1-chloro-(dimethyl)silyl-1-borylalkenes and 2-lithio-1-methylimidazole and 1-lithio-indazoles, 45

## Homogeneous catalysis

Catalytic activity of cationic diphospalladium(II) complexes in the alkene/CO copolymerization in organic solvents and water in dependence on the length of the alkyl chain at the phosphine ligands, 173

## Hydrido complexes

Reactivity of dinuclear arene ruthenium complexes: reactions of the hydrido complex  $[(p\text{-Me-C}_6\text{H}_4\text{-Pr})_2\text{Ru}_2\text{Cl}_2(\mu\text{-Cl})(\mu\text{-H})]$  with NaX and HX (X = F, Cl, Br, I), 188

## Hydroboration

New zwitterionic heterocycles derived from 1-chloro-(dimethyl)silyl-1-borylalkenes and 2-lithio-1-methylimidazole and 1-lithio-indazole, 45

## Hydrogen bonding

Crystal and molecular structure of di-*tert*-butylhydridosilanol, 170

## Hydroxymethyl phosphines

Synthesis and reactions of the ferrocene derived hydroxymethyl phosphine  $\text{FcCH}(\text{CH}_3)\text{P}(\text{CH}_2\text{OH})_2$  and its sulfide: crystal structures of  $[\text{FcCH}(\text{CH}_3)\text{P}(\text{S})\text{R}_2]$ , R = CH<sub>2</sub>OH, CH<sub>2</sub>CH<sub>2</sub>CN and  $\text{FcCH}(\text{CH}_3)\text{P}(\text{S})(\text{CH}_2\text{O})_2\text{PPh}$  (Fc = ferrocenyl), 125

## Imidazole NMR multinuclear

New zwitterionic heterocycles derived from 1-chloro-(dimethyl)silyl-1-borylalkenes and 2-lithio-1-methylimidazole and 1-lithio-indazole, 45

## Indium

Precursor chemistry of Group III nitrides: Part XVI. Synthesis and structure of monomeric penta-coordinated intramolecularly adduct-stabilized amidobisazides of aluminum, gallium and indium with an all-nitrogen coordination sphere: OMCDV of GaN using  $(\text{N}_3)_2\text{Ga}\{\text{N}[\text{CH}_2\text{CH}_2(\text{NEt}_2)]_2\}$ , 29

## Infrared spectroscopy

Auropophilic association in *endo*-dicyclopentadienechlorogold(I), 133

Copper(I) complexes with conjugated dienes, 5

## Insertion

$\pi$ -Allyliridium(I) complexes,  $\text{Ir}(\eta^3\text{-CH}_2\text{CHCHAR})(\text{CO})(\text{PPh}_3)_2$  (Ar = Ph, C<sub>6</sub>H<sub>4</sub>Me-*p*, C<sub>6</sub>H<sub>4</sub>Br-*p*). Comparison of their structures and chemical properties with analogous Rh complexes, 144

## Intramolecular adduct

Precursor chemistry of Group III nitrides: Part XVI. Synthesis and structure of monomeric penta-coordinated intramolecularly adduct-stabilized amidobisazides of aluminum, gallium and indium with an all-nitrogen coordination sphere: OMCDV of GaN using  $(\text{N}_3)_2\text{Ga}\{\text{N}[\text{CH}_2\text{CH}_2(\text{NEt}_2)]_2\}$ , 29

## Iridium

$\pi$ -Allyliridium(I) complexes,  $\text{Ir}(\eta^3\text{-CH}_2\text{CHCHAR})(\text{CO})(\text{PPh}_3)_2$  (Ar = Ph, C<sub>6</sub>H<sub>4</sub>Me-*p*, C<sub>6</sub>H<sub>4</sub>Br-*p*). Comparison of their structures and chemical properties with analogous Rh complexes, 144

Iridium carbonyl cluster complexes with bidentate phosphine ligands; the X-ray crystal structure of  $[\text{Ir}_4(\text{CO})_8(\text{dppa})_2] \cdot 3(\text{THF})$ , 137

## Isocyanide complexes

Use of the aqua complex  $[\text{fac-Mn}(\text{CO})_3(\text{dppe})\text{OH}_2]\text{BF}_4$  for the preparation of a series of *fac*-Mn(CO)<sub>3</sub>(dppe)Z and *[fac-Mn(CO)<sub>3</sub>(dppe)Z]BF<sub>4</sub>* complexes. X-ray crystal structures of three representative examples, 97

## Isoprene

Copper(I) complexes with conjugated dienes, 5

## Isopropenylacetylene

Copper(I) complexes with conjugated dienes, 5

## Kinetics

Rh(acac)(CO)(PR<sub>3</sub>) and Rh(oxinate)(CO)(PR<sub>3</sub>) complexes—substitution chemistry and structural aspects, 59

## Ligand properties

Water-soluble phosphines: Part XII. Pd catalyzed P–C coupling reactions: a novel synthetic route to cationic phosphines with *para*- and *meta*-guanidiniumphenyl moieties, 158

## Ligand substitution

Rh(acac)(CO)(PR<sub>3</sub>) and Rh(oxinate)(CO)(PR<sub>3</sub>) complexes—substitution chemistry and structural aspects, 59

## Linear dichroism spectrum

Zur Elektronenstruktur metallorganischer Komplexe der f-Elemente IL. Erstmalige Aufklärung der Elektronenstruktur eines metallorganischen  $\sigma$ -Komplexes der f-Elemente: Tris(bis(trimethylsilyl)methyl)erbium(III), 65

## Manganese complexes

Use of the aqua complex  $[\text{fac-Mn}(\text{CO})_3(\text{dppe})\text{OH}_2]\text{BF}_4$  for the preparation of a series of *fac*-Mn(CO)<sub>3</sub>(dppe)Z and *[fac-Mn(CO)<sub>3</sub>(dppe)Z]BF<sub>4</sub>* complexes. X-ray crystal structures of three representative examples, 97

## Mercury(II)

Heterobimetallische Komplexe von Titan(IV)–Quecksilber(II) und Platin(II)–Quecksilber(II), 91  
*meta*-, *para*-Guanidiniumphenyl phosphines

Water-soluble phosphines: Part XII. Pd catalyzed P–C coupling reactions: a novel synthetic route to cationic phosphines with *para*- and *meta*-guanidiniumphenyl moieties, 158

## Molecular orbital schemes

Zur Elektronenstruktur metallorganischer Komplexe der f-Elemente IL. Erstmalige Aufklärung der Elektronenstruktur eines metallorganischen  $\sigma$ -Komplexes der f-Elemente: Tris(bis(trimethylsilyl)methyl)erbium(III), 65

## Molybdenum

Halogenation and oxygenation of the metal and oxidation of phosphorus in complexes of the type M(dppe)<sub>2</sub>(N<sub>2</sub>)<sub>2</sub> (M = Mo, W) by reaction with thionylhalides, 24

## Molybdenum dioxide

A mononuclear alkyne-stabilized Mo<sup>(IV)</sup>O<sub>2</sub> moiety, 72

## Molybdenum(II)

*Cis*- and *trans*-bis(diphenylphosphino)ethene seven-coordinate complexes of molybdenum(II) and tungsten(II). Crystal structures of  $[\text{MI}_2(\text{CO})_3(\text{cis-dppen})]$  {M = Mo or W; dppe = bis(diphenylphosphino)ethene},  $[\text{WI}(\text{CO})_2(\text{cis-dppen})_2]\text{J}$ ,  $[\text{MI}_2(\text{CO})_2\{\text{P}(\text{OR})_3\}(\text{cis-dppen})]$  (M = Mo, R = Me; M = W, R = Et or Ph), 115

## Monocyclopentadienyl

Monocyclopentadienyl and mono(2-methoxyethyl)cyclopentadienyl zirconium methyl complexes, 51

## Mono zirconium complexes

Monocyclopentadienyl and mono(2-methoxyethyl)cyclopentadienyl zirconium methyl complexes, 51

## Nitrato complexes

Use of the aqua complex  $[\text{fac-Mn}(\text{CO})_3(\text{dppe})\text{OH}_2]\text{BF}_4$  for the preparation of a series of *fac*-Mn(CO)<sub>3</sub>(dppe)Z and *[fac-Mn(CO)<sub>3</sub>(dppe)Z]BF<sub>4</sub>* complexes. X-ray crystal structures of three representative examples, 97

## Nitrile complexes

Use of the aqua complex  $[\text{fac-Mn}(\text{CO})_3(\text{dppe})\text{OH}_2]\text{BF}_4$  for the preparation of a series of *fac*-Mn(CO)<sub>3</sub>(dppe)Z and *[fac-Mn(CO)<sub>3</sub>(dppe)Z]BF<sub>4</sub>* complexes. X-ray crystal structures of three representative examples, 97

## NMR spectroscopy

Electrochemistry of ruthenium metallocenes: Part 3. Synthesis and properties of ruthenium [2<sub>n</sub>]paracyclophane complexes with methacrylic acid and methacrylate ester substituents, 75

## Oligosilane

Formation and characterization of cyclic and polycyclic silthianes containing Si–Si bonds, 193

## OMCVD

Precursor chemistry of Group III nitrides: Part XVI. Synthesis and structure of monomeric penta-coordinated intramolecularly adduct-stabilized amidobisazides of aluminum, gallium and indium with an all-nitrogen coordination sphere: OMCVD of GaN using  $(\text{N}_3)_2\text{Ga}\{\text{N}[\text{CH}_2\text{CH}_2(\text{NEt}_2)]_2\}$ , 29

Organometallic  $\pi$ -tweezers

Heterobimetallic Komplexe von Titan(IV)–Quecksilber(II) und Platin(II)–Quecksilber(II), 91

## Orthometallated

Reactivity of Pd(II) complexes containing the orthometallated C,C-chelating ligand  $C_6\text{H}_4\text{-}2\text{-PPh}_2\text{C}(\text{H})\text{COCH}_2\text{PPh}_3$  towards deprotonating reagents. Part 2, 151

## Oxygenation

Halogenation and oxygenation of the metal and oxidation of phosphorus in complexes of the type  $\text{M}(\text{dppe})_2(\text{N}_2)_2$  ( $\text{M} = \text{Mo, W}$ ) by reaction with thionylhalides, 24

## Palladium

Reactivity of Pd(II) complexes containing the orthometallated C,C-chelating ligand  $C_6\text{H}_4\text{-}2\text{-PPh}_2\text{C}(\text{H})\text{COCH}_2\text{PPh}_3$  towards deprotonating reagents. Part 2, 151

## Palladium complexes

Catalytic activity of cationic diphospalladium(II) complexes in the alkene/CO copolymerization in organic solvents and water in dependence on the length of the alkyl chain at the phosphine ligands, 173

## Pd catalyzed P–C coupling

Water-soluble phosphines: Part XII. Pd catalyzed P–C coupling reactions: a novel synthetic route to cationic phosphines with *para*- and *meta*-guanidiniumphenyl moieties, 158

## 1,3-Pentadiene

Copper(I) complexes with conjugated dienes, 5

## Phosphine sulfide

Synthesis and reactions of the ferrocene derived hydroxymethyl phosphine  $\text{FcCH}(\text{CH}_3)\text{P}(\text{CH}_2\text{OH})_2$  and its sulfide: crystal structures of  $[\text{FcCH}(\text{CH}_3)\text{P}(\text{S})\text{R}_2]$   $\text{R} = \text{CH}_2\text{OH, CH}_2\text{CH}_2\text{CN}$  and  $\text{FcCH}(\text{CH}_3)\text{P}(\text{S})(\text{CH}_2\text{O})_2\text{PPh}$  ( $\text{Fc} = \text{ferrocenyl}$ ), 125

## Phosphoylides

Reactivity of Pd(II) complexes containing the orthometallated C,C-chelating ligand  $C_6\text{H}_4\text{-}2\text{-PPh}_2\text{C}(\text{H})\text{COCH}_2\text{PPh}_3$  towards deprotonating reagents. Part 2, 151

## Platinum(II)

Heterobimetallic Komplexe von Titan(IV)–Quecksilber(II) und Platin(II)–Quecksilber(II), 91

## Polyketones

Catalytic activity of cationic diphospalladium(II) complexes in the alkene/CO copolymerization in organic solvents and water in dependence on the length of the alkyl chain at the phosphine ligands, 173

## Racemic phosphine

Synthesis and reactions of the ferrocene derived hydroxymethyl phosphine  $\text{FcCH}(\text{CH}_3)\text{P}(\text{CH}_2\text{OH})_2$  and its sulfide: crystal structures of  $[\text{FcCH}(\text{CH}_3)\text{P}(\text{S})\text{R}_2]$   $\text{R} = \text{CH}_2\text{OH, CH}_2\text{CH}_2\text{CN}$  and  $\text{FcCH}(\text{CH}_3)\text{P}(\text{S})(\text{CH}_2\text{O})_2\text{PPh}$  ( $\text{Fc} = \text{ferrocenyl}$ ), 125

## Reduction

The behavior of 3-ferrocenyl-1-methyl-1,2-pentamethyleneallyl and 1,5-diferrocenyl-3-methyl-2,4-tetramethylene-1,4-dienyl carbocations in the cationic dimerization of 1,3-dienes, 105

## Rhodium

$\text{Rh}(\text{acac})(\text{CO})(\text{PR}_3)$  and  $\text{Rh}(\text{oxinate})(\text{CO})(\text{PR}_3)$  complexes—substitution chemistry and structural aspects, 59

## Ruthenium complexes

Electrochemistry of ruthenium metallocenes: Part 3. Synthesis and properties of ruthenium [2<sub>2</sub>]paracyclophane complexes with methacrylic acid and methacrylate ester substituents, 75

## Ruthenium complexes

Reactivity of dinuclear arene ruthenium complexes: reactions of the hydrido complex  $[(p\text{-Me-C}_6\text{H}_4\text{-Pr})_2\text{Ru}_2\text{Cl}_2(\mu\text{-Cl})(\mu\text{-H})]$  with NaX and HX ( $X = \text{F, Cl, Br, I}$ ), 188

## Seven-coordinate

*Cis*- and *trans*-bis(diphenylphosphino)ethene seven-coordinate complexes of molybdenum(II) and tungsten(II). Crystal structures of  $[\text{MI}_2(\text{CO})_3\{\text{cis-dppen}\}]$  ( $\text{M} = \text{Mo or W}$ ; dppen = bis(diphenylphosphino)ethene),  $[\text{WI}(\text{CO})_2\{\text{cis-dppen}\}_2]\text{I}$ ,  $[\text{MI}_2(\text{CO})_2\{\text{P}(\text{OR})_3\}\{\text{cis-dppen}\}]$  ( $\text{M} = \text{Mo, R = Me; M = W, R = Et or Ph}$ ), 115

## Silane

Formation and characterization of cyclic and polycyclic silthianes containing Si–Si bonds, 193

## Silane dendrimers

Silane dendrimers containing titanium complexes on their periphery, 208

## Silanes

New zwitterionic heterocycles derived from 1-chloro-(dimethyl)silyl-1-borylalkenes and 2-lithio-1-methylimidazole and 1-lithio-indazole, 45

## Silanol

Crystal and molecular structure of di-*tert*-butylhydridosilanol, 170

## Silicon

Crystal and molecular structure of di-*tert*-butylhydridosilanol, 170

## Silthiane

Formation and characterization of cyclic and polycyclic silthianes containing Si–Si bonds, 193

<sup>29</sup>Si-NMR

Formation and characterization of cyclic and polycyclic silthianes containing Si–Si bonds, 193

## Spirocycles

Organotransition metal-modified sugars: Part 16. Diastereoselective synthesis of carbohydrate-derived spirocyclic 2-oxacyclohexylidene chromium and tungsten complexes, 37

## Suzuki coupling

Water-soluble phosphines: Part XII. Pd catalyzed P–C coupling reactions: a novel synthetic route to cationic phosphines with *para*- and *meta*-guanidiniumphenyl moieties, 158

## Tertiary phosphines

$\text{Rh}(\text{acac})(\text{CO})(\text{PR}_3)$  and  $\text{Rh}(\text{oxinate})(\text{CO})(\text{PR}_3)$  complexes—substitution chemistry and structural aspects, 59

## Thionylhalides

Halogenation and oxygenation of the metal and oxidation of phosphorus in complexes of the type  $\text{M}(\text{dppe})_2(\text{N}_2)_2$  ( $\text{M} = \text{Mo, W}$ ) by reaction with thionylhalides, 24

## Titanium complexes

Silane dendrimers containing titanium complexes on their periphery, 208

## Titanium(IV)

Heterobimetallic Komplexe von Titan(IV)–Quecksilber(II) und Platin(II)–Quecksilber(II), 91

## Titanocenes

A mononuclear alkyne-stabilized  $\text{Mo}^{(\text{IV})}\text{O}_2$  moiety, 72

## Tungsten

Halogenation and oxygenation of the metal and oxidation of phosphorus in complexes of the type  $\text{M}(\text{dppe})_2(\text{N}_2)_2$  ( $\text{M} = \text{Mo, W}$ ) by reaction with thionylhalides, 24

## Tungsten(II)

*Cis-* and *trans*-bis(diphenylphosphino)ethene seven-coordinate complexes of molybdenum(II) and tungsten(II). Crystal structures of  $[M\text{I}_2(\text{CO})_3\{\text{cis}\text{-dppen}\}]$  { $M = \text{Mo}$  or  $\text{W}$ ; dppen = bis(diphenylphosphino)ethene},  $[\text{WI}(\text{CO})_2\{\text{cis}\text{-dppen}\}_2]\text{I}$ ,  $[\text{MI}_2(\text{CO})_2\{\text{P}(\text{OR})_3\}\{\text{cis}\text{-dppen}\}]$  ( $M = \text{Mo}$ ,  $R = \text{Me}$ ;  $M = \text{W}$ ,  $R = \text{Et}$  or  $\text{Ph}$ ), 115

## Water solubility

Catalytic activity of cationic diphospalladium(II) complexes in the alkene/CO copolymerization in organic solvents and water in dependence on the length of the alkyl chain at the phosphine ligands, 173

## X-ray

New zwitterionic heterocycles derived from 1-chloro-(dimethyl)silyl-1-borylalkenes and 2-lithio-1-methylimidazole and 1-lithio-indazole, 45

## X-ray structure

Crystal and molecular structure of di-*tert*-butylhydridosilanol, 170

Formation and characterization of cyclic and polycyclic silthianes containing Si–Si bonds, 193

Water-soluble phosphines: Part XII. Pd catalyzed P–C coupling reactions: a novel synthetic route to cationic phosphines with *para*- and *meta*-guanidiniumphenyl moieties, 158