



ELSEVIER

www.elsevier.nl/locate/jorgchem

Journal of Organometallic Chemistry 612 (2000) 187–190

Journal  
of Organo  
metallic  
Chemistry

## Subject Index of Volume 612

### Adduct

Triphenylphosphine oxide adducts of diphenylantimony(V) and diorganotin(IV) Lewis acids: structures of  $\text{SnPh}_2\text{Cl}_2 \cdot \text{OPPh}_3$ ,  $\text{SnPh}_2\text{Cl}_2 \cdot 2\text{OPPh}_3$ ,  $\text{SnPh}_2\text{Br}_2 \cdot \text{OPPh}_3$  and  $\text{SbPh}_2\text{Cl}_3 \cdot \text{OPPh}_3$ , 53

### Alkyltin

Wurtz-type reductive coupling reaction of primary alkyl iodides and haloorganotins in cosolvent/ $\text{H}_2\text{O}(\text{NH}_4\text{Cl})/\text{Zn}$  media as a route to mixed alkylstannanes and hexaalkyldistannanes, 78

### Alkyne

The photolysis of  $\text{Fe}(\text{CO})_3\{\text{P}(\text{OPh})_3\}_2$  and reaction of the photoproduct with alkynes: crystal structures of  $\text{Fe}(\text{CO})_2\{\text{P}(\text{OPh})_3\}_2(\eta^2\text{-PhCCPh})$ ,  $\text{Fe}(\text{CO})_2\{\text{P}(\text{OPh})_3\}_2(\eta^1:\eta^1\text{-C(O)C(Me)C[CH(OEt)]_2C(O)})$  and  $\text{Fe}_2(\text{CO})_4\{\text{P}(\text{OPh})_3\}_2(\text{PhCCH}_2)$ , 61

### Bis{2-(*N*-morpholino)-ethyl} telluride

$N\text{-}\{2\text{-}(4\text{-Methoxyphenyltelluro)ethyl}\}\text{morpholine (L}^1\text{)}$  and bis{2-(*N*-morpholino)ethyl} telluride ( $\text{L}^2$ ): synthesis and complexation with palladium(II) and mercury(II). Crystal structures of *trans*-[ $\text{PdCl}_2(\text{L}^1)_2$ ] and *trans*-[ $\text{PdCl}_2(\text{L}^2)_2$ ], 46

### 1,1'-bis(Butylstannyl)ferrocene

A novel synthesis of 1,1'-unsymmetrical disubstituted heteroaryl-ferrocene, 172

### Carbenoid

*ortho*-Metalated dirhodium(II)-catalyzed  $\alpha$ -diazocarbonyl transformation. Diastereoselective cyclopropanation of menthyl- $\alpha$ -diazo- $\beta$ -keto ester and C–H insertion of  $\alpha$ -diazo ester, 36

### Carbonyl

Reactivities of  $\pi$ -electron-rich phosphorus–nitrogen ligands toward Groups VIB and VIIB metal carbonyl complexes, 160

### Carbonyl compounds

Unbridged homo and hetero dinuclear complexes of Group 6 and 8 metals: synthesis, characterization and comparison of X-ray crystallographic data, 106

### Carbonyl ligands

Transition metal complexes of 2,2,2-trifluoroethyl isocyanide and 1*H*,1*H*-perfluoroctyl isocyanide, 96

### Chalcogenides

Antiferromagnetic complexes with metal–metal bonds: Part XXIX. Synthesis and molecular structures of heterochalcogenide binuclear complex [ $\pi\text{-}(\text{CH}_3\text{C}_5\text{H}_4)\text{Cr}(\mu\text{-SPh})_2\text{Se}$  and trinuclear mixed-metal cluster [ $\pi\text{-}(\text{CH}_3\text{C}_5\text{H}_4)_2\text{Cr}_2(\mu\text{-SPh})(\mu_3\text{-S})(\mu_3\text{-Se})\text{Co}(\text{CO})_2$ ], 9

### Chromium

Antiferromagnetic complexes with metal–metal bonds: Part XXIX. Synthesis and molecular structures of heterochalcogenide binuclear complex [ $\pi\text{-}(\text{CH}_3\text{C}_5\text{H}_4)\text{Cr}(\mu\text{-SPh})_2\text{Se}$  and trinuclear mixed-metal cluster [ $\pi\text{-}(\text{CH}_3\text{C}_5\text{H}_4)_2\text{Cr}_2(\mu\text{-SPh})(\mu_3\text{-S})(\mu_3\text{-Se})\text{Co}(\text{CO})_2$ ], 9

Reactivities of  $\pi$ -electron-rich phosphorus–nitrogen ligands toward Groups VIB and VIIB metal carbonyl complexes, 160

### Cobalt

Antiferromagnetic complexes with metal–metal bonds: Part XXIX. Synthesis and molecular structures of heterochalcogenide binuclear complex [ $\pi\text{-}(\text{CH}_3\text{C}_5\text{H}_4)\text{Cr}(\mu\text{-SPh})_2\text{Se}$  and trinuclear mixed-metal cluster [ $\pi\text{-}(\text{CH}_3\text{C}_5\text{H}_4)_2\text{Cr}_2(\mu\text{-SPh})(\mu_3\text{-S})(\mu_3\text{-Se})\text{Co}(\text{CO})_2$ ], 9

### Complex

$N\text{-}\{2\text{-}(4\text{-Methoxyphenyltelluro)ethyl}\}\text{morpholine (L}^1\text{)}$  and bis{2-(*N*-morpholino)ethyl} telluride ( $\text{L}^2$ ): synthesis and complexation with palladium(II) and mercury(II). Crystal structures of *trans*-[ $\text{PdCl}_2(\text{L}^1)_2$ ] and *trans*-[ $\text{PdCl}_2(\text{L}^2)_2$ ], 46

### Coupling

Wurtz-type reductive coupling reaction of primary alkyl iodides and haloorganotins in cosolvent/ $\text{H}_2\text{O}(\text{NH}_4\text{Cl})/\text{Zn}$  media as a route to mixed alkylstannanes and hexaalkyldistannanes, 78

### Crystal structures

$N\text{-}\{2\text{-}(4\text{-Methoxyphenyltelluro)ethyl}\}\text{morpholine (L}^1\text{)}$  and bis{2-(*N*-morpholino)ethyl} telluride ( $\text{L}^2$ ): synthesis and complexation with palladium(II) and mercury(II). Crystal structures of *trans*-[ $\text{PdCl}_2(\text{L}^1)_2$ ] and *trans*-[ $\text{PdCl}_2(\text{L}^2)_2$ ], 46

Synthesis and studies on reactivities of tetrahydrofurfurylcyclopentadienyl lanthanide chlorides  $\text{Cp}^\delta\text{LnCl}_2\text{·}2\text{THF}$  ( $\text{Ln} = \text{Dy}, \text{Ho}, \text{Gd}$ ) and  $\text{Cp}_2^\delta\text{LnCl}\text{·}(\text{Ln} = \text{Sm}, \text{Dy})$ . The crystal structures of  $\text{Cp}_2^\delta\text{SmCl}$  and  $\text{Cp}_2^\delta\text{Sm}\text{·}(\text{C}_3\text{N}_2\text{HMe}_2)$ , 176

Unbridged homo and hetero dinuclear complexes of Group 6 and 8 metals: synthesis, characterization and comparison of X-ray crystallographic data, 106

### Cyclometallation

Polymeric cyclometallated palladium (II) complexes derived from potentially hexadentate Schiff base ligands. Crystal structures of  $[(\text{Cl})\text{Pd}\{\text{Me}_2\text{NCH}_2\text{CH}_2\text{N}=\text{H}\text{C}\}\text{C}_6\text{H}_2\{\text{C}(\text{H})=\text{NCH}_2\text{CH}_2\text{NMe}_2\}\text{Pd}(\text{Cl})]$  and  $[(\{\text{Ph}_2\text{PCH}_2\text{CH}_2\}_2\text{PPh-P}, P, P)\text{Pd}\{\text{Me}_2\text{NCH}_2\text{CH}_2\text{N}=\text{H}\text{C}\}\text{C}_6\text{H}_2\{\text{C}(\text{H})=\text{NCH}_2\text{CH}_2\text{NMe}_2\}\text{Pd}\{\text{Ph}_2\text{PCH}_2\text{CH}_2\}_2\text{PPh-P}, P, P]\}][\text{Cl}]_2$ , 85

### Cyclopentadienyl complexes

Antiferromagnetic complexes with metal–metal bonds: Part XXIX. Synthesis and molecular structures of heterochalcogenide binuclear complex [ $\pi\text{-}(\text{CH}_3\text{C}_5\text{H}_4)\text{Cr}(\mu\text{-SPh})_2\text{Se}$  and trinuclear mixed-metal cluster [ $\pi\text{-}(\text{CH}_3\text{C}_5\text{H}_4)_2\text{Cr}_2(\mu\text{-SPh})(\mu_3\text{-S})(\mu_3\text{-Se})\text{Co}(\text{CO})_2$ ], 9

### DFT calculations

Vibrational spectroscopy studies and density functional theory calculations on square-planar vinylidene, carbonyl and ethylene rhodium(I) complexes, 125

### Diazo compounds

*ortho*-Metalated dirhodium(II)-catalyzed  $\alpha$ -diazocarbonyl transformation. Diastereoselective cyclopropanation of menthyl- $\alpha$ -diazo- $\beta$ -keto ester and C–H insertion of  $\alpha$ -diazo ester, 36

### $\mu$ -Dimethylaminocarbonyne

The reactions of *cis*-[ $\text{Fe}_2(\eta\text{-C}_5\text{H}_5)_2(\text{CO})_2(\mu\text{-CNMe}_2)_2\}[\text{SO}_3\text{CF}_3]_2$  and related compounds with nucleophiles, 18

## Dinuclear complexes

Polymeric cyclometallated palladium (II) complexes derived from potentially hexadentate Schiff base ligands. Crystal structures of  $[(Cl)Pd\{Me_2NCH_2CH_2N(H)C\}C_6H_2\{C(H)=NCH_2CH_2NMe_2\}Pd(Cl)]$  and  $[(Ph_2PCH_2CH_2)_2PPh-P,P;Pd\{Me_2NCH_2CH_2N(H)C\}C_6H_2\{C(H)=NCH_2NMe_2\}Pd\{(Ph_2PCH_2CH_2)_2PPh-P,P\}][Cl]_2$ , 85

## Disilanes

1,2-Dithiolate derivatives of monosilanes and disilanes, 133

## Distannane

Wurtz-type reductive coupling reaction of primary alkyl iodides and haloorganotins in cosolvent/H<sub>2</sub>O(NH<sub>4</sub>Cl)/Zn media as a route to mixed alkylstannanes and hexaalkyldistannanes, 78

## Disupersilyldisilanes

Disupersilyldisilane R\*X<sub>2</sub>Si–SiX<sub>2</sub>R\*: Darstellung, Charakterisierung und Strukturen; sterische Effekte der Substituenten, 141

## Ethyne

<sup>57</sup>Fe-Mössbauer spectroscopic study of monomeric and polymeric ferrocenyl- and octamethylferrocenyl-substituted ethynes, 1

## Ferrocenyl polymers

<sup>57</sup>Fe-Mössbauer spectroscopic study of monomeric and polymeric ferrocenyl- and octamethylferrocenyl-substituted ethynes, 1

## Ferrole

The photolysis of Fe(CO)<sub>3</sub>{P(OPh)<sub>3</sub>}<sub>2</sub> and reaction of the photoproduct with alkynes: crystal structures of Fe(CO)<sub>2</sub>{P(OPh)<sub>3</sub>}<sub>2</sub>(η<sup>2</sup>-PhCCPh), Fe(CO)<sub>2</sub>{P(OPh)<sub>3</sub>}<sub>2</sub>[η<sup>1</sup>:η<sup>1</sup>-C(Me)C[CH(OEt)<sub>2</sub>]C(O)} and Fe<sub>2</sub>(CO)<sub>4</sub>{P(OPh)<sub>3</sub>}<sub>2</sub>(PhCCH)<sub>2</sub>, 61

## Hetero bimetallic complexes

Unbridged homo and hetero dinuclear complexes of Group 6 and 8 metals: synthesis, characterization and comparison of X-ray crystallographic data, 106

## IR and FT-Raman spectroscopy

Vibrational spectroscopy studies and density functional theory calculations on square-planar vinylidene, carbonyl and ethylene rhodium(I) complexes, 125

## Iron

The reactions of cis-[Fe<sub>2</sub>(η-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>(CO)<sub>2</sub>(μ-CNMe<sub>2</sub>)<sub>2</sub>][SO<sub>3</sub>CF<sub>3</sub>]<sub>2</sub> and related compounds with nucleophiles, 18

## Iron carbonyl

The photolysis of Fe(CO)<sub>3</sub>{P(OPh)<sub>3</sub>}<sub>2</sub> and reaction of the photoproduct with alkynes: crystal structures of Fe(CO)<sub>2</sub>{P(OPh)<sub>3</sub>}<sub>2</sub>(η<sup>2</sup>-PhCCPh), Fe(CO)<sub>2</sub>{P(OPh)<sub>3</sub>}<sub>2</sub>[η<sup>1</sup>:η<sup>1</sup>-C(Me)C[CH(OEt)<sub>2</sub>]C(O)} and Fe<sub>2</sub>(CO)<sub>4</sub>{P(OPh)<sub>3</sub>}<sub>2</sub>(PhCCH)<sub>2</sub>, 61

## Isocyanide ligands

Transition metal complexes of 2,2,2-trifluoroethyl isocyanide and 1H,1H-perfluorooctyl isocyanide, 96

## Isotopic substitution

Vibrational spectroscopy studies and density functional theory calculations on square-planar vinylidene, carbonyl and ethylene rhodium(I) complexes, 125

## Magnetic properties

Antiferromagnetic complexes with metal–metal bonds: Part XXIX. Synthesis and molecular structures of heterochalcogenide binuclear complex [π-(CH<sub>3</sub>C<sub>5</sub>H<sub>4</sub>)Cr(μ-SPh)]<sub>2</sub>Se and trinuclear mixed-metal cluster [π-(CH<sub>3</sub>C<sub>5</sub>H<sub>4</sub>)<sub>2</sub>Cr<sub>2</sub>(μ-SPh)][μ<sub>3</sub>-S](μ<sub>3</sub>-Se)Co(CO)<sub>2</sub>, 9

## Maleoyl

The photolysis of Fe(CO)<sub>3</sub>{P(OPh)<sub>3</sub>}<sub>2</sub> and reaction of the photoproduct with alkynes: crystal structures of Fe(CO)<sub>2</sub>{P(OPh)<sub>3</sub>}<sub>2</sub>(η<sup>2</sup>-PhCCPh), Fe(CO)<sub>2</sub>{P(OPh)<sub>3</sub>}<sub>2</sub>[η<sup>1</sup>:η<sup>1</sup>-C(Me)C[CH(OEt)<sub>2</sub>]C(O)} and Fe<sub>2</sub>(CO)<sub>4</sub>{P(OPh)<sub>3</sub>}<sub>2</sub>(PhCCH)<sub>2</sub>, 61

## Manganese

Reactivities of π-electron-rich phosphorus–nitrogen ligands toward Groups VIB and VIIB metal carbonyl complexes, 160

## Mercury(II)

N-{2-(4-Methoxyphenyltelluro)ethyl}morpholine (**L**<sup>1</sup>) and bis{2-(N-morpholino)ethyl}telluride (**L**<sup>2</sup>): synthesis and complexation with palladium(II) and mercury(II). Crystal structures of trans-[PdCl<sub>2</sub>(**L**<sup>1</sup>)<sub>2</sub>] and trans-[PdCl<sub>2</sub>(**L**<sup>2</sup>)<sub>2</sub>], 46

## Metal carbonyls

Antiferromagnetic complexes with metal–metal bonds: Part XXIX. Synthesis and molecular structures of heterochalcogenide binuclear complex [π-(CH<sub>3</sub>C<sub>5</sub>H<sub>4</sub>)Cr(μ-SPh)]<sub>2</sub>Se and trinuclear mixed-metal cluster [π-(CH<sub>3</sub>C<sub>5</sub>H<sub>4</sub>)<sub>2</sub>Cr<sub>2</sub>(μ-SPh)][μ<sub>3</sub>-S](μ<sub>3</sub>-Se)Co(CO)<sub>2</sub>, 9

## Mixed-metal clusters

Antiferromagnetic complexes with metal–metal bonds: Part XXIX. Synthesis and molecular structures of heterochalcogenide binuclear complex [π-(CH<sub>3</sub>C<sub>5</sub>H<sub>4</sub>)Cr(μ-SPh)]<sub>2</sub>Se and trinuclear mixed-metal cluster [π-(CH<sub>3</sub>C<sub>5</sub>H<sub>4</sub>)<sub>2</sub>Cr<sub>2</sub>(μ-SPh)][μ<sub>3</sub>-S](μ<sub>3</sub>-Se)Co(CO)<sub>2</sub>, 9

## Molybdenum

Reactivities of π-electron-rich phosphorus–nitrogen ligands toward Groups VIB and VIIB metal carbonyl complexes, 160

Unbridged homo and hetero dinuclear complexes of Group 6 and 8 metals: synthesis, characterization and comparison of X-ray crystallographic data, 106

## Monosilanes

1,2-Dithiolate derivatives of monosilanes and disilanes, 133

## Mössbauer

Triphenylphosphine oxide adducts of diphenylantimony(V) and diorganotin(IV) Lewis acids: structures of SnPh<sub>2</sub>Cl<sub>2</sub>·OPPh<sub>3</sub>, SnPh<sub>2</sub>Cl<sub>2</sub>·2OPPh<sub>3</sub>, SnPh<sub>2</sub>Br<sub>2</sub>·OPPh<sub>3</sub> and SbPh<sub>2</sub>Cl<sub>3</sub>·OPPh<sub>3</sub>, 53

## Mössbauer spectroscopy

<sup>57</sup>Fe-Mössbauer spectroscopic study of monomeric and polymeric ferrocenyl- and octamethylferrocenyl-substituted ethynes, 1

## N-{2-(4-Methoxyphenyltelluro)ethyl}morpholine

N-{2-(4-Methoxyphenyltelluro)ethyl}morpholine (**L**<sup>1</sup>) and bis{2-(N-morpholino)ethyl}telluride (**L**<sup>2</sup>): synthesis and complexation with palladium(II) and mercury(II). Crystal structures of trans-[PdCl<sub>2</sub>(**L**<sup>1</sup>)<sub>2</sub>] and trans-[PdCl<sub>2</sub>(**L**<sup>2</sup>)<sub>2</sub>], 46

## NMR spectra

Disupersilyldisilane R\*X<sub>2</sub>Si–SiX<sub>2</sub>R\*: Darstellung, Charakterisierung und Strukturen; sterische Effekte der Substituenten, 141

## Nucleophiles

The reactions of cis-[Fe<sub>2</sub>(η-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>(CO)<sub>2</sub>(μ-CNMe<sub>2</sub>)<sub>2</sub>][SO<sub>3</sub>CF<sub>3</sub>]<sub>2</sub> and related compounds with nucleophiles, 18

## Octamethyl ferrocenyl polymers

<sup>57</sup>Fe-Mössbauer spectroscopic study of monomeric and polymeric ferrocenyl- and octamethylferrocenyl-substituted ethynes, 1

## Organoantimony

Triphenylphosphine oxide adducts of diphenylantimony(V) and diorganotin(IV) Lewis acids: structures of SnPh<sub>2</sub>Cl<sub>2</sub>·OPPh<sub>3</sub>, SnPh<sub>2</sub>Cl<sub>2</sub>·2OPPh<sub>3</sub>, SnPh<sub>2</sub>Br<sub>2</sub>·OPPh<sub>3</sub> and SbPh<sub>2</sub>Cl<sub>3</sub>·OPPh<sub>3</sub>, 53

## Organolanthanide

Synthesis and studies on reactivities of tetrahydrofurfurylclopentadienyl lanthanide chlorides Cp<sup>s</sup>LnCl<sub>2</sub>·2THF (Ln = Dy, Ho, Gd) and Cp<sup>s</sup>LnCl<sub>2</sub> (Ln = Sm, Dy). The crystal structures of Cp<sup>s</sup>SmCl and Cp<sup>s</sup>Sm(C<sub>3</sub>N<sub>2</sub>HMe<sub>2</sub>), 176

## Organotin

Triphenylphosphine oxide adducts of diphenylantimony(V) and diorganotin(IV) Lewis acids: structures of  $\text{SnPh}_2\text{Cl}_2 \cdot \text{OPPh}_3$ ,  $\text{SnPh}_2\text{Cl}_2 \cdot 2\text{OPPh}_3$ ,  $\text{SnPh}_2\text{Br}_2 \cdot \text{OPPh}_3$  and  $\text{SbPh}_2\text{Cl}_3 \cdot \text{OPPh}_3$ , 53

*ortho*-Metalated

*ortho*-Metalated dirhodium(II)-catalyzed  $\alpha$ -diazocarbonyl transformation. Diastereoselective cyclopropanation of menthyl- $\alpha$ -diao- $\beta$ -keto ester and C–H insertion of  $\alpha$ -diao ester, 36

## Orthometallation

The photolysis of  $\text{Fe}(\text{CO})_3\{\text{P}(\text{OPh})_3\}_2$  and reaction of the photo-product with alkynes: crystal structures of  $\text{Fe}(\text{CO})_2\{\text{P}(\text{OPh})_3\}_2(\eta^2\text{-PhCCPh})$ ,  $\text{Fe}(\text{CO})_2\{\text{P}(\text{OPh})_3\}_2(\eta^1\text{:}\eta^1\text{-C(O)-C(Me)C}[\text{CH}(\text{OEt})_2]\text{C(O)})$  and  $\text{Fe}_2(\text{CO})_4\{\text{P}(\text{OPh})_3\}_2(\text{PhC-CH})_2$ , 61

## Oxidative addition

Polynuclear cyclometallated palladium (II) complexes derived from potentially hexadentate Schiff base ligands. Crystal structures of  $[(\text{Cl})\text{Pd}\{\text{Me}_2\text{NCH}_2\text{CH}_2\text{N}=(\text{H})\text{C}\}\text{C}_6\text{H}_2\{\text{C(H)}=\text{NCH}_2\text{CH}_2\text{NMe}_2\}\text{Pd}(\text{Cl})]$  and  $[(\{\text{Ph}_2\text{PCH}_2\text{CH}_2\}_2)\text{PPh-P,P,P}\}\text{Pd}\{\text{Me}_2\text{NCH}_2\text{CH}_2\text{N}=(\text{H})\text{C}\}\text{C}_6\text{H}_2\{\text{C(H)}=\text{NCH}_2\text{CH}_2\text{NMe}_2\}\text{Pd}\{(\text{Ph}_2\text{PCH}_2\text{CH}_2)_2\text{PPh-P,P,P}\}][\text{Cl}]_2$ , 85

## Palladium

Polynuclear cyclometallated palladium (II) complexes derived from potentially hexadentate Schiff base ligands. Crystal structures of  $[(\text{Cl})\text{Pd}\{\text{Me}_2\text{NCH}_2\text{CH}_2\text{N}=(\text{H})\text{C}\}\text{C}_6\text{H}_2\{\text{C(H)}=\text{NCH}_2\text{CH}_2\text{NMe}_2\}\text{Pd}(\text{Cl})]$  and  $[(\{\text{Ph}_2\text{PCH}_2\text{CH}_2\}_2)\text{PPh-P,P,P}\}\text{Pd}\{\text{Me}_2\text{NCH}_2\text{CH}_2\text{N}=(\text{H})\text{C}\}\text{C}_6\text{H}_2\{\text{C(H)}=\text{NCH}_2\text{CH}_2\text{NMe}_2\}\text{Pd}\{(\text{Ph}_2\text{PCH}_2\text{CH}_2)_2\text{PPh-P,P,P}\}][\text{Cl}]_2$ , 85

## Palladium(II)

*N*-{2-(4-Methoxyphenyltelluro)ethyl}morpholine (**L**<sup>1</sup>) and bis{2-(*N*-morpholino)ethyl}telluride (**L**<sup>2</sup>): synthesis and complexation with palladium(II) and mercury(II). Crystal structures of *trans*-[PdCl<sub>2</sub>(**L**<sup>1</sup>)<sub>2</sub>] and *trans*-[PdCl<sub>2</sub>(**L**<sup>2</sup>)<sub>2</sub>], 46

## Phosphine complexes

Polynuclear cyclometallated palladium (II) complexes derived from potentially hexadentate Schiff base ligands. Crystal structures of  $[(\text{Cl})\text{Pd}\{\text{Me}_2\text{NCH}_2\text{CH}_2\text{N}=(\text{H})\text{C}\}\text{C}_6\text{H}_2\{\text{C(H)}=\text{NCH}_2\text{CH}_2\text{NMe}_2\}\text{Pd}(\text{Cl})]$  and  $[(\{\text{Ph}_2\text{PCH}_2\text{CH}_2\}_2)\text{PPh-P,P,P}\}\text{Pd}\{\text{Me}_2\text{NCH}_2\text{CH}_2\text{N}=(\text{H})\text{C}\}\text{C}_6\text{H}_2\{\text{C(H)}=\text{NCH}_2\text{CH}_2\text{NMe}_2\}\text{Pd}\{(\text{Ph}_2\text{PCH}_2\text{CH}_2)_2\text{PPh-P,P,P}\}][\text{Cl}]_2$ , 85

## Phosphines

Sulfur(IV) compounds as ligands: Part XXV. Halfsandwich ruthenium thiosulfonato complexes. Crystal and molecular structure of [CpRu(dppe){SSO<sub>2</sub>(4-C<sub>6</sub>H<sub>4</sub>Cl)}], 14

## Phosphate

The photolysis of  $\text{Fe}(\text{CO})_3\{\text{P}(\text{OPh})_3\}_2$  and reaction of the photo-product with alkynes: crystal structures of  $\text{Fe}(\text{CO})_2\{\text{P}(\text{OPh})_3\}_2(\eta^2\text{-PhCCPh})$ ,  $\text{Fe}(\text{CO})_2\{\text{P}(\text{OPh})_3\}_2(\eta^1\text{:}\eta^1\text{-C(O)-C(Me)C}[\text{CH}(\text{OEt})_2]\text{C(O)})$  and  $\text{Fe}_2(\text{CO})_4\{\text{P}(\text{OPh})_3\}_2(\text{PhC-CH})_2$ , 61

## Polycarbodiimidogermylene

Polycarbodiimidogermylene: synthesis, characterization, properties and theoretical studies, 69

## Pyrazolate

Synthesis and studies on reactivities of tetrahydrofurfurylclopentadienyl lanthanide chlorides  $\text{Cp}^*\text{LnCl}_2 \cdot 2\text{THF}$  ( $\text{Ln} = \text{Dy, Ho, Gd}$ ) and  $\text{Cp}_2^*\text{LnCl} \cdot (\text{Ln} = \text{Sm, Dy})$ . The crystal structures of  $\text{Cp}_2^*\text{SmCl}$  and  $\text{Cp}_2^*\text{Sm} \cdot (\text{C}_3\text{N}_2\text{HMe}_2)$ , 176

## Rhenium

Reactivities of  $\pi$ -electron-rich phosphorus–nitrogen ligands toward Groups VIB and VIIIB metal carbonyl complexes, 160

## Rhenium complexes

Ruthenium and rhenium complexes with silyl-substituted bipyridyl ligands, 117

## Rhodium

*ortho*-Metalated dirhodium(II)-catalyzed  $\alpha$ -diazocarbonyl transformation. Diastereoselective cyclopropanation of menthyl- $\alpha$ -diao- $\beta$ -keto ester and C–H insertion of  $\alpha$ -diao ester, 36

Vibrational spectroscopy studies and density functional theory calculations on square-planar vinylidene, carbonyl and ethylene rhodium(I) complexes, 125

## Ruthenium

Sulfur(IV) compounds as ligands: Part XXV. Halfsandwich ruthenium thiosulfonato complexes. Crystal and molecular structure of [CpRu(dppe){SSO<sub>2</sub>(4-C<sub>6</sub>H<sub>4</sub>Cl)}], 14

Unbridged homo and hetero dinuclear complexes of Group 6 and 8 metals: synthesis, characterization and comparison of X-ray crystallographic data, 106

## Ruthenium complexes

Ruthenium and rhenium complexes with silyl-substituted bipyridyl ligands, 117

## Schiff base

Polynuclear cyclometallated palladium (II) complexes derived from potentially hexadentate Schiff base ligands. Crystal structures of  $[(\text{Cl})\text{Pd}\{\text{Me}_2\text{NCH}_2\text{CH}_2\text{N}=(\text{H})\text{C}\}\text{C}_6\text{H}_2\{\text{C(H)}=\text{NCH}_2\text{CH}_2\text{NMe}_2\}\text{Pd}(\text{Cl})]$  and  $[(\{\text{Ph}_2\text{PCH}_2\text{CH}_2\}_2)\text{PPh-P,P,P}\}\text{Pd}\{\text{Me}_2\text{NCH}_2\text{CH}_2\text{N}=(\text{H})\text{C}\}\text{C}_6\text{H}_2\{\text{C(H)}=\text{NCH}_2\text{CH}_2\text{NMe}_2\}\text{Pd}\{(\text{Ph}_2\text{PCH}_2\text{CH}_2)_2\text{PPh-P,P,P}\}][\text{Cl}]_2$ , 85

## Selectivity

*ortho*-Metalated dirhodium(II)-catalyzed  $\alpha$ -diazocarbonyl transformation. Diastereoselective cyclopropanation of menthyl- $\alpha$ -diao- $\beta$ -keto ester and C–H insertion of  $\alpha$ -diao ester, 36

## Semiconductors

Polycarbodiimidogermylene: synthesis, characterization, properties and theoretical studies, 69

## Silicon

Disupersilyldisilane  $\text{R}^*\text{X}_2\text{Si-SiX}_2\text{R}^*$ : Darstellung, Charakterisierung und Strukturen; sterische Effekte der Substituenten, 141

## Silyl substituted bipyridyl ligands

Ruthenium and rhenium complexes with silyl-substituted bipyridyl ligands, 117

## Steric effects

Disupersilyldisilane  $\text{R}^*\text{X}_2\text{Si-SiX}_2\text{R}^*$ : Darstellung, Charakterisierung und Strukturen; sterische Effekte der Substituenten, 141

## Stille reaction

A novel synthesis of 1,1'-unsymmetrical disubstituted heteroaryl-ferrrocene, 172

## Structure

Sulfur(IV) compounds as ligands: Part XXV. Halfsandwich ruthenium thiosulfonato complexes. Crystal and molecular structure of [CpRu(dppe){SSO<sub>2</sub>(4-C<sub>6</sub>H<sub>4</sub>Cl)}], 14

## Substitution

The reactions of *cis*-[Fe<sub>2</sub>( $\eta$ -C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>](CO)<sub>2</sub>( $\mu$ -CNMe<sub>2</sub>)<sub>2</sub>][SO<sub>3</sub>CF<sub>3</sub>]<sub>2</sub> and related compounds with nucleophiles, 18

## Sulfur

Sulfur(IV) compounds as ligands: Part XXV. Halfsandwich ruthenium thiosulfonato complexes. Crystal and molecular structure of [CpRu(dppe){SSO<sub>2</sub>(4-C<sub>6</sub>H<sub>4</sub>Cl)}], 14

## Synthesis

Synthesis and studies on reactivities of tetrahydrofurfurylclopentadienyl lanthanide chlorides  $\text{Cp}^*\text{LnCl}_2 \cdot 2\text{THF}$  ( $\text{Ln} = \text{Dy, Ho, Gd}$ ) and  $\text{Cp}_2^*\text{LnCl} \cdot (\text{Ln} = \text{Sm, Dy})$ . The crystal structures of  $\text{Cp}_2^*\text{SmCl}$  and  $\text{Cp}_2^*\text{Sm} \cdot (\text{C}_3\text{N}_2\text{HMe}_2)$ , 176

## T-dependent

$^{57}\text{Fe}$ -Mössbauer spectroscopic study of monomeric and polymeric ferrocenyl- and octamethylferrocenyl-substituted ethynes, 1

## Tetrahydrofurfurylcyclopentadienyl

Synthesis and studies on reactivities of tetrahydrofurfurylcyclopentadienyl lanthanide chlorides  $\text{Cp}^*\text{LnCl}_2\cdot 2\text{THF}$  ( $\text{Ln} = \text{Dy}, \text{Ho}, \text{Gd}$ ) and  $\text{Cp}_2^*\text{LnCl}\cdot (\text{Ln} = \text{Sm}, \text{Dy})$ . The crystal structures of  $\text{Cp}_2^*\text{SmCl}$  and  $\text{Cp}_2^*\text{Sm}\cdot (\text{C}_3\text{N}_2\text{HMe}_2)$ , 176

## Theoretical studies

Polycarbodiimidogermmylene: synthesis, characterization, properties and theoretical studies, 69

## Thiosulfonate

Sulfur(IV) compounds as ligands: Part XXV. Halfsandwich ruthenium thiosulfonato complexes. Crystal and molecular structure of  $[\text{CpRu}(\text{dppe})\{\text{SSO}_2(4-\text{C}_6\text{H}_4\text{Cl})\}]$ , 14

## Transition metal complexes

Transition metal complexes of 2,2,2-trifluoroethyl isocyanide and 1H,1H-perfluoroctyl isocyanide, 96

## 1,1'-Unsymmetrical diheteroaryl-ferrocene

A novel synthesis of 1,1'-unsymmetrical disubstituted heteroaryl-ferrocene, 172

## Vinylidene complexes

Vibrational spectroscopy studies and density functional theory calculations on square-planar vinylidene, carbonyl and ethylene rhodium(I) complexes, 125

## X-ray

Antiferromagnetic complexes with metal–metal bonds: Part XXIX. Synthesis and molecular structures of heterochalcogenide binuclear complex  $[\pi-(\text{CH}_3\text{C}_5\text{H}_4)\text{Cr}(\mu\text{-SPh})_2\text{Se}$  and trinuclear mixed-metal cluster  $[\pi-(\text{CH}_3\text{C}_5\text{H}_4)_2\text{Cr}_2(\mu\text{-SPh})(\mu_3\text{-S})(\mu_3\text{-Se})\text{Co}(\text{CO})_2]$ , 9

## X-ray structure analyses

Disupersilyldisilane  $\text{R}^*\text{X}_2\text{Si}-\text{SiX}_2\text{R}^*$ : Darstellung, Charakterisierung und Strukturen; sterische Effekte der Substituenten, 141