

## Subject Index of Volume 619

### Acaricidal activity

Synthesis and structural characterization of several tris(2-methyl-2-phenylpropyl)tin carboxylates containing germanium, 43

### Acetyl complex

Structural and spectroscopic characterization of ruthenium(II) complexes with methyl, formyl, and acetyl groups as model species in multi-step CO<sub>2</sub> reduction, 299

### Acetylide

Synthesis of highly ethynylated mono and dinuclear Pt(II) tethers bearing the 4,4'-bis(ethynyl)biphenyl (debp) unit as central core, 49

### Alcohol

Direct synthesis of alkyldialkoxysilanes by the reaction of silicon, alcohol and alkene using a high-pressure flow reactor, 235

### Alkene

Direct synthesis of alkyldialkoxysilanes by the reaction of silicon, alcohol and alkene using a high-pressure flow reactor, 235

### 1-Alkylaminopyrazoles

Organometallic rhodium (I) complexes with 1-alkylaminopyrazole ligands, 14

### Alkyne

Stabilisation of alkynyl dithiocarboxylates at a ruthenium centre: X-ray crystal structures of [Ru(H)(CO)(S<sub>2</sub>CC≡CPh)(PPh<sub>3</sub>)<sub>2</sub>], [Ru(CO)(CPh=CHPh)(S<sub>2</sub>CC≡CPh)(PPh<sub>3</sub>)<sub>2</sub>] and [Ru(CO)(C≡CMes)(S<sub>2</sub>CC≡CPh)(PPh<sub>3</sub>)<sub>2</sub>], 209

### Alkyne cyclisation

New trisubstituted cyclopentadienyl ligands: synthesis, characterisation and catalytic properties of mono and dinuclear cobalt, rhodium, iron and ruthenium complexes, 179

### Alkynyl

Donor-funktionalisierte Alkynyle von Titan(IV), Kupfer(I) und Silber(I); Festkörperstrukturen von [Ti](C≡CCH<sub>2</sub>NMe<sub>2</sub>)<sub>2</sub> und {[Ti](C≡C'Bu)<sub>2</sub>}CuC≡CC≡CC<sub>2</sub>H<sub>5</sub>, 99

### 1-Allyl-2-formylpyrrole

A shortcut hydroformylation route to 7-formyl-5,6-dihydroindolizine from 1-allyl-2-formylpyrrole, 241

### Allylgermanes

Palladium complex-catalyzed germylation of allylic halides using (germyl)stannanes, 313

### Allyl halides

Palladium complex-catalyzed germylation of allylic halides using (germyl)stannanes, 313

### Aluminium

Aluminium zirconium (+ 3 and + 4) heterometallic hydrido complexes of compositions [(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>Zr(μ-H)]<sub>2</sub>(μ-H)AlCl<sub>2</sub> and [(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>ZrH(μ-H)]<sub>2</sub>Al, 36

### ansa-Metallocenes

Synthesis and X-ray crystal structures of *ansa*-complexes of titanium, zirconium, and hafnium involving methylene-bis(indenyl) ligand, 280

### Arsaphosphene

New routes to diarsenes and arsaphosphenes involving fluorinated arsines, 275

### Arylpalladium complexes

Thioether- and selenoether-carboxylates in palladium chemistry: conclusive proof of hemilabile properties of O–Se ligands, 164

### Asymmetric catalysis

Synthesis and crystal structure of bis[(4*S*,5*S*)-4,5-dihydro-4,5-diphenyl-2-(2'-oxidophenyl-χO)oxazole-χN]copper(II) and its application in the asymmetric Baeyer–Villiger reaction, 204

### Baeyer–Villiger reaction

Synthesis and crystal structure of bis[(4*S*,5*S*)-4,5-dihydro-4,5-diphenyl-2-(2'-oxidophenyl-χO)oxazole-χN]copper(II) and its application in the asymmetric Baeyer–Villiger reaction, 204

### Benzonitrile

Organometallic complexes for second-order non-linear optics: synthesis and molecular quadratic hyperpolarizabilities of η<sup>5</sup>-monocyclopentadienyliron(II) nitrile derivatives with different phosphines. X-ray crystal structure of [FeCp(DPPE)(*p*-NCC<sub>6</sub>H<sub>4</sub>NO<sub>2</sub>)] [PF<sub>6</sub>]-CH<sub>2</sub>Cl<sub>2</sub>, 252

### Benzyl group abstraction

Benzyl group abstraction from Ti(CH<sub>2</sub>Ph)<sub>4</sub> by B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> and [Ph<sub>3</sub>C][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>] to form Ziegler–Natta and carbocationic alkene polymerization initiators, 62

### bicyclo- and tetrahydro-Tetrasilanes

Disilene R\*XS<sub>i</sub>–SiXR\* (R\*–Si*t*Bu<sub>3</sub>) mit siliciumgebundenen H- und Hal-Atomen X: Bildung, Isomerisierung, Reaktionen, 110

### bis-(Ferroceny)butadiyne

Insertion of bis-ferrocenylbutadiyne into an osmium hydride bond. The synthesis, structure and electrochemical response of Os<sub>4</sub>(CO)<sub>11</sub>(μ-η-Z-FcCC(H)C<sub>2</sub>Fc)(μ-H)<sub>3</sub>, 271

### Bithienyl

The effect of different bithienyl spacers on the linear and non-linear optical properties of organometallic chromophores derived from [Fe<sub>2</sub>(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>(CO)<sub>2</sub>(μ-CO)(μ-C–CH<sub>3</sub>)]<sup>+</sup>[BF<sub>4</sub>]<sup>–</sup>. Crystal structure of [Fe<sub>2</sub>(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>(CO)<sub>2</sub>(μ-CO)(μ-(*E*)-C–CH=CH-7-(thieno[3,2-*e*][1]benzothio-phenyl))]<sup>+</sup>[BF<sub>4</sub>]<sup>–</sup>, 141

### Boron/bis(cyclopentadienyl)boranes

Synthesis and structure of aminobis(η<sup>1</sup>-cyclopentadienyl)boranes and related compounds, 305

### Carbocationic initiator

Benzyl group abstraction from Ti(CH<sub>2</sub>Ph)<sub>4</sub> by B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> and [Ph<sub>3</sub>C][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>] to form Ziegler–Natta and carbocationic alkene polymerization initiators, 62

### Carbohydrate

Synthesis, solution and solid-state NMR spectra of methyl 4,6-*O*-benzylidene-2,3-bis-*O*-(organostannylmethyl)-α-D-glucopyranosides: Crystal structure of methyl 4,6-*O*-benzylidene-2,3-bis-*O*-[(dimethylphenylstannyl)methyl]-α-D-glucopyranoside, 226

## Carbon–carbon bond formation

Preparations and selectivities of magnesium-based mixed cuprates, 31

## Carbon–copper bond

Preparations and selectivities of magnesium-based mixed cuprates, 31

## Carbon monoxide

New catalysts for the alternating copolymerization of 4-*tert*-butylstyrene/CO, 287

## Carbonyl

Selective CO oxidation of heterometallic carbonyl clusters with the oxygen transfer reagent (*p*-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>)<sub>2</sub>TeO. Crystal structures of (μ<sub>3</sub>-S)FeCoMo(CO)<sub>7</sub>(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>COCH<sub>3</sub>)(PPh<sub>3</sub>) and (μ<sub>3</sub>-Se)FeCoMo(CO)<sub>7</sub>(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>COCH<sub>3</sub>)(PPh<sub>3</sub>), 194

## Catalysis

A shortcut hydroformylation route to 7-formyl-5,6-dihydroindolizine from 1-allyl-2-formylpyrrole, 241

New catalysts for the alternating copolymerization of 4-*tert*-butylstyrene/CO, 287

## Chiral-at-metal complexes

Synthesis of rhodium(III) complexes with the chiral phosphine (*S*)-Ph<sub>2</sub>PCH<sub>2</sub>CHMeCH<sub>2</sub>OH. Crystal structure of (S<sub>Rh</sub>,S<sub>C</sub>)-[(η<sup>5</sup>-C<sub>5</sub>Me<sub>5</sub>)RhCl(η<sup>2</sup>-PPh<sub>2</sub>CH<sub>2</sub>CHMeCH<sub>2</sub>OH-P,O)]BF<sub>4</sub>, 1

## Chiral bidentate ligand

Synthesis of rhodium(III) complexes with the chiral phosphine (*S*)-Ph<sub>2</sub>PCH<sub>2</sub>CHMeCH<sub>2</sub>OH. Crystal structure of (S<sub>Rh</sub>,S<sub>C</sub>)-[(η<sup>5</sup>-C<sub>5</sub>Me<sub>5</sub>)RhCl(η<sup>2</sup>-PPh<sub>2</sub>CH<sub>2</sub>CHMeCH<sub>2</sub>OH-P,O)]BF<sub>4</sub>, 1

## Chirality

Funktionelle Phosphane: Part XI. Optisch reine β-Aminophosphane und β-Aminophosphinite für die komplexkatalysierte Reduktion organischer Carbonylverbindungen. Molekülstrukturen von [(1*R*,2*R*)-Ph<sub>2</sub>PCH(Ph)CH(Me)-NH<sub>2</sub>Me]Cl, (1*R*,2*S*)-Ph<sub>2</sub>PCH(Ph)CH(Me)NHSO<sub>2</sub>Me und [(1*R*,2*R*)-Ph<sub>2</sub>PCH(Ph)CH(Me)NHMe-κ*N*,κ*P*];Rh(η<sup>4</sup>-1,5-C<sub>8</sub>H<sub>12</sub>)]BF<sub>4</sub>, 88

## Chiral oxazoline

Synthesis and crystal structure of bis[(4*S*,5*S*)-4,5-dihydro-4,5-diphenyl-2-(2'-oxidophenyl-χO)oxazole-χN]copper(II) and its application in the asymmetric Baeyer–Villiger reaction, 204

## Chiral phosphine

Synthesis of rhodium(III) complexes with the chiral phosphine (*S*)-Ph<sub>2</sub>PCH<sub>2</sub>CHMeCH<sub>2</sub>OH. Crystal structure of (S<sub>Rh</sub>,S<sub>C</sub>)-[(η<sup>5</sup>-C<sub>5</sub>Me<sub>5</sub>)RhCl(η<sup>2</sup>-PPh<sub>2</sub>CH<sub>2</sub>CHMeCH<sub>2</sub>OH-P,O)]BF<sub>4</sub>, 1

## Cluster

Insertion of bis-ferrocenylbutadiyne into an osmium hydride bond. The synthesis, structure and electrochemical response of Os<sub>4</sub>(CO)<sub>11</sub>(μ-η-Z-FcCC(H)C<sub>2</sub>Fc)(μ-H)<sub>3</sub>, 271

Transition metal clusters containing carbosilane dendrimers, 245

## Conformation analysis

Multiply trimethylstannyl substituted ferrocenes synthesis, NMR studies, X-ray structural analysis and electrochemistry, 74

## Conjugate addition

Preparations and selectivities of magnesium-based mixed cuprates, 31

## Constitutional isomers

Synthesis and structure of aminobis(η<sup>1</sup>-cyclopentadienyl)boranes and related compounds, 305

## Coordination bond

Nature of the metal–ligand bond in trivalent neodymium complexes with neutral π-donor ligands. A theoretical study, 24

## Copolymerization

New catalysts for the alternating copolymerization of 4-*tert*-butylstyrene/CO, 287

## Copper

Synthesis, spectroscopic and structural characterisation of complexes of the new ditelluroether 1,2-bis(methyltelluro-

methyl)benzene, *o*-C<sub>6</sub>H<sub>4</sub>(CH<sub>2</sub>TeMe)<sub>2</sub>. X-ray structures of [Mn(CO)<sub>3</sub>Cl{*o*-C<sub>6</sub>H<sub>4</sub>(CH<sub>2</sub>TeMe)<sub>2</sub>}] and [W(CO)<sub>4</sub>{*o*-C<sub>6</sub>H<sub>4</sub>(CH<sub>2</sub>TeMe)<sub>2</sub>}], 218

## Copper catalyst

Direct synthesis of alkyldialkoxysilanes by the reaction of silicon, alcohol and alkene using a high-pressure flow reactor, 235

## Copper(I)

Donor-funktionalisierte Alkinyne von Titan(IV), Kupfer(I) und Silber(I); Festkörperstrukturen von [Ti](C≡CCH<sub>2</sub>NMe<sub>2</sub>)<sub>2</sub> und {[Ti](C≡C'Bu)<sub>2</sub>}CuC≡CC≡CC<sub>2</sub>H<sub>5</sub>, 99

## Crystal structures

Modelling nucleophilic substitution at silicon using hypervalent silicon compounds based on urea ligands, 132

Organometallic complexes for second-order non-linear optics: synthesis and molecular quadratic hyperpolarizabilities of η<sup>5</sup>-monocyclopentadienyliron(II) nitrile derivatives with different phosphines. X-ray crystal structure of [FeCp-(DPPE)(*p*-NCC<sub>6</sub>H<sub>4</sub>NO<sub>2</sub>)]PF<sub>6</sub>·CH<sub>2</sub>Cl<sub>2</sub>, 252

Selective CO oxidation of heterometallic carbonyl clusters with the oxygen transfer reagent (*p*-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>)<sub>2</sub>TeO. Crystal structures of (μ<sub>3</sub>-S)FeCoMo(CO)<sub>7</sub>(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>COCH<sub>3</sub>)(PPh<sub>3</sub>) and (μ<sub>3</sub>-Se)FeCoMo(CO)<sub>7</sub>(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>COCH<sub>3</sub>)(PPh<sub>3</sub>), 194

Stabilisation of alkynyl dithiocarboxylates at a ruthenium centre: X-ray crystal structures of [Ru(H)(CO)(S<sub>2</sub>CC≡CPh)(PPh<sub>3</sub>)<sub>2</sub>], [Ru(CO)(CPh=CHPh)(S<sub>2</sub>CC≡CPh)(PPh<sub>3</sub>)<sub>2</sub>] and [Ru(CO)(C≡CMes)(S<sub>2</sub>CC≡CPh)(PPh<sub>3</sub>)<sub>2</sub>], 209

Structural studies of diorganotin(IV) sulphonates. The crystal structures of [(*n*-C<sub>4</sub>H<sub>9</sub>)<sub>2</sub>Sn{μ-OSO<sub>2</sub>C<sub>6</sub>H<sub>2</sub>(CH<sub>3</sub>)<sub>3</sub>}<sub>2</sub>]<sub>*n*</sub> and its partially hydrolyzed product [(*n*-C<sub>4</sub>H<sub>9</sub>)<sub>2</sub>Sn{(μ-OH)(μ-OSO<sub>2</sub>C<sub>6</sub>H<sub>2</sub>(CH<sub>3</sub>)<sub>3</sub>)}]<sub>*n*</sub>, 157

Synthesis and crystal structure of bis[(4*S*,5*S*)-4,5-dihydro-4,5-diphenyl-2-(2'-oxidophenyl-χO)oxazole-χN]copper(II) and its application in the asymmetric Baeyer–Villiger reaction, 204

Synthesis of rhodium(III) complexes with the chiral phosphine (*S*)-Ph<sub>2</sub>PCH<sub>2</sub>CHMeCH<sub>2</sub>OH. Crystal structure of (S<sub>Rh</sub>,S<sub>C</sub>)-[(η<sup>5</sup>-C<sub>5</sub>Me<sub>5</sub>)RhCl(η<sup>2</sup>-PPh<sub>2</sub>CH<sub>2</sub>CHMeCH<sub>2</sub>OH-P,O)]BF<sub>4</sub>, 1

Synthesis, solution and solid-state NMR spectra of methyl 4,6-*O*-benzylidene-2,3-bis-*O*-(organostannylmethyl)-α-D-glucopyranosides: Crystal structure of methyl 4,6-*O*-benzylidene-2,3-bis-*O*-[(dimethylphenylstannyl)methyl]-α-D-glucopyranoside, 226

## Cyclopentadienyl complexes: cobalt, rhodium, iron, ruthenium

New trisubstituted cyclopentadienyl ligands: synthesis, characterisation and catalytic properties of mono and dinuclear cobalt, rhodium, iron and ruthenium complexes, 179

## Cyclosilanes

Disilene R\*XS<sub>i</sub>=SiXR\* (R\*=Si*t*Bu<sub>3</sub>) mit siliciumgebundenen H- und Hal-Atomen X: Bildung, Isomerisierung, Reaktionen, 110

## Dendrimer

Transition metal clusters containing carbosilane dendrimers, 245

## DFT calculation

Nature of the metal–ligand bond in trivalent neodymium complexes with neutral π-donor ligands. A theoretical study, 24

## Diarsene

New routes to diarsenes and arsaphosphenes involving fluorinated arsines, 275

## Diboroles

Syntheses and ligand properties of 1,3-di*t*-butyl-2,3-dihydro-1,3-diboroles. Rhodium-(η<sup>5</sup>-2,3-dihydro-1,3-diborole) and rhodium-(η<sup>5</sup>-2,3-dihydro-1,3-diborolyl) complexes, 7

## Diindenylboranes

Synthesis and structure of aminobis(η<sup>1</sup>-cyclopentadienyl)boranes and related compounds, 305

## Dinuclear

Synthesis of highly ethynylated mono and dinuclear Pt(II) tethers bearing the 4,4'-bis(ethynyl)biphenyl (debp) unit as central core, 49

## Diorganotin compounds

Structural studies of diorganotin(IV) sulphonates. The crystal structures of  $[(n-C_4H_9)_2Sn\{\mu-OSO_2C_6H_2(CH_3)_3\}_2]_n$  and its partially hydrolyzed product  $[(n-C_4H_9)_2Sn\{(\mu-OH)(\mu-OSO_2C_6H_2(CH_3)_3)\}]_n$ , 157

## Direct synthesis

Direct synthesis of alkyldialkoxysilanes by the reaction of silicon, alcohol and alkene using a high-pressure flow reactor, 235

## Disilanides

Disilene  $R^*XSi-SiXR^*$  ( $R^*=Si^tBu_3$ ) mit siliciumgebundenen H- und Hal-Atomen X: Bildung, Isomerisierung, Reaktionen, 110

## Disilenes

Disilene  $R^*XSi-SiXR^*$  ( $R^*=Si^tBu_3$ ) mit siliciumgebundenen H- und Hal-Atomen X: Bildung, Isomerisierung, Reaktionen, 110

## Early-late complexes

Donor-funktionalisierte Alkinyne von Titan(IV), Kupfer(I) und Silber(I); Festkörperstrukturen von  $[Ti](C\equiv CCH_2NMe_2)_2$  und  $\{[Ti](C\equiv C^tBu)_2\}CuC\equiv CC\equiv CC_2H_5$ , 99

## Electrochemistry

Multiply trimethylstannyl substituted ferrocenes synthesis, NMR studies, X-ray structural analysis and electrochemistry, 74

## Electronic communication

Insertion of bis-ferrocenylbutadiyne into an osmium hydride bond. The synthesis, structure and electrochemical response of  $Os_4(CO)_{11}(\mu-\eta-Z-FcCC(H)C_2Fc)(\mu-H)_3$ , 271

## Electrospray mass spectra

Organometallic rhodium (I) complexes with 1-alkylaminopyrazole ligands, 14

## Ferrocene

Synthesis of highly ethynylated mono and dinuclear Pt(II) tethers bearing the 4,4'-bis(ethynyl)biphenyl (debp) unit as central core, 49

The effect of different bithienyl spacers on the linear and non-linear optical properties of organometallic chromophores derived from  $[Fe_2(\eta^5-C_5H_5)_2(CO)_2(\mu-CO)(\mu-C-CH_3)]^+[BF_4]^-$ . Crystal structure of  $[Fe_2(\eta^5-C_5H_5)_2(CO)_2(\mu-CO)(\mu-E)-C-CH=CH-7-(thieno[3,2-e][1]benzothiophene)]^+[BF_4]^-$ , 141

## First hyperpolarizability

Organometallic complexes for second-order non-linear optics: synthesis and molecular quadratic hyperpolarizabilities of  $\eta^5$ -monocyclopentadienyliron(II) nitrile derivatives with different phosphines. X-ray crystal structure of  $[FeCp(DPPE)(p-NCC_6H_4NO_2)][PF_6]\cdot CH_2Cl_2$ , 252

## Fluorinated arsines

New routes to diarsenes and arsaphosphenes involving fluorinated arsines, 275

## Fluoroarsinylphosphine

New routes to diarsenes and arsaphosphenes involving fluorinated arsines, 275

## Formyl complex

Structural and spectroscopic characterization of ruthenium(II) complexes with methyl, formyl, and acetyl groups as model species in multi-step  $CO_2$  reduction, 299

## Germylstannanes

Palladium complex-catalyzed germylation of allylic halides using (germyl)stannanes, 313

## Gold

Transition metal clusters containing carbosilane dendrimers, 245

## Hafnium

Synthesis and X-ray crystal structures of *ansa*-complexes of titanium, zirconium, and hafnium involving methylene-bis(indenyl) ligand, 280

## Hemilability

Thioether- and selenoether-carboxylates in palladium chemistry: conclusive proof of hemilabile properties of O-Se ligands, 164

## Heterocycles

A new class of triazepine-ruthenium(II) complexes: synthesis, crystal structure and NMR spectroscopic properties, 265

## Heterometallic cluster

Selective CO oxidation of heterometallic carbonyl clusters with the oxygen transfer reagent (*p*- $CH_3OC_6H_4$ ) $_2TeO$ . Crystal structures of  $(\mu_3-S)FeCoMo(CO)_7(\eta^5-C_5H_4COCH_3)(PPh_3)$  and  $(\mu_3-Se)FeCoMo(CO)_7(\eta^5-C_5H_4COCH_3)(PPh_3)$ , 194

## Hydride

Stabilisation of alkynyl dithiocarboxylates at a ruthenium centre: X-ray crystal structures of  $[Ru(H)(CO)(S_2CC\equiv CPh)(PPh_3)_2]$ ,  $[Ru(CO)(CPh=CHPh)(S_2CC\equiv CPh)(PPh_3)_2]$  and  $[Ru(CO)(C\equiv CMes)(S_2CC\equiv CPh)(PPh_3)_2]$ , 209

## Hydrides

Aluminium zirconium (+3 and +4) heterometallic hydrido complexes of compositions  $[(\eta^5-C_5H_5)_2Zr(\mu-H)_2(\mu-H)AlCl_2]$  and  $[(\eta^5-C_5H_5)_2ZrH(\mu-H)_2Al]$ , 36

## Hydroformylation

A shortcut hydroformylation route to 7-formyl-5,6-dihydroindolizine from 1-allyl-2-formylpyrrole, 241

New trisubstituted cyclopentadienyl ligands: synthesis, characterisation and catalytic properties of mono and dinuclear cobalt, rhodium, iron and ruthenium complexes, 179

## Hydrogenation

Funktionelle Phosphane: Part XI. Optisch reine  $\beta$ -Aminophosphane und  $\beta$ -Aminophosphinite für die komplexkatalysierte Reduktion organischer Carbonylverbindungen. Molekülstrukturen von  $[(1R,2R)-Ph_2PCH(Ph)CH(Me)-NH_2Me]Cl$ ,  $(1R,2S)-Ph_2PCH(Ph)CH(Me)NHSO_2Me$  und  $\{[(1R,2R)-Ph_2PCH(Ph)CH(Me)NHMe-\kappa N, \kappa P]Rh(\eta^4-1,5-C_8H_{12})\}BF_4$ , 88

## Hyperpolarisability

The effect of different bithienyl spacers on the linear and non-linear optical properties of organometallic chromophores derived from  $[Fe_2(\eta^5-C_5H_5)_2(CO)_2(\mu-CO)(\mu-C-CH_3)]^+[BF_4]^-$ . Crystal structure of  $[Fe_2(\eta^5-C_5H_5)_2(CO)_2(\mu-CO)(\mu-E)-C-CH=CH-7-(thieno[3,2-e][1]benzothiophene)]^+[BF_4]^-$ , 141

## Hyper-Raleigh scattering

The effect of different bithienyl spacers on the linear and non-linear optical properties of organometallic chromophores derived from  $[Fe_2(\eta^5-C_5H_5)_2(CO)_2(\mu-CO)(\mu-C-CH_3)]^+[BF_4]^-$ . Crystal structure of  $[Fe_2(\eta^5-C_5H_5)_2(CO)_2(\mu-CO)(\mu-E)-C-CH=CH-7-(thieno[3,2-e][1]benzothiophene)]^+[BF_4]^-$ , 141

## Indolizine

A shortcut hydroformylation route to 7-formyl-5,6-dihydroindolizine from 1-allyl-2-formylpyrrole, 241

## Insertion

Insertion of bis-ferrocenylbutadiyne into an osmium hydride bond. The synthesis, structure and electrochemical response of  $Os_4(CO)_{11}(\mu-\eta-Z-FcCC(H)C_2Fc)(\mu-H)_3$ , 271

## IR

Synthesis and structural characterization of several tris(2-methyl-2-phenylpropyl)tin carboxylates containing germanium, 43

## IR and Raman spectroscopy

Structural and spectroscopic characterization of ruthenium(II) complexes with methyl, formyl, and acetyl groups as model species in multi-step CO<sub>2</sub> reduction, 299

## Iridium

Funktionelle Phosphane: Part XI. Optisch reine β-Aminophosphane und β-Aminophosphinite für die komplex-katalysierte Reduktion organischer Carbonylverbindungen. Molekülstrukturen von [(1*R*,2*R*)-Ph<sub>2</sub>PCH(Ph)CH(Me)-NH<sub>2</sub>Me]Cl, (1*R*,2*S*)-Ph<sub>2</sub>PCH(Ph)CH(Me)NHSO<sub>2</sub>Me und [{(1*R*,2*R*)-Ph<sub>2</sub>PCH(Ph)CH(Me)NHMe-κ*N*,κ*P*};Rh(η<sup>4</sup>-1,5-C<sub>8</sub>H<sub>12</sub>)]BF<sub>4</sub>, 88

## Iron

Multiply trimethylstannyl substituted ferrocenes synthesis, NMR studies, X-ray structural analysis and electrochemistry, 74

Organometallic complexes for second-order non-linear optics: synthesis and molecular quadratic hyperpolarizabilities of η<sup>5</sup>-monocyclopentadienyliron(II) nitrile derivatives with different phosphines. X-ray crystal structure of [FeCp(DPPE)(*p*-NCC<sub>6</sub>H<sub>4</sub>NO<sub>2</sub>)]PF<sub>6</sub>·CH<sub>2</sub>Cl<sub>2</sub>, 252

The effect of different bithienyl spacers on the linear and non-linear optical properties of organometallic chromophores derived from [Fe<sub>2</sub>(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>(CO)<sub>2</sub>(μ-CO)(μ-C-CH<sub>3</sub>)]<sup>+</sup>[BF<sub>4</sub>]<sup>-</sup>. Crystal structure of [Fe<sub>2</sub>(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>(CO)<sub>2</sub>(μ-CO)(μ-*E*)-C-CH=CH-7-(thieno[3,2-*e*]1]benzothiophene)]<sup>+</sup>[BF<sub>4</sub>]<sup>-</sup>, 141

## Manganese

Synthesis, spectroscopic and structural characterisation of complexes of the new ditelluroether 1,2-bis(methyltelluromethyl)benzene, *o*-C<sub>6</sub>H<sub>4</sub>(CH<sub>2</sub>TeMe)<sub>2</sub>. X-ray structures of [Mn(CO)<sub>3</sub>Cl{*o*-C<sub>6</sub>H<sub>4</sub>(CH<sub>2</sub>TeMe)<sub>2</sub>}] and [W(CO)<sub>4</sub>{*o*-C<sub>6</sub>H<sub>4</sub>(CH<sub>2</sub>TeMe)<sub>2</sub>}], 218

Transition metal clusters containing carbosilane dendrimers, 245

## Mercury

Phenyl derivatives of mercury functionalised in the 2-position with either a formyl or a vinyl group. Crystal structures of Hg[C<sub>6</sub>H<sub>4</sub>(CHO)-2]Cl and Hg[C<sub>6</sub>H<sub>4</sub>(CH=CH<sub>2</sub>)-2]<sub>2</sub>, 293

## Metallic silicon

Direct synthesis of alkyldialkoxysilanes by the reaction of silicon, alcohol and alkene using a high-pressure flow reactor, 235

## Metallocenes

Aluminium zirconium (+ 3 and + 4) heterometallic hydrido complexes of compositions [(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>Zr(μ-H)]<sub>2</sub>(μ-H)AlCl<sub>2</sub> and [(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>ZrH(μ-H)]<sub>2</sub>Al, 36

## Metathesis

Palladium complex-catalyzed germylation of allylic halides using (germyl)stannanes, 313

## Methyl complex

Structural and spectroscopic characterization of ruthenium(II) complexes with methyl, formyl, and acetyl groups as model species in multi-step CO<sub>2</sub> reduction, 299

## Mixed cuprates

Preparations and selectivities of magnesium-based mixed cuprates, 31

## Molecular dynamics

Multiply trimethylstannyl substituted ferrocenes synthesis, NMR studies, X-ray structural analysis and electrochemistry, 74

## Molybdenum

Transition metal clusters containing carbosilane dendrimers, 245

## Multinuclear NMR

Synthesis and structural characterization of several tris(2-methyl-2-phenylpropyl)tin carboxylates containing germanium, 43

## Neodymium

Nature of the metal–ligand bond in trivalent neodymium complexes with neutral π-donor ligands. A theoretical study, 24

## Nickel(0)

Donor-funktionalisierte Alkinyne von Titan(IV), Kupfer(I) und Silber(I); Festkörperstrukturen von [Ti](C≡CCH<sub>2</sub>NMe<sub>2</sub>)<sub>2</sub> und {[Ti](C≡C<sup>*t*</sup>Bu)<sub>2</sub>}CuC≡CC≡CC<sub>2</sub>H<sub>5</sub>, 99

## Nitrogen ligands

New catalysts for the alternating copolymerization of 4-*tert*-butylstyrene/CO, 287

## NLO materials

Organometallic complexes for second-order non-linear optics: synthesis and molecular quadratic hyperpolarizabilities of η<sup>5</sup>-monocyclopentadienyliron(II) nitrile derivatives with different phosphines. X-ray crystal structure of [FeCp(DPPE)(*p*-NCC<sub>6</sub>H<sub>4</sub>NO<sub>2</sub>)]PF<sub>6</sub>·CH<sub>2</sub>Cl<sub>2</sub>, 252

## NMR

A new class of triazepine–ruthenium(II) complexes: synthesis, crystal structure and NMR spectroscopic properties, 265

## Non-linear optics

Organometallic complexes for second-order non-linear optics: synthesis and molecular quadratic hyperpolarizabilities of η<sup>5</sup>-monocyclopentadienyliron(II) nitrile derivatives with different phosphines. X-ray crystal structure of [FeCp(DPPE)(*p*-NCC<sub>6</sub>H<sub>4</sub>NO<sub>2</sub>)]PF<sub>6</sub>·CH<sub>2</sub>Cl<sub>2</sub>, 252

The effect of different bithienyl spacers on the linear and non-linear optical properties of organometallic chromophores derived from [Fe<sub>2</sub>(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>(CO)<sub>2</sub>(μ-CO)(μ-C-CH<sub>3</sub>)]<sup>+</sup>[BF<sub>4</sub>]<sup>-</sup>. Crystal structure of [Fe<sub>2</sub>(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>(CO)<sub>2</sub>(μ-CO)(μ-*E*)-C-CH=CH-7-(thieno[3,2-*e*]1]benzothiophene)]<sup>+</sup>[BF<sub>4</sub>]<sup>-</sup>, 141

*o*-Formylphenyl complexes

Phenyl derivatives of mercury functionalised in the 2-position with either a formyl or a vinyl group. Crystal structures of Hg[C<sub>6</sub>H<sub>4</sub>(CHO)-2]Cl and Hg[C<sub>6</sub>H<sub>4</sub>(CH=CH<sub>2</sub>)-2]<sub>2</sub>, 293

## Olefin

Nature of the metal–ligand bond in trivalent neodymium complexes with neutral π-donor ligands. A theoretical study, 24

## Organogermanium

Synthesis and structural characterization of several tris(2-methyl-2-phenylpropyl)tin carboxylates containing germanium, 43

## Organometallic complexes

Phenyl derivatives of mercury functionalised in the 2-position with either a formyl or a vinyl group. Crystal structures of Hg[C<sub>6</sub>H<sub>4</sub>(CHO)-2]Cl and Hg[C<sub>6</sub>H<sub>4</sub>(CH=CH<sub>2</sub>)-2]<sub>2</sub>, 293

## Organosilane

Direct synthesis of alkyldialkoxysilanes by the reaction of silicon, alcohol and alkene using a high-pressure flow reactor, 235

## Organotin

Synthesis and structural characterization of several tris(2-methyl-2-phenylpropyl)tin carboxylates containing germanium, 43

Synthesis, solution and solid-state NMR spectra of methyl 4,6-*O*-benzylidene-2,3-bis-*O*-(organostannylmethyl)-α-D-glucopyranosides: Crystal structure of methyl 4,6-*O*-benzylidene-2,3-bis-*O*-[(dimethylphenylstannyl)methyl]-α-D-glucopyranoside, 226

## Osmium

Insertion of bis-ferrocenylbutadiene into an osmium hydride bond. The synthesis, structure and electrochemical response of Os<sub>4</sub>(CO)<sub>11</sub>(μ-η-Z-FcCC(H)C<sub>2</sub>Fc)(μ-H)<sub>3</sub>, 271

*o*-Vinylphenyl complexes

Phenyl derivatives of mercury functionalised in the 2-position with either a formyl or a vinyl group. Crystal structures of Hg[C<sub>6</sub>H<sub>4</sub>(CHO)-2]Cl and Hg[C<sub>6</sub>H<sub>4</sub>(CH=CH<sub>2</sub>)-2]<sub>2</sub>, 293

## Oxidation

Selective CO oxidation of heterometallic carbonyl clusters with the oxygen transfer reagent (*p*-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>)<sub>2</sub>TeO. Crystal structures of (μ<sub>3</sub>-S)FeCoMo(CO)<sub>7</sub>(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>COCH<sub>3</sub>)(PPh<sub>3</sub>) and (μ<sub>3</sub>-Se)FeCoMo(CO)<sub>7</sub>(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>COCH<sub>3</sub>)(PPh<sub>3</sub>), 194

## Oxygen–selenium ligands

Thioether- and selenoether-carboxylates in palladium chemistry: conclusive proof of hemilabile properties of O–Se ligands, 164

## Oxygen–sulphur ligands

Thioether- and selenoether-carboxylates in palladium chemistry: conclusive proof of hemilabile properties of O–Se ligands, 164

## Oxygen transfer reagent

Selective CO oxidation of heterometallic carbonyl clusters with the oxygen transfer reagent (*p*-CH<sub>3</sub>OC<sub>6</sub>H<sub>4</sub>)<sub>2</sub>TeO. Crystal structures of (μ<sub>3</sub>-S)FeCoMo(CO)<sub>7</sub>(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>COCH<sub>3</sub>)(PPh<sub>3</sub>) and (μ<sub>3</sub>-Se)FeCoMo(CO)<sub>7</sub>(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>COCH<sub>3</sub>)(PPh<sub>3</sub>), 194

## Palladium

Thioether- and selenoether-carboxylates in palladium chemistry: conclusive proof of hemilabile properties of O–Se ligands, 164

## Palladium catalysis

Palladium complex-catalyzed germylation of allylic halides using (germyl)stannanes, 313

## Palladium complexes

New catalysts for the alternating copolymerization of 4-*tert*-butylstyrene/CO, 287

## Pentacoordination

Modelling nucleophilic substitution at silicon using hypervalent silicon compounds based on urea ligands, 132

## Platinum

Synthesis of highly ethynylated mono and dinuclear Pt(II) tethers bearing the 4,4'-bis(ethynyl)biphenyl (debp) unit as central core, 49

## Platinum metal

Synthesis, spectroscopic and structural characterisation of complexes of the new ditelluroether 1,2-bis(methyltelluromethyl)benzene, *o*-C<sub>6</sub>H<sub>4</sub>(CH<sub>2</sub>TeMe)<sub>2</sub>. X-ray structures of [Mn(CO)<sub>3</sub>Cl{*o*-C<sub>6</sub>H<sub>4</sub>(CH<sub>2</sub>TeMe)<sub>2</sub>}] and [W(CO)<sub>4</sub>{*o*-C<sub>6</sub>H<sub>4</sub>(CH<sub>2</sub>TeMe)<sub>2</sub>}], 218

*P,N* ligands

Funktionelle Phosphane: Part XI. Optisch reine β-Aminophosphane und β-Aminophosphinite für die komplexkatalysierte Reduktion organischer Carbonylverbindungen. Molekülstrukturen von [(1*R*,2*R*)-Ph<sub>2</sub>PCH(Ph)CH(Me)-NH<sub>2</sub>Me]Cl, (1*R*,2*S*)-Ph<sub>2</sub>PCH(Ph)CH(Me)NHSO<sub>2</sub>Me und [{(1*R*,2*R*)-Ph<sub>2</sub>PCH(Ph)CH(Me)NHMe-κ*N*,κ*P*}Rh(η<sup>4</sup>-1,5-C<sub>8</sub>H<sub>12</sub>)]BF<sub>4</sub>, 88

## Reduction

Aluminium zirconium (+ 3 and + 4) heterometallic hydrido complexes of compositions [(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>Zr(μ-H)]<sub>2</sub>(μ-H)AlCl<sub>2</sub> and [(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>ZrH(μ-H)]<sub>2</sub>Al, 36

## Rhodium

A shortcut hydroformylation route to 7-formyl-5,6-dihydroindolizine from 1-allyl-2-formylpyrrole, 241

Funktionelle Phosphane: Part XI. Optisch reine β-Aminophosphane und β-Aminophosphinite für die komplexkatalysierte Reduktion organischer Carbonylverbindungen. Molekülstrukturen von [(1*R*,2*R*)-Ph<sub>2</sub>PCH(Ph)CH(Me)-NH<sub>2</sub>Me]Cl, (1*R*,2*S*)-Ph<sub>2</sub>PCH(Ph)CH(Me)NHSO<sub>2</sub>Me und [{(1*R*,2*R*)-Ph<sub>2</sub>PCH(Ph)CH(Me)NHMe-κ*N*,κ*P*}Rh(η<sup>4</sup>-1,5-C<sub>8</sub>H<sub>12</sub>)]BF<sub>4</sub>, 88

Syntheses and ligand properties of 1,3-*dit*-butyl-2,3-dihydro-1,3-diboroles. Rhodium-(η<sup>5</sup>-2,3-dihydro-1,3-diborole) and rhodium-(η<sup>5</sup>-2,3-dihydro-1,3-diboroly) complexes, 7

Synthesis of rhodium(III) complexes with the chiral phosphine (*S*)-Ph<sub>2</sub>PCH<sub>2</sub>CHMeCH<sub>2</sub>OH. Crystal structure of (S<sub>Rh</sub>,S<sub>C</sub>)-[(η<sup>5</sup>-C<sub>5</sub>Me<sub>5</sub>)RhCl(η<sup>2</sup>-PPh<sub>2</sub>CH<sub>2</sub>CHMeCH<sub>2</sub>OH-P,O)]BF<sub>4</sub>, 1

## Rhodium–amine complexes

Organometallic rhodium (I) complexes with 1-alkylaminopyrazole ligands, 14

## Rhodium complexes

Organometallic rhodium (I) complexes with 1-alkylaminopyrazole ligands, 14

## Ruthenium

Stabilisation of alkynyl dithiocarboxylates at a ruthenium centre: X-ray crystal structures of [Ru(H)(CO)(S<sub>2</sub>CC≡CPh)(PPh<sub>3</sub>)<sub>2</sub>], [Ru(CO)(CPh=CHPh)(S<sub>2</sub>CC≡CPh)(PPh<sub>3</sub>)<sub>2</sub>] and [Ru(CO)(C≡CMes)(S<sub>2</sub>CC≡CPh)(PPh<sub>3</sub>)<sub>2</sub>], 209

Structural and spectroscopic characterization of ruthenium(II) complexes with methyl, formyl, and acetyl groups as model species in multi-step CO<sub>2</sub> reduction, 299

## Ruthenium(II)

A new class of triazepine–ruthenium(II) complexes: synthesis, crystal structure and NMR spectroscopic properties, 265

## Sandwich complexes

Multiply trimethylstannyl substituted ferrocenes synthesis, NMR studies, X-ray structural analysis and electrochemistry, 74

## Sandwich compounds

Syntheses and ligand properties of 1,3-*dit*-butyl-2,3-dihydro-1,3-diboroles. Rhodium-(η<sup>5</sup>-2,3-dihydro-1,3-diborole) and rhodium-(η<sup>5</sup>-2,3-dihydro-1,3-diboroly) complexes, 7

## Selectivity

Preparations and selectivities of magnesium-based mixed cuprates, 31

## Silicon

Disilene R<sup>\*</sup>XSi=SiXR<sup>\*</sup> (R<sup>\*</sup>=*Sit*Bu<sub>3</sub>) mit siliciumgebundenen H- und Hal-Atomen X: Bildung, Isomerisierung, Reaktionen, 110

Modelling nucleophilic substitution at silicon using hypervalent silicon compounds based on urea ligands, 132

## Silver(I)

Donor-funktionalisierte Alkinyne von Titan(IV), Kupfer(I) und Silber(I); Festkörperstrukturen von [Ti](C≡CCH<sub>2</sub>NMe<sub>2</sub>)<sub>2</sub> und {[Ti](C≡C<sup>*Bu*</sup>)<sub>2</sub>}CuC≡CC≡CC<sub>2</sub>H<sub>5</sub>, 99

## Silylenes

Disilene R<sup>\*</sup>XSi=SiXR<sup>\*</sup> (R<sup>\*</sup>=*Sit*Bu<sub>3</sub>) mit siliciumgebundenen H- und Hal-Atomen X: Bildung, Isomerisierung, Reaktionen, 110

## Solid-state NMR

Synthesis, solution and solid-state NMR spectra of methyl 4,6-*O*-benzylidene-2,3-bis-*O*-(organostannylmethyl)-α-*D*-glucopyranosides: Crystal structure of methyl 4,6-*O*-benzylidene-2,3-bis-*O*-[(dimethylphenylstannyl)methyl]-α-*D*-glucopyranoside, 226

## Styrene

New catalysts for the alternating copolymerization of 4-*tert*-butylstyrene/CO, 287

## Substituted cyclopentadiene

New trisubstituted cyclopentadienyl ligands: synthesis, characterisation and catalytic properties of mono and dinuclear cobalt, rhodium, iron and ruthenium complexes, 179

## Substitution

Modelling nucleophilic substitution at silicon using hypervalent silicon compounds based on urea ligands, 132

## Sulphonates

Structural studies of diorganotin(IV) sulphonates. The crystal structures of [(*n*-C<sub>4</sub>H<sub>9</sub>)<sub>2</sub>Sn{μ-OSO<sub>2</sub>C<sub>6</sub>H<sub>2</sub>(CH<sub>3</sub>)<sub>3</sub>}<sub>2</sub>]<sub>*n*</sub> and its partially hydrolyzed product [(*n*-C<sub>4</sub>H<sub>9</sub>)<sub>2</sub>Sn{(μ-OH)(μ-OSO<sub>2</sub>C<sub>6</sub>H<sub>2</sub>(CH<sub>3</sub>)<sub>3</sub>)}]<sub>*n*</sub>, 157

## Synthesis

Synthesis and structural characterization of several tris(2-methyl-2-phenylpropyl)tin carboxylates containing germanium, 43

## Telluroether

Synthesis, spectroscopic and structural characterisation of complexes of the new ditelluroether 1,2-bis(methyltelluro-methyl)benzene,  $o$ -C<sub>6</sub>H<sub>4</sub>(CH<sub>2</sub>TeMe)<sub>2</sub>. X-ray structures of [Mn(CO)<sub>3</sub>Cl{*o*-C<sub>6</sub>H<sub>4</sub>(CH<sub>2</sub>TeMe)<sub>2</sub>}] and [W(CO)<sub>4</sub>{*o*-C<sub>6</sub>H<sub>4</sub>(CH<sub>2</sub>TeMe)<sub>2</sub>}], 218

## Tin

Multiply trimethylstannyl substituted ferrocenes synthesis, NMR studies, X-ray structural analysis and electrochemistry, 74  
 Synthesis and X-ray crystal structures of *ansa*-complexes of titanium, zirconium, and hafnium involving methylene-bis(indenyl) ligand, 280

## Titanium

Aluminium zirconium (+ 3 and + 4) heterometallic hydrido complexes of compositions [(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>Zr(μ-H)]<sub>2</sub>(μ-H)AlCl<sub>2</sub> and [(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>ZrH(μ-H)]<sub>3</sub>Al, 36  
 Synthesis and X-ray crystal structures of *ansa*-complexes of titanium, zirconium, and hafnium involving methylene-bis(indenyl) ligand, 280

## Titanocene

Donor-funktionalisierte Alkinyne von Titan(IV), Kupfer(I) und Silber(I); Festkörperstrukturen von [Ti](C≡CCH<sub>2</sub>NMe<sub>2</sub>)<sub>2</sub> und {[Ti](C≡C'Bu)<sub>2</sub>}CuC≡CC≡CC<sub>2</sub>H<sub>5</sub>, 99

## Triazepine

A new class of triazepine–ruthenium(II) complexes: synthesis, crystal structure and NMR spectroscopic properties, 265

## Urea

Modelling nucleophilic substitution at silicon using hypervalent silicon compounds based on urea ligands, 132

## Vinyl

Stabilisation of alkynyl dithiocarboxylates at a ruthenium centre: X-ray crystal structures of [Ru(H)(CO)(S<sub>2</sub>CC≡CPh)(PPh<sub>3</sub>)<sub>2</sub>], [Ru(CO)(CPh=CHPh)(S<sub>2</sub>CC≡CPh)(PPh<sub>3</sub>)<sub>2</sub>] and [Ru(CO)(C≡CMes)(S<sub>2</sub>CC≡CPh)(PPh<sub>3</sub>)<sub>2</sub>], 209

## X-ray analyses

A new class of triazepine–ruthenium(II) complexes: synthesis, crystal structure and NMR spectroscopic properties, 265

## X-ray analysis

Donor-funktionalisierte Alkinyne von Titan(IV), Kupfer(I) und Silber(I); Festkörperstrukturen von [Ti](C≡CCH<sub>2</sub>NMe<sub>2</sub>)<sub>2</sub> und {[Ti](C≡C'Bu)<sub>2</sub>}CuC≡CC≡CC<sub>2</sub>H<sub>5</sub>, 99

## X-ray crystallography

Structural and spectroscopic characterization of ruthenium(II) complexes with methyl, formyl, and acetyl groups as model species in multi-step CO<sub>2</sub> reduction, 299

## X-ray diffraction

Synthesis and structural characterization of several tris(2-methyl-2-phenylpropyl)tin carboxylates containing germanium, 43

## X-ray structure analyses

Disilene R\*XS<sub>2</sub>–SiXR\* (R\* = Si*t*Bu<sub>3</sub>) mit siliciumgebundenen H- und Hal-Atomen X: Bildung, Isomerisierung, Reaktionen, 110

## X-ray structure analysis

Funktionelle Phosphane: Part XI. Optisch reine β-Aminophosphane und β-Aminophosphinite für die komplex-katalysierte Reduktion organischer Carbonylverbindungen. Molekülstrukturen von [(1*R*,2*R*)-Ph<sub>2</sub>PCH(Ph)CH(Me)-NH<sub>2</sub>Me]Cl, (1*R*,2*S*)-Ph<sub>2</sub>PCH(Ph)CH(Me)NHSO<sub>2</sub>Me und [{(1*R*,2*R*)-Ph<sub>2</sub>PCH(Ph)CH(Me)NHMe-κ*N*,κ*P*}Rh(η<sup>4</sup>-1,5-C<sub>8</sub>H<sub>12</sub>)]BF<sub>4</sub>, 88

## Ziegler–Natta catalyst

Benzyl group abstraction from Ti(CH<sub>2</sub>Ph)<sub>4</sub> by B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> and [Ph<sub>3</sub>C][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>] to form Ziegler–Natta and carbocationic alkene polymerization initiators, 62

## Zirconium

Aluminium zirconium (+ 3 and + 4) heterometallic hydrido complexes of compositions [(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>Zr(μ-H)]<sub>2</sub>(μ-H)AlCl<sub>2</sub> and [(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>ZrH(μ-H)]<sub>3</sub>Al, 36

Synthesis and X-ray crystal structures of *ansa*-complexes of titanium, zirconium, and hafnium involving methylene-bis(indenyl) ligand, 280