

Subject Index of Volume 621

- Air**
Dihydroxylation of olefins using air as the terminal oxidant, 70
- Alkeneinylhalogensilanes/lithium-*tert*-butyl**
Silaheterocyclen XXXVII. Die Kombination Alkeninylhalogensilane/Lithium-*tert*-Butyl als Synthesebaustein zum Aufbau von Silaheterocyclen mit kumulierten CC-Doppelbindungen, 10
- Alkoxides**
Preparation of bistrimethylsilylmethylniobiumtetrafluoride and the application of KHF_2 and $n\text{-Bu}_4\text{NHF}_2$ as fluorinating reagents, 310
- Alkylation**
Ligands with cycloalkane backbones VIII. Steric effects in the ligand accelerated enantioselective alkylation of benzaldehyde, 153
- Alkyl polyglycosides**
Synthesis of achiral, but unsymmetric, seven-membered rhodium(I)-chelates for hydrogenation in the chiral environment of alkyl polyglucoside micelles, 120
- Alkyne cyclotrimerisation**
Stoichiometric cyclotrimerisation of chiral alkynes at a ruthenium centre: preparation of optically active $(\eta^6\text{-arene})(\eta^4\text{-cycloocta-1,5-diene})\text{ruthenium}(0)$ complexes, 246
- Alkynyl complexes**
Stereoselective synthesis of chiral terminal (*E*)-1,3-enynes derived from the aldehydes (*1R*)-(–)-myrtenal and (*S*)-(–)-perillaldehyde using the alkynyl–phosphonio complex $[\text{Ru}\{\text{C}\equiv\text{CCH}_2(\text{PPh}_3)\}(\eta^5\text{-C}_9\text{H}_7)(\text{PPh}_3)_2][\text{PF}_6]$ as synthon, 39
- Allenylidene complexes**
Stereoselective synthesis of chiral terminal (*E*)-1,3-enynes derived from the aldehydes (*1R*)-(–)-myrtenal and (*S*)-(–)-perillaldehyde using the alkynyl–phosphonio complex $[\text{Ru}\{\text{C}\equiv\text{CCH}_2(\text{PPh}_3)\}(\eta^5\text{-C}_9\text{H}_7)(\text{PPh}_3)_2][\text{PF}_6]$ as synthon, 39
- Allyl**
Komplekxkatalyse: LVII. Vereinfachte Synthese des $\text{Nd}(\pi\text{-C}_3\text{H}_5)_3\text{-C}_4\text{H}_8\text{O}_2$ nach der Grignard-Methode und Darstellung der neuen Allylneodym(III)-Komplexe $[\text{Nd}(\pi\text{-C}_5\text{Me}_5)(\pi\text{-C}_3\text{H}_5)_2\text{-C}_4\text{H}_8\text{O}_2]$ und $[\text{Nd}(\pi\text{-C}_3\text{H}_5)\text{Cl}(\text{THF})_5]\text{B}(\text{C}_6\text{H}_5)_4$; THF als Präkatalysatoren für die stereospezifische Butadienpolymerisation, 327
- Aluminium**
Synthese und Charakterisierung chiraler Organogallium- und -aluminiumalkoholate mit intramolekularer Stickstoffstabilisierung. Röntgenstrukturanalyse von Dimethylgallium-(*S*)-2-dimethylamino-3-methyl-1-butanolat, Dimethylgallium-(*S*)-2-dimethylamino-3-phenyl-1-propanolat und Dimethylgallium-(*R*)-2-dimethylamino-1-butanolat, 317
- Amine alcoholate**
Synthese und Charakterisierung chiraler Organogallium- und -aluminiumalkoholate mit intramolekularer Stickstoffstabilisierung. Röntgenstrukturanalyse von Dimethylgallium-(*S*)-2-dimethylamino-3-methyl-1-butanolat, Dimethylgallium-(*S*)-2-dimethylamino-3-phenyl-1-propanolat und Dimethylgallium-(*R*)-2-dimethylamino-1-butanolat, 317
- α -Amino acids**
Metallkomplexe mit biologisch wichtigen Liganden, CXXVII: Halbsandwich-Komplexe von Ruthenium, Rhodium und Iridium sowie phosphanhaltige Palladium- und Platin(II)-Komplexe mit Schiff-Basen aus α -Aminosäureanionen und Aldehyden, 109
- Aminotroponimate**
Cyclooctatetraene complexes of yttrium and the lanthanides with chiral and achiral aminotroponimates, 277
- Amphiphiles**
Aqueous micellar and non-micellar effects during the asymmetric hydrogenation of dehydroamino acid derivatives: influence of amphiphiles on enantioselectivity and $\alpha\text{-CH/CD}$ exchange, 158
- Aqueous biphasic catalysis**
Transition metal complexes with the C_1 -symmetric diphosphines (*R*)-(*R*)-3-benzyl-2,4-bis(diphenylphosphino)pentane and (*R*)-(*R*)-3-benzyl(*p*-sulphonate)-2,4-bis(diphenylphosphino)pentane sodium salt. Applications to enantioselective catalysis in different phase systems, 26
- Arene complexes**
Stoichiometric cyclotrimerisation of chiral alkynes at a ruthenium centre: preparation of optically active $(\eta^6\text{-arene})(\eta^4\text{-cycloocta-1,5-diene})\text{ruthenium}(0)$ complexes, 246
- Arene ligand**
Catalytic hydrogenation of aromatics under biphasic conditions: isolation and structural characterisation of the cluster intermediate $[(\eta^6\text{-C}_6\text{Me}_6)_2(\eta^6\text{-C}_6\text{H}_6)\text{Ru}_3(\mu_2\text{-H})_2(\mu_2\text{-OH})(\mu_3\text{-O})]^+$, 103
- Aromatic hydrogenation**
Catalytic hydrogenation of aromatics under biphasic conditions: isolation and structural characterisation of the cluster intermediate $[(\eta^6\text{-C}_6\text{Me}_6)_2(\eta^6\text{-C}_6\text{H}_6)\text{Ru}_3(\mu_2\text{-H})_2(\mu_2\text{-OH})(\mu_3\text{-O})]^+$, 103
- Aryl ethers**
Meta, *para* and *meta*, *ortho* double *exo* nucleophilic additions of trimethylsilyl ester enolates derived from saturated and unsaturated carboxylic acids to tricarbonylchromium complexes of aryl ethers: dearomatizing cyclization to lactones, 284
- Asymmetric catalysis**
Transition metal complexes with the C_1 -symmetric diphosphines (*R*)-(*R*)-3-benzyl-2,4-bis(diphenylphosphino)pentane and (*R*)-(*R*)-3-benzyl(*p*-sulphonate)-2,4-bis(diphenylphosphino)pentane sodium salt. Applications to enantioselective catalysis in different phase systems, 26
- Asymmetric dihydroxylation**
Dihydroxylation of olefins using air as the terminal oxidant, 70
- Asymmetric hydrogenation**
Aqueous micellar and non-micellar effects during the asymmetric hydrogenation of dehydroamino acid derivatives: influence of amphiphiles on enantioselectivity and $\alpha\text{-CH/CD}$ exchange, 158

- Synthesis of achiral, but unsymmetric, seven-membered rhodium(I)-chelates for hydrogenation in the chiral environment of alkyl polyglucoside micelles, 120
- Binaphtholato aluminates
A study of Noyori's reagent, 231
- Bis(alkynyl) titanocenes
Alkinstabilisierte monomere Organo-Kupfer(I)-Verbindungen als Organyl-Transfer Reagenzien, 143
- Bisiminopyridines
Trimethylammonium[tetrakis(pentafluorophenoxy)borate], a safe way to synthesize novel co-catalysts for olefin polymerization, 184
- Bis[2,2,6,6-tetramethylpiperidyl-(1)]-disulfide
Improved synthesis of the radical source bis[2,2,6,6-tetramethylpiperidyl-(1)]-disulfide and photoinduced reactions with $M_2(CO)_{10}$ ($M = Mn, Re$), 173
- Bonding
The electronic structure and bonding in $W_2(\mu-H)_2(O^iPr)_4(DMPE)_2$ and a comparison with $Mo_2(O^iPr)_4(DMPE)_2$, 66
- Borates
Syntheses and properties of the novel co-catalysts *N,N*-dimethylanilinium- and trityl{tetrakis[4-(trifluoromethyl)-2,3,5,6-tetrafluorophenyl]borate}, 177
Trimethylammonium[tetrakis(pentafluorophenoxy)borate], a safe way to synthesize novel co-catalysts for olefin polymerization, 184
- Butadiene
Komplekxkatalyse: LVII. Vereinfachte Synthese des $Nd(\pi-C_3H_5)_3 \cdot C_4H_8O_2$ nach der Grignard-Methode und Darstellung der neuen Allylneodym(III)-Komplexe $[Nd(\pi-C_5Me_5)(\pi-C_3H_5)_2 \cdot C_4H_8O_2]$ und $[Nd(\pi-C_3H_5)Cl(THF)_5]B(C_6H_5)_4$; THF als Prækatalysatoren für die stereospezifische Butadienpolymerisation, 327
- Carbamate compounds
Insertion of carbon dioxide and isocyanide into tantalum–amide and tantalum–methyl bonds, 299
- Carbene complexes
Synthesis of chiral β -aminoalcohol-substituted carbene complexes of manganese and influence of the chiral carbene ligand on the diastereoselectivity of the CO/PR₃ exchange, 344
- Carbon dioxide
Insertion of carbon dioxide and isocyanide into tantalum–amide and tantalum–methyl bonds, 299
- Carbosilanes
Synthesis and characterization of platinum(II)-terminated dendritic carbosilanes: X-ray crystal structure of the model species $[PtCl(C_6H_3\{CH_2NMe_2\}-2-SiMe_3-5)(PPh_3)]$, 190
- Catalysis
Unbridged metallocene dichloride complexes with mono-substituted indenyl ligands and their application for the polymerization of propene, 304
- Catalyst
Komplekxkatalyse: LVII. Vereinfachte Synthese des $Nd(\pi-C_3H_5)_3 \cdot C_4H_8O_2$ nach der Grignard-Methode und Darstellung der neuen Allylneodym(III)-Komplexe $[Nd(\pi-C_5Me_5)(\pi-C_3H_5)_2 \cdot C_4H_8O_2]$ und $[Nd(\pi-C_3H_5)Cl(THF)_5]B(C_6H_5)_4$; THF als Prækatalysatoren für die stereospezifische Butadienpolymerisation, 327
- CC-double bonds
Silaheterocyclen XXXVII. Die Kombination Alkeninylhalogensilan/Lithium-*tert*-Butyl als Synthesebaustein zum Aufbau von Silaheterocyclen mit kumulierten CC-Doppelbindungen, 10
- C–H activation
Synthesis and characterization of platinum(II)-terminated dendritic carbosilanes: X-ray crystal structure of the model species $[PtCl(C_6H_3\{CH_2NMe_2\}-2-SiMe_3-5)(PPh_3)]$, 190
- Chelate ligands
Ligands with cycloalkane backbones VIII. Steric effects in the ligand accelerated enantioselective alkylation of benzaldehyde, 153
- Chiral complexes
Synthesis of chiral β -aminoalcohol-substituted carbene complexes of manganese and influence of the chiral carbene ligand on the diastereoselectivity of the CO/PR₃ exchange, 344
- Chiral ligands
Chelating behaviour of a phosphinoalcohol leading to a stable alcohol palladium complex, 218
Chiral bis(oxazoline) and pyridyl alcoholate dioxo-molybdenum(VI) complexes: synthesis, characterization and catalytic examinations, 207
- Cluster
Catalytic hydrogenation of aromatics under biphasic conditions: isolation and structural characterisation of the cluster intermediate $[(\eta^6-C_6Me_6)_2(\eta^6-C_6H_6)Ru_3(\mu_2-H)_2(\mu_2-OH)(\mu_3-O)]^+$, 103
- Co-catalyst
Syntheses and properties of the novel co-catalysts *N,N*-dimethylanilinium- and trityl{tetrakis[4-(trifluoromethyl)-2,3,5,6-tetrafluorophenyl]borate}, 177
- Co-catalysts
Trimethylammonium[tetrakis(pentafluorophenoxy)borate], a safe way to synthesize novel co-catalysts for olefin polymerization, 184
- Copper (I) cyanide
Self-assembly of $[Cu(CN)_4]^{3-}$ ions with cationic $\{Me_3Sn\}^+$ or $\{Me_3Sn(CH_2)_3SnMe_2\}^{2+}$ fragments in the presence of a $^tBu_4N^+$ template, 254
- Counter-ions
Synthesis of achiral, but unsymmetric, seven-membered rhodium(I)-chelates for hydrogenation in the chiral environment of alkyl polyglucoside micelles, 120
- Crystal structure
Cationic derivatives of $RhCl[P(\eta^2-C_7H_7)_3]$. An intramolecular Diels–Alder rearrangement in the tripodal ligand tri(1-cyclohepta-2,4,6-trienyl)phosphane, $P(C_7H_7)_3$, 166
- Crystal structures
Preparation and crystal structures of the complexes $(\eta^5-C_5H_5)TMS-CMe_2-\eta^5-C_{13}H_8)MCl_2$ and $[3,6-di^tBuC_{13}H_6-SiMe_2-N^tBu]MCl_2$ ($M = Hf, Zr$ or Ti): mechanistic aspects of the catalytic formation of a isotactic–syndiotactic stereoblock-type polypropylene, 267
- C_1 -symmetric diphosphines
Transition metal complexes with the C_1 -symmetric diphosphines (*R*)-(*R*)-3-benzyl-2,4-bis(diphenylphosphino)pentane and (*R*)-(*R*)-3-benzyl(*p*-sulphonate)-2,4-bis(diphenylphosphino)pentane sodium salt. Applications to enantioselective catalysis in different phase systems, 26
- 1,5-Cyclooctadiene complexes
Stoichiometric cyclotrimerisation of chiral alkynes at a ruthenium centre: preparation of optically active $(\eta^6\text{-arene})(\eta^4\text{-cycloocta-1,5-diene})$ ruthenium(0) complexes, 246
- Cyclooctatetraene
Cyclooctatetraene complexes of yttrium and the lanthanides with chiral and achiral aminotroponimines, 277
- Cyclopentadienyl ligand
Mono(cyclopentadienyl)titanium complexes containing a sulfide-bridged bis(phenolato) ligand. Molecular structure of $Ti\{2,2'-S(OC_6H_2-4-Me-6^tBu)_2\}(\eta^5-C_5H_5)Cl$, 3

Dearomatization

Meta, para and *meta, ortho* double *exo* nucleophilic additions of trimethylsilyl ester enolates derived from saturated and unsaturated carboxylic acids to tricarbonylchromium complexes of aryl ethers: dearomatizing cyclization to lactones, 284

Dendrimers

Synthesis and characterization of platinum(II)-terminated dendritic carbosilanes: X-ray crystal structure of the model species [PtCl(C₆H₃{CH₂NMe₂}-2-SiMe₃-5)(PPh₃)], 190

Deuteration

Aqueous micellar and non-micellar effects during the asymmetric hydrogenation of dehydroamino acid derivatives: influence of amphiphiles on enantioselectivity and α -CH/CD exchange, 158

Diastereoselectivity

Diastereoselective oxidative addition of cyclic thiosulfates to platinum(0) compounds: chiral platinum(II) complexes with sulfenato ligands Crystal structures of S(O)-CH₂-CH(OAc)-CH(OAc)-CH₂-S and dppePt-[S(O)-(CH₂)₄-S]; Part 16. Metal complexes of functionalized sulfur-containing ligands, 352

Synthesis of chiral β -aminoalcohol-substituted carbene complexes of manganese and influence of the chiral carbene ligand on the diastereoselectivity of the CO/PR₃ exchange, 344

Diels–Alder cycloaddition2

Cationic derivatives of RhCl[P(η^2 -C₇H₇)₃]. An intramolecular Diels–Alder rearrangement in the tripodal ligand tri(1-cyclohepta-2,4,6-trienyl)phosphane, P(C₇H₇)₃, 166

Diene complexes

'*Supine*' or '*prone*' ligands: geometric preference of conjugated diene, 1-azadiene, and 1,4-diazadiene ligands on half-metallocene complexes of early transition metals, 224

Dienol

Meta, para and *meta, ortho* double *exo* nucleophilic additions of trimethylsilyl ester enolates derived from saturated and unsaturated carboxylic acids to tricarbonylchromium complexes of aryl ethers: dearomatizing cyclization to lactones, 284

Diethylzinc

Ligands with cycloalkane backbones VIII. Steric effects in the ligand accelerated enantioselective alkylation of benzaldehyde, 153

Ditopic ligands

Chelating behaviour of a phosphinoalcohol leading to a stable alcohol palladium complex, 218

Electronic structures

The electronic structure and bonding in W₂(μ -H)₂(O^{*i*}Pr)₄(DMPE)₂ and a comparison with Mo₂(O^{*i*}Pr)₄(DMPE)₂, 66

Enantioselective catalysis

Ligands with cycloalkane backbones VIII. Steric effects in the ligand accelerated enantioselective alkylation of benzaldehyde, 153

Enantioselective hydroformylation

Highly efficient enantioselective catalysis in supercritical carbon dioxide using the perfluoroalkyl-substituted ligand (*R,S*)-3-H²F⁶-BINAPHOS, 130

Enantioselective hydrogenation

Highly efficient enantioselective catalysis in supercritical carbon dioxide using the perfluoroalkyl-substituted ligand (*R,S*)-3-H²F⁶-BINAPHOS, 130

Tunable ferrocenyl diphosphine ligands for the Ir-catalyzed enantioselective hydrogenation of *N*-aryl imines, 34

1,3-Enynes

Stereoselective synthesis of chiral terminal (*E*)-1,3-enynes derived from the aldehydes (*1R*)-(–)-myrtenal and (*S*)-(–)-peril-

aldehyde using the alkylnyl–phosphonio complex [Ru{C=CCH₂(PPh₃)}(η^5 -C₉H₇)(PPh₃)₂][PF₆] as synthon, 39

Epoxidation

Chiral bis(oxazoline) and pyridyl alcoholate dioxo-molybdenum(VI) complexes: synthesis, characterization and catalytic examinations, 207

1,1'-Ferrocenedialkoxysilanes

Synthesis of poly(ferrocenyldialkoxysilane) polymers and their introduction into oxide matrices, 46

Ferrocenyl diphosphines

Tunable ferrocenyl diphosphine ligands for the Ir-catalyzed enantioselective hydrogenation of *N*-aryl imines, 34

Fluorinated ligands

Highly efficient enantioselective catalysis in supercritical carbon dioxide using the perfluoroalkyl-substituted ligand (*R,S*)-3-H²F⁶-BINAPHOS, 130

Fluorinating reagents

Preparation of bistrimethylsilylmethylniobiumtetrafluoride and the application of KHF₂ and *n*-Bu₄NHF₂ as fluorinating reagents, 310

Functional phosphines

Chelating behaviour of a phosphinoalcohol leading to a stable alcohol palladium complex, 218

Gallium

Synthese und Charakterisierung chiraler Organogallium- und -aluminiumalkoholate mit intramolekularer Stickstoffstabilisierung. Röntgenstrukturanalyse von Dimethylgallium-(*S*)-2-dimethylamino-3-methyl-1-butanolat, Dimethylgallium-(*S*)-2-dimethylamino-3-phenyl-1-propanolat und Dimethylgallium-(*R*)-2-dimethylamino-1-butanolat, 317

Group 4 and 5 metals

Preparation of bistrimethylsilylmethylniobiumtetrafluoride and the application of KHF₂ and *n*-Bu₄NHF₂ as fluorinating reagents, 310

Hafnium

Preparation and crystal structures of the complexes (η^5 -C₅H₃TMS-CMe₂- η^5 -C₁₃H₈)MCl₂ and [3,6-di^{*i*}ButC₁₃H₆-SiMe₂-N^{*i*}Bu]MCl₂ (M = Hf, Zr or Ti): mechanistic aspects of the catalytic formation of an isotactic–syndiotactic stereoblock-type polypropylene, 267

Half-metallocene fragments

'*Supine*' or '*prone*' ligands: geometric preference of conjugated diene, 1-azadiene, and 1,4-diazadiene ligands on half-metallocene complexes of early transition metals, 224

High performance liquid chromatography

High performance liquid chromatography for facile analytical separation of the enantiomers of chiral organometallic complexes, 242

Homogeneous catalysis

Highly efficient enantioselective catalysis in supercritical carbon dioxide using the perfluoroalkyl-substituted ligand (*R,S*)-3-H²F⁶-BINAPHOS, 130

Homogeneous Ziegler–Natta catalysts

Preparation and structural features of 1-(2-pyridyl)- and 1-(2-furyl)indenyl-zirconium complexes, 197

Hydridoaluminates

A study of Noyori's reagent, 231

Hydrodesulfurization

Reactions of Cp*Ir(2,5-dimethylthiophene) with electrophiles, acids, and maleic anhydride, 55

Hydroformylation

Catalytic conversions in water: Part 19. Smooth hydroformylation of *N*-allylacetamide in mono- and biphasic aqueous media, 337

Hydrogen–deuterium exchange

Aqueous micellar and non-micellar effects during the asymmetric hydrogenation of dehydroamino acid derivatives: influence of amphiphiles on enantioselectivity and α -CH/CD exchange, 158

Indenyl

Stereoselective synthesis of chiral terminal (*E*)-1,3-enynes derived from the aldehydes (1*R*)-(–)-myrtenal and (*S*)-(–)-perillaldehyde using the alkynyl–phosphonio complex $[\text{Ru}\{\text{C}\equiv\text{CCH}_2(\text{PPh}_3)\}(\eta^5\text{-C}_9\text{H}_7)(\text{PPh}_3)_2][\text{PF}_6^-]$ as synthon, 39

Indenyl lithium compounds

Preparation and structural features of 1-(2-pyridyl)- and 1-(2-furyl)indenyl-zirconium complexes, 197

Insertion reactions

Insertion of carbon dioxide and isocyanide into tantalum–amide and tantalum–methyl bonds, 299

Intramolecularly donor-stabilized

Intramolecularly donor-stabilized silenes: Part 3. Synthesis and structure of 1-[2,6-bis(dimethylaminomethyl)phenyl]-1,2,2-tris(trimethylsilyl)silene, 261

Ir-diphosphine catalysts

Tunable ferrocenyl diphosphine ligands for the Ir-catalyzed enantioselective hydrogenation of *N*-aryl imines, 34

Iridium

High performance liquid chromatography for facile analytical separation of the enantiomers of chiral organometallic complexes, 242

Metallkomplexe mit biologisch wichtigen Liganden, CXXVII: Halbsandwich-Komplexe von Ruthenium, Rhodium und Iridium sowie phosphanhaltige Palladium- und Platin(II)-Komplexe mit Schiff-Basen aus α -Aminosäureanionen und Aldehyden, 109

Iridium complexes

Reactions of $\text{Cp}^*\text{Ir}(2,5\text{-dimethylthiophene})$ with electrophiles, acids, and maleic anhydride, 55

Lanthanides

Cyclooctatetraene complexes of yttrium and the lanthanides with chiral and achiral aminotroponimines, 277

Maleic anhydride

Reactions of $\text{Cp}^*\text{Ir}(2,5\text{-dimethylthiophene})$ with electrophiles, acids, and maleic anhydride, 55

Mechanistic investigations

Aqueous micellar and non-micellar effects during the asymmetric hydrogenation of dehydroamino acid derivatives: influence of amphiphiles on enantioselectivity and α -CH/CD exchange, 158

Metal complexes

Metallkomplexe mit biologisch wichtigen Liganden, CXXVII: Halbsandwich-Komplexe von Ruthenium, Rhodium und Iridium sowie phosphanhaltige Palladium- und Platin(II)-Komplexe mit Schiff-Basen aus α -Aminosäureanionen und Aldehyden, 109

Metallocene

Syntheses and properties of the novel co-catalysts *N,N*-dimethylanilinium- and trityl{tetrakis[4-(trifluoromethyl)-2,3,5,6-tetrafluorophenyl]borate}, 177

Micelles

Aqueous micellar and non-micellar effects during the asymmetric hydrogenation of dehydroamino acid derivatives: influence of amphiphiles on enantioselectivity and α -CH/CD exchange, 158

Synthesis of achiral, but unsymmetric, seven-membered rhodium(I)-chelates for hydrogenation in the chiral environment of alkyl polyglucoside micelles, 120

Molybdenum

Chiral bis(oxazoline) and pyridyl alcoholate dioxo-molybdenum(VI) complexes: synthesis, characterization and catalytic examinations, 207

High performance liquid chromatography for facile analytical separation of the enantiomers of chiral organometallic complexes, 242

 $\text{Mo}_2(\text{O}^i\text{Pr})_4(\text{DMPE})_2$

The electronic structure and bonding in $\text{W}_2(\mu\text{-H})_2(\text{O}^i\text{Pr})_4(\text{DMPE})_2$ and a comparison with $\text{Mo}_2(\text{O}^i\text{Pr})_4(\text{DMPE})_2$, 66

N-Allylacetamide

Catalytic conversions in water: Part 19. Smooth hydroformylation of *N*-allylacetamide in mono- and biphasic aqueous media, 337

N-Aryl imines

Tunable ferrocenyl diphosphine ligands for the Ir-catalyzed enantioselective hydrogenation of *N*-aryl imines, 34

 $^n\text{Bu}_4\text{N}^+$ template

Self-assembly of $[\text{Cu}(\text{CN})_4]^{3-}$ ions with cationic $\{\text{Me}_3\text{Sn}\}^+$ or $\{\text{Me}_3\text{Sn}(\text{CH}_2)_3\text{SnMe}_2\}^{2+}$ fragments in the presence of a $^n\text{Bu}_4\text{N}^+$ template, 254

N-donor ligands

Ligands with cycloalkane backbones VIII. Steric effects in the ligand accelerated enantioselective alkylation of benzaldehyde, 153

Neodymium

Komplekxkatalyse: LVII. Vereinfachte Synthese des $\text{Nd}(\pi\text{-C}_3\text{H}_5)_3\cdot\text{C}_4\text{H}_8\text{O}_2$ nach der Grignard-Methode und Darstellung der neuen Allylneodym(III)-Komplexe $[\text{Nd}(\pi\text{-C}_5\text{Me}_5)(\pi\text{-C}_3\text{H}_5)_2\cdot\text{C}_4\text{H}_8\text{O}_2]$ und $[\text{Nd}(\pi\text{-C}_3\text{H}_5)\text{Cl}(\text{THF})_3]\text{B}(\text{C}_6\text{H}_5)_4\cdot\text{THF}$ als Präkatalysatoren für die stereospezifische Butadienpolymerisation, 327

N ligands

Cyclooctatetraene complexes of yttrium and the lanthanides with chiral and achiral aminotroponimines, 277

NMR

A study of Noyori's reagent, 231

Cationic derivatives of $\text{RhCl}[\text{P}(\eta^2\text{-C}_7\text{H}_7)_3]$. An intramolecular Diels–Alder rearrangement in the tripodal ligand tri(1-cyclohepta-2,4,6-trienyl)phosphane, $\text{P}(\text{C}_7\text{H}_7)_3$, 166

Non-hydrolytic sol–gel condensation

Synthesis of poly(ferrocenyldialkoxysilane) polymers and their introduction into oxide matrices, 46

Olefinic phosphane complexes

Cationic derivatives of $\text{RhCl}[\text{P}(\eta^2\text{-C}_7\text{H}_7)_3]$. An intramolecular Diels–Alder rearrangement in the tripodal ligand tri(1-cyclohepta-2,4,6-trienyl)phosphane, $\text{P}(\text{C}_7\text{H}_7)_3$, 166

Olefin polymerization

Syntheses and properties of the novel co-catalysts *N,N*-dimethylanilinium- and trityl{tetrakis[4-(trifluoromethyl)-2,3,5,6-tetrafluorophenyl]borate}, 177

Trimethylammonium[tetrakis(pentafluorophenoxy)borate], a safe way to synthesize novel co-catalysts for olefin polymerization, 184

Optically active complexes

Stoichiometric cyclotrimerisation of chiral alkynes at a ruthenium centre: preparation of optically active $(\eta^6\text{-arene})(\eta^4\text{-cycloocta-1,5-diene})\text{ruthenium}(0)$ complexes, 246

Optically active ligand

Synthese und Charakterisierung chiraler Organogallium- und -aluminiumalkoholate mit intramolekularer Stickstoffstabilisierung. Röntgenstrukturanalyse von Dimethyl-gallium-(*S*)-2-dimethylamino-3-methyl-1-butanolat, Dimethylgallium-(*S*)-2-dimethylamino-3-phenyl-1-propanolat und Dimethylgallium-(*R*)-2-dimethylamino-1-butanolat, 317

Organic synthesis

Alkinstabilisierte monomere Organo-Kupfer(I)-Verbindungen als Organyl-Transfer Reagenzien, 143

Organo-copper(I)

Alkinstabilisierte monomere Organo-Kupfer(I)-Verbindungen als Organyl-Transfer Reagenzien, 143

Organometallic fluorides

Preparation of bistrimethylsilylmethylniobiumtetrafluoride and the application of KHF_2 and $n\text{-Bu}_4\text{NHF}_2$ as fluorinating reagents, 310

Organotin(IV)

Self-assembly of $[\text{Cu}(\text{CN})_4]^{3-}$ ions with cationic $\{\text{Me}_3\text{Sn}\}^+$ or $\{\text{Me}_2\text{Sn}(\text{CH}_2)_3\text{SnMe}_2\}^{2+}$ fragments in the presence of a $^t\text{Bu}_4\text{N}^+$ template, 254

Osmium

Dihydroxylation of olefins using air as the terminal oxidant, 70

Oxazoline

Chiral bis(oxazoline) and pyridyl alcoholate dioxo-molybdenum(VI) complexes: synthesis, characterization and catalytic examinations, 207

Oxidation

Dihydroxylation of olefins using air as the terminal oxidant, 70

Oxidative addition

Diastereoselective oxidative addition of cyclic thiosulfates to platinum(0) compounds: chiral platinum(II) complexes with sulfenato ligands Crystal structures of $\text{S}(\text{O})\text{-(CH}_2\text{)-CH(OAc)-CH(OAc)-CH}_2\text{-S}$ and $\text{dppePt-[S(O)-(CH}_2\text{)}_4\text{-S]}$: Part 16. Metal complexes of functionalized sulfur-containing ligands, 352

Oxide complexes

Chiral bis(oxazoline) and pyridyl alcoholate dioxo-molybdenum(VI) complexes: synthesis, characterization and catalytic examinations, 207

Oxo ligand

Catalytic hydrogenation of aromatics under biphasic conditions: isolation and structural characterisation of the cluster intermediate $[(\eta^6\text{-C}_6\text{Me}_6)_2(\eta^6\text{-C}_6\text{H}_6)\text{Ru}_3(\mu_2\text{-H})_2(\mu_2\text{-OH})(\mu_3\text{-O})]^+$, 103

Palladium

Chelating behaviour of a phosphinoalcohol leading to a stable alcohol palladium complex, 218

Metallkomplexe mit biologisch wichtigen Liganden, CXXVII: Halbsandwich-Komplexe von Ruthenium, Rhodium und Iridium sowie phosphanhaltige Palladium- und Platin(II)-Komplexe mit Schiff-Basen aus α -Aminosäureanionen und Aldehyden, 109

Pentafluorophenol

Trimethylammonium[tetrakis(pentafluorophenoxy)borate], a safe way to synthesize novel co-catalysts for olefin polymerization, 184

Pentamethylcyclopentadienyl

Reactions of $\text{Cp}^*\text{Ir}(2,5\text{-dimethylthiophene)}$ with electrophiles, acids, and maleic anhydride, 55

Phenolato ligand

Mono(cyclopentadienyl)titanium complexes containing a sulfide-bridged bis(phenolato) ligand. Molecular structure of $\text{Ti}\{2,2'\text{-S}(\text{OC}_6\text{H}_2\text{-4-Me-6-}^t\text{Bu})_2\}(\eta^5\text{-C}_5\text{H}_5)\text{Cl}$, 3

Photoreactions with $\text{M}_2(\text{CO})_{10}$ (M = Mn, Re)

Improved synthesis of the radical source bis[2,2,6,6-tetramethylpiperidyl(1)]-disulfide and photoinduced reactions with $\text{M}_2(\text{CO})_{10}$ (M = Mn, Re), 173

Planar chiral

High performance liquid chromatography for facile analytical separation of the enantiomers of chiral organometallic complexes, 242

Platinum

Diastereoselective oxidative addition of cyclic thiosulfates to platinum(0) compounds: chiral platinum(II) complexes with sulfenato ligands Crystal structures of $\text{S}(\text{O})\text{-(CH}_2\text{)-CH(OAc)-CH(OAc)-CH}_2\text{-S}$ and $\text{dppePt-[S(O)-(CH}_2\text{)}_4\text{-S]}$: Part 16. Metal complexes of functionalized sulfur-containing ligands, 352

Metallkomplexe mit biologisch wichtigen Liganden, CXXVII: Halbsandwich-Komplexe von Ruthenium, Rhodium und Iridium sowie phosphanhaltige Palladium- und Platin(II)-Komplexe mit Schiff-Basen aus α -Aminosäureanionen und Aldehyden, 109

Synthesis and characterization of platinum(II)-terminated dendritic carbosilanes: X-ray crystal structure of the model species $[\text{PtCl}(\text{C}_6\text{H}_3\{\text{CH}_2\text{NMe}_2\}_2\text{-2-SiMe}_3\text{-5})(\text{PPh}_3)]$, 190

Poly(ferrocenylsilanes)

Synthesis of poly(ferrocenyldialkoxysilane) polymers and their introduction into oxide matrices, 46

Polymerization

Komplekxkatalyse: LVII. Vereinfachte Synthese des $\text{Nd}(\pi\text{-C}_3\text{H}_5)_2\text{-C}_4\text{H}_8\text{O}_2$ nach der Grignard-Methode und Darstellung der neuen Allylneodym(III)-Komplexe $[\text{Nd}(\pi\text{-C}_3\text{Me}_5)(\pi\text{-C}_3\text{H}_5)_2\text{-C}_4\text{H}_8\text{O}_2]$ und $[\text{Nd}(\pi\text{-C}_3\text{H}_5)\text{Cl}(\text{THF})_3]\text{B}(\text{C}_6\text{H}_5)_4$: THF als Prækatalysatoren für die stereospezifische Butadienpolymerisation, 327

Polymer properties

Unbridged metallocene dichloride complexes with mono-substituted indenyl ligands and their application for the polymerization of propene, 304

Polypropylene

Preparation and structural features of 1-(2-pyridyl)- and 1-(2-furyl)indenyl-zirconium complexes, 197

Potassium enolates

Meta, *para* and *meta*, *ortho* double *exo* nucleophilic additions of trimethylsilyl ester enolates derived from saturated and unsaturated carboxylic acids to tricarbonylchromium complexes of aryl ethers: dearomatizing cyclization to lactones, 284

Propene polymerization

Unbridged metallocene dichloride complexes with mono-substituted indenyl ligands and their application for the polymerization of propene, 304

2'-Pyridyl alcoholate

Chiral bis(oxazoline) and pyridyl alcoholate dioxo-molybdenum(VI) complexes: synthesis, characterization and catalytic examinations, 207

Rearrangements

Intramolecularly donor-stabilized silenes: Part 3. Synthesis and structure of 1-[2,6-bis(dimethylaminomethyl)phenyl]-1,2,2-tris(trimethylsilyl)silene, 261

Rhodium

Cationic derivatives of $\text{RhCl}[\text{P}(\eta^2\text{-C}_7\text{H}_7)_3]$. An intramolecular Diels–Alder rearrangement in the tripodal ligand tri(1-cyclohepta-2,4,6-trienyl)phosphane, $\text{P}(\text{C}_7\text{H}_7)_3$, 166

Metallkomplexe mit biologisch wichtigen Liganden, CXXVII: Halbsandwich-Komplexe von Ruthenium, Rhodium und Iridium sowie phosphanhaltige Palladium- und Platin(II)-Komplexe mit Schiff-Basen aus α -Aminosäureanionen und Aldehyden, 109

Rhodium(I) chelates

Synthesis of achiral, but unsymmetric, seven-membered rhodium(I)-chelates for hydrogenation in the chiral environment of alkyl polyglucoside micelles, 120

Rhodium/phosphine complexes

Catalytic conversions in water: Part 19. Smooth hydroformylation of *N*-allylacetamide in mono- and biphasic aqueous media, 337

Ruthenium

Catalytic hydrogenation of aromatics under biphasic conditions: isolation and structural characterisation of the cluster intermediate $[(\eta^6\text{-C}_6\text{Me}_6)_2(\eta^6\text{-C}_6\text{H}_6)\text{Ru}_3(\mu_2\text{-H})_2(\mu_2\text{-OH})(\mu_3\text{-O})]^+$, 103

Metallkomplexe mit biologisch wichtigen Liganden, CXXVII: Halbsandwich-Komplexe von Ruthenium, Rhodium und Iridium sowie phosphanhaltige Palladium- und Platin(II)-Komplexe mit Schiff-Basen aus α -Aminosäureanionen und Aldehyden, 109

Stereoselective synthesis of chiral terminal (*E*)-1,3-enynes derived from the aldehydes (1*R*)-(–)-myrtenal and (*S*)-(–)-perilaldehyde using the alkynyl–phosphonio complex $[\text{Ru}\{\text{C}\equiv\text{CCH}_2(\text{PPh}_3)\}(\eta^5\text{-C}_9\text{H}_7)(\text{PPh}_3)_2][\text{PF}_6]$ as synthon, 39

Stoichiometric cyclotrimerisation of chiral alkynes at a ruthenium centre: preparation of optically active $(\eta^6\text{-arene})(\eta^4\text{-cycloocta-1,5-diene})\text{ruthenium}(0)$ complexes, 246

Schiff bases

Metallkomplexe mit biologisch wichtigen Liganden, CXXVII: Halbsandwich-Komplexe von Ruthenium, Rhodium und Iridium sowie phosphanhaltige Palladium- und Platin(II)-Komplexe mit Schiff-Basen aus α -Aminosäureanionen und Aldehyden, 109

Self assembly

Self-assembly of $[\text{Cu}(\text{CN})_4]^{3-}$ ions with cationic $\{\text{Me}_3\text{Sn}\}^+$ or $\{\text{Me}_2\text{Sn}(\text{CH}_2)_3\text{SnMe}_2\}^{2+}$ fragments in the presence of a $^t\text{Bu}_4\text{N}^+$ template, 254

Silaethenes

Intramolecularly donor-stabilized silenes: Part 3. Synthesis and structure of 1-[2,6-bis(dimethylaminomethyl)phenyl]-1,2,2-tris(trimethylsilyl)silene, 261

Silaheterocycles

Silaheterocyclen XXXVII. Die Kombination Alkeninylhalogensilan/Lithium-*tert*-Butyl als Synthesebaustein zum Aufbau von Silaheterocyclen mit kumulierten CC-Doppelbindungen, 10

Silanes

Intramolecularly donor-stabilized silenes: Part 3. Synthesis and structure of 1-[2,6-bis(dimethylaminomethyl)phenyl]-1,2,2-tris(trimethylsilyl)silene, 261

Silenes

Intramolecularly donor-stabilized silenes: Part 3. Synthesis and structure of 1-[2,6-bis(dimethylaminomethyl)phenyl]-1,2,2-tris(trimethylsilyl)silene, 261

Silicon

Intramolecularly donor-stabilized silenes: Part 3. Synthesis and structure of 1-[2,6-bis(dimethylaminomethyl)phenyl]-1,2,2-tris(trimethylsilyl)silene, 261

Sol–gel process

Synthesis of poly(ferrocenyldialkoxysilane) polymers and their introduction into oxide matrices, 46

Stereospecific polymerization

Preparation and crystal structures of the complexes $(\eta^5\text{-C}_5\text{H}_3\text{TMS-CMe}_2\text{-}\eta^5\text{-C}_{13}\text{H}_8)\text{MCl}_2$ and $[3,6\text{-di}^t\text{BuC}_{13}\text{H}_6\text{-SiMe}_2\text{-N}^t\text{Bu}]\text{MCl}_2$ (M = Hf, Zr or Ti): mechanistic aspects of the catalytic formation of a isotactic–syndiotactic stereoblock-type polypropylene, 267

Structure–activity correlation

Tunable ferrocenyl diphosphine ligands for the Ir-catalyzed enantioselective hydrogenation of *N*-aryl imines, 34

Substituted zirconocenes

Preparation and structural features of 1-(2-pyridyl)- and 1-(2-furyl)indenyl-zirconium complexes, 197

Substitution

Synthesis of chiral β -aminoalcohol-substituted carbene complexes of manganese and influence of the chiral carbene ligand on the diastereoselectivity of the CO/PR₃ exchange, 344

Sulfenato complexes

Diastereoselective oxidative addition of cyclic thiosulfonates to platinum(0) compounds: chiral platinum(II) complexes with sulfenato ligands Crystal structures of $\text{S}(\text{O})\text{-CH}_2\text{-CH}(\text{OAc})\text{-CH}(\text{OAc})\text{-CH}_2\text{-S}$ and $\text{dppePt}[\text{S}(\text{O})\text{-}(\text{CH}_2)_4\text{-S}]$: Part 16. Metal complexes of functionalized sulfur-containing ligands, 352

Sulfur donor

Mono(cyclopentadienyl)titanium complexes containing a sulfide-bridged bis(phenolato) ligand. Molecular structure of $\text{Ti}\{2,2'\text{-S}(\text{OC}_6\text{H}_2\text{-4-Me-6-}^t\text{Bu})_2\}(\eta^5\text{-C}_5\text{H}_5)\text{Cl}$, 3

Supercritical carbon dioxide

Highly efficient enantioselective catalysis in supercritical carbon dioxide using the perfluoroalkyl-substituted ligand (*R,S*)-3-H²F⁶-BINAPHOS, 130

Supine and *prone*

'*Supine*' or '*prone*' ligands: geometric preference of conjugated diene, 1-azadiene, and 1,4-diazadiene ligands on half-metallocene complexes of early transition metals, 224

Synthesis

Chelating behaviour of a phosphinoalcohol leading to a stable alcohol palladium complex, 218

Thiolato complexes

Diastereoselective oxidative addition of cyclic thiosulfonates to platinum(0) compounds: chiral platinum(II) complexes with sulfenato ligands Crystal structures of $\text{S}(\text{O})\text{-CH}_2\text{-CH}(\text{OAc})\text{-CH}(\text{OAc})\text{-CH}_2\text{-S}$ and $\text{dppePt}[\text{S}(\text{O})\text{-}(\text{CH}_2)_4\text{-S}]$: Part 16. Metal complexes of functionalized sulfur-containing ligands, 352

Thiophene

Reactions of Cp*Ir(2,5-dimethylthiophene) with electrophiles, acids, and maleic anhydride, 55

Thiosulfonates

Diastereoselective oxidative addition of cyclic thiosulfonates to platinum(0) compounds: chiral platinum(II) complexes with sulfenato ligands Crystal structures of $\text{S}(\text{O})\text{-CH}_2\text{-CH}(\text{OAc})\text{-CH}(\text{OAc})\text{-CH}_2\text{-S}$ and $\text{dppePt}[\text{S}(\text{O})\text{-}(\text{CH}_2)_4\text{-S}]$: Part 16. Metal complexes of functionalized sulfur-containing ligands, 352

Titanium complexes

Mono(cyclopentadienyl)titanium complexes containing a sulfide-bridged bis(phenolato) ligand. Molecular structure of $\text{Ti}\{2,2'\text{-S}(\text{OC}_6\text{H}_2\text{-4-Me-6-}^t\text{Bu})_2\}(\eta^5\text{-C}_5\text{H}_5)\text{Cl}$, 3

Transfer reagents

Alkinstabilisierte monomere Organo-Kupfer(I)-Verbindungen als Organyl-Transfer Reagenzien, 143

Tricarbonylchromium complexes

Meta, *para* and *meta*, *ortho* double *exo* nucleophilic additions of trimethylsilyl ester enolates derived from saturated and unsaturated carboxylic acids to tricarbonylchromium complexes of aryl ethers: dearomatizing cyclization to lactones, 284

Trimethylene tethers

Self-assembly of $[\text{Cu}(\text{CN})_4]^{3-}$ ions with cationic $\{\text{Me}_3\text{Sn}\}^+$ or $\{\text{Me}_2\text{Sn}(\text{CH}_2)_3\text{SnMe}_2\}^{2+}$ fragments in the presence of a $^t\text{Bu}_4\text{N}^+$ template, 254

Trinuclear

Catalytic hydrogenation of aromatics under biphasic conditions: isolation and structural characterisation of the cluster intermediate $[(\eta^6\text{-C}_6\text{Me}_6)_2(\eta^6\text{-C}_6\text{H}_6)\text{Ru}_3(\mu_2\text{-H})_2(\mu_2\text{-OH})(\mu_3\text{-O})]^+$, 103

Tripodal ligands

Cationic derivatives of $\text{RhCl}[\text{P}(\eta^2\text{-C}_7\text{H}_7)_3]$. An intramolecular Diels–Alder rearrangement in the tripodal ligand tri(1-cyclohepta-2,4,6-trienyl)phosphane, $\text{P}(\text{C}_7\text{H}_7)_3$, 166

Unbridged metallocene complexes

Unbridged metallocene dichloride complexes with mono-substituted indenyl ligands and their application for the polymerization of propene, 304

Unsymmetric bisphosphines

Synthesis of achiral, but unsymmetric, seven-membered rhodium(I)-chelates for hydrogenation in the chiral environment of alkyl polyglucoside micelles, 120

Vinylidene complexes

Stereoselective synthesis of chiral terminal (*E*)-1,3-enynes derived from the aldehydes (1*R*)-(–)-myrtenal and (*S*)-(–)-perilaldehyde using the alkynyl–phosphonio complex $[\text{Ru}\{\text{C}\equiv\text{CCH}_2(\text{PPh}_3)\}(\eta^5\text{-C}_9\text{H}_7)(\text{PPh}_3)_2][\text{PF}_6]$ as synthon, 39

Water-soluble catalyst

Catalytic hydrogenation of aromatics under biphasic conditions: isolation and structural characterisation of the cluster intermediate $[(\eta^6\text{-C}_6\text{Me}_6)_2(\eta^6\text{-C}_6\text{H}_6)\text{Ru}_3(\mu_2\text{-H})_2(\mu_2\text{-OH})(\mu_3\text{-O})]^+$, 103

Wittig reactions

Stereoselective synthesis of chiral terminal (*E*)-1,3-enynes derived from the aldehydes (1*R*)-(–)-myrtenal and (*S*)-(–)-perilaldehyde using the alkynyl–phosphonio complex

$[\text{Ru}\{\text{C}\equiv\text{CCH}_2(\text{PPh}_3)\}(\eta^5\text{-C}_9\text{H}_7)(\text{PPh}_3)_2][\text{PF}_6]$ as synthon, 39

X-ray structure

Synthese und Charakterisierung chiraler Organogallium- und -aluminiumalkoholate mit intramolekularer Stickstoffstabilisierung. Röntgenstrukturanalyse von Dimethylgallium-(*S*)-2-dimethylamino-3-methyl-1-butanolat, Dimethylgallium-(*S*)-2-dimethylamino-3-phenyl-1-propanolat und Dimethylgallium-(*R*)-2-dimethylamino-1-butanolat, 317

X-ray structures

A study of Noyori's reagent, 231

Chelating behaviour of a phosphinoalcohol leading to a stable alcohol palladium complex, 218

Yttrium

Cyclooctatetraene complexes of yttrium and the lanthanides with chiral and achiral aminotroponimines, 277

Zirconium

Preparation and crystal structures of the complexes $(\eta^5\text{-C}_5\text{H}_3\text{TMS-CMe}_2\text{-}\eta^5\text{-C}_{13}\text{H}_8)\text{MCl}_2$ and $[3,6\text{-di}^i\text{ButC}_{13}\text{H}_6\text{-SiMe}_2\text{-N}^i\text{Bu}]\text{MCl}_2$ ($\text{M} = \text{Hf, Zr or Ti}$): mechanistic aspects of the catalytic formation of a isotactic–syndiotactic stereoblock-type polypropylene, 267

Syntheses and properties of the novel co-catalysts *N,N*-dimethylanilinium- and trityl{tetrakis[4-(trifluoromethyl)-2,3,5,6-tetrafluorophenyl]borate}, 177