

Subject Index of Volume 623

Alkynes

The reaction of ethynylferrocene with Cp*Rh half-sandwich complexes containing a chelating 1,2-dicarba-*closo*-dodecaborane(12)-1,2-dichalcogenolato ligand, Cp*Rh[E₂C₂-(B₁₀H₁₀)], E = S, Se, 149

Aminothiolate ligands

Schiff bases derived from mercury(II)-aminothiolate complexes as metalloligands for transition metals, 137

Aromatic diselenides

The oxidative addition of diphenyl diselenide and ditelluride to tetrakis(triphenylphosphine)palladium, 168

Aromatic ditellurides

The oxidative addition of diphenyl diselenide and ditelluride to tetrakis(triphenylphosphine)palladium, 168

Benzylselenolate

Synthesis and characterization of benzylselenolate complexes of palladium(II) and platinum(II), 48

B–H Activation

The reaction of ethynylferrocene with Cp*Rh half-sandwich complexes containing a chelating 1,2-dicarba-*closo*-dodecaborane(12)-1,2-dichalcogenolato ligand, Cp*Rh[E₂C₂-(B₁₀H₁₀)], E = S, Se, 149

Binuclear compounds

Synthesis and characterization of benzylselenolate complexes of palladium(II) and platinum(II), 48

Cadmium sulfide

Transition metal complexes of 2,4,6-trimercapto-1,3,5-triazine (TMT): potential precursors to nanoparticulate metal sulfides, 185

Cadmium telluride

Preparation of CdTe coatings using the chemical deposition method, 81

Carborane

The reaction of ethynylferrocene with Cp*Rh half-sandwich complexes containing a chelating 1,2-dicarba-*closo*-dodecaborane(12)-1,2-dichalcogenolato ligand, Cp*Rh[E₂C₂-(B₁₀H₁₀)], E = S, Se, 149

Carboxylic esters

Synthesis of α -phenylseleno- α,β -unsaturated esters by Wittig-type reactions. Studies on the Diels–Alder reaction, 131

Catalysis

The reaction of ethynylferrocene with Cp*Rh half-sandwich complexes containing a chelating 1,2-dicarba-*closo*-dodecaborane(12)-1,2-dichalcogenolato ligand, Cp*Rh[E₂C₂-(B₁₀H₁₀)], E = S, Se, 149

Chalcogen

Hypervalent organotin(IV) derivatives containing [2-(Me₂NCH₂)-C₆H₄]₂Sn moieties. Competition between nitrogen and chalcogen atoms for coordination to the metal centre, 161

Cluster

Formation of rhodium carbonyl thiolate dimers via elimination of dihydrogen; crystal and molecular structure of [Rh₂(CO)₄(μ -SC₆H₄CH₃)₂], 65

Synthesis and characterization of new thioether derivatives of [Os₃(CO)₁₂] and [H₂Os₃(CO)₁₀]; crystal and molecular structures of [Os₃(CO)₁₁(L)] [Et₂S, Pr₂S, S(CH₂)₅] and [Os₃(CO)₉(μ -H)(μ -X){SMe(Bu')}]} (X = H, OH), 191

CO-bridged anions

Synthesis of double and multiple butterfly Fe/E (E = S, Se, Te) cluster complexes via reactions of complex anions [(μ -RE)(μ -CO)Fe₂(CO)₆]⁻ (E = S, Se, Te). Crystal structures of (μ -*p*-MeC₆H₄Se)(μ -MeS)[Fe₂(CO)₆]₂(μ -S) and {(μ -*t*-BuS)[Fe₂(CO)₆]₂(μ -S)}₂[(μ -*p*-SCH₂)₂C₆H₄], 56

Complexes

A comparison of the coordination chemistry of Ph₂SNH and Ph₂SNCH₂CH₂CN; the preparation and X-ray crystal structures of [PPh₄][PdBr₃(Ph₂SNCH₂CH₂CN)] and *trans*-[PdBr₂(Ph₂SNCH₂CH₂CN)₂], the first fully characterised complexes of a N-substituted sulfimide, 120

Conversions

The general course of reactions and structural correlations in S–N chemistry discussed in terms of a carbon and sulfur(IV) equivalence, 3

Coupling reactions

Carbon–carbon bond formation by means of organotellurium compounds, 43

Cryptands

Synthesis of some macrocycles/bicycles from bis(*o*-formylphenyl) selenide: X-ray crystal structure of bis(*o*-formylphenyl) selenide and the first 28-membered selenium containing macrocyclic ligand, 87

Crystal and molecular structures

The key role of sulfur in thiosemicarbazone compounds. Crystal and molecular structure of [Pd{4-MeOC₆H₄C(Me)=NN=C(S)NHPPh}₂], 176

Crystal structure

Synthesis and characterization of new thioether derivatives of [Os₃(CO)₁₂] and [H₂Os₃(CO)₁₀]; crystal and molecular structures of [Os₃(CO)₁₁(L)] [Et₂S, Pr₂S, S(CH₂)₅] and [Os₃(CO)₉(μ -H)(μ -X){SMe(Bu')}]} (X = H, OH), 191

Crystal structures

Tellurium complexes with new, large-bite dithio ligands. The crystal structures of *cis*- and *trans*- [Te{Ph₂P(S)–N–P(S)(OPh)₂}₂] and [(4-MeOC₆H₄TeCl₃)₂{ μ -^{*i*}Pr₂P(S)–Fc–P(S)^{*i*}Pr₂}], 124

Detected and assumed intermediates

The general course of reactions and structural correlations in S–N chemistry discussed in terms of a carbon and sulfur(IV) equivalence, 3

Dichalcogenoimidodiphosphinates

Thiophosphorus and related ligands in coordination, organometallic and supramolecular chemistry. A personal account, 29

Diels–Alder reactions

Synthesis of α -phenylseleno- α,β -unsaturated esters by Wittig-type reactions. Studies on the Diels–Alder reaction, 131

Dithioarsinates

Thiophosphorus and related ligands in coordination, organometallic and supramolecular chemistry. A personal account, 29

Dithiocarbamates

Synthesis, spectroscopic characterization and structural studies of dialkyl dithiophosphinate and *N,N*-dialkyl dithio- and monothio-carbamate derivatives of 1-iodo-1,1,2,3,4,5-hexahydrotellurophene, 74

Dithiophosphates

Thiophosphorus and related ligands in coordination, organometallic and supramolecular chemistry. A personal account, 29

Dithiophosphinates

Synthesis, spectroscopic characterization and structural studies of dialkyl dithiophosphinate and *N,N*-dialkyl dithio- and monothio-carbamate derivatives of 1-iodo-1,1,2,3,4,5-hexahydrotellurophene, 74

Thiophosphorus and related ligands in coordination, organometallic and supramolecular chemistry. A personal account, 29

Early transition metals

Chemical applications of topology and group theory: Part 35. Non-octahedral six-coordinate tris(dithiolene) and related complexes of the early transition metals, 95

Ferrocene

Electrochemical and NMR spectroscopic studies of selenium- and tellurium-substituted ferrocenes I: ferrocenyl alkyl chalcogenides [Fe(η -C₅H₅)(η -C₅H₄ER)], 101

The reaction of ethynylferrocene with Cp*Rh half-sandwich complexes containing a chelating 1,2-dicarba-*closo*-dodecaborane(12)-1,2-dichalcogenolato ligand, Cp*Rh[E₂C₂(B₁₀H₁₀)], E = S, Se, 149

Halogenations

The general course of reactions and structural correlations in S–N chemistry discussed in terms of a carbon and sulfur(IV) equivalence, 3

Heterocycles

Bridge cleavage of [PhP(Se)(μ -Se)]₂ by 1,2-C₆H₄(EH)(E'H) (E, E' = O or NH). X-ray crystal structure of PhP(Se)-(NHC₆H₄NH-1,2), 116

Hydride complex

Reaction between the thiocarbonyl complex, Os(CS)(CO)(PPh₃)₃, and propyne: crystal structure of a new sulfur-substituted osmabenzene, 109

Hydrodesulfurization

Formation of rhodium carbonyl thiolate dimers via elimination of dihydrogen; crystal and molecular structure of [Rh₂(CO)₄(μ -SC₆H₄CH₃)₂], 65

Synthesis and characterization of new thioether derivatives of [Os₃(CO)₁₂] and [H₂Os₃(CO)₁₀]; crystal and molecular structures of [Os₃(CO)₁₁(L)] [Et₂S, Pr₂S, S(CH₂)₃] and [Os₃(CO)₉(μ -H)(μ -X){SMe(Bu^r)}] (X = H, OH), 191

Hypervalency

Tuning selenium–iodine contacts: from secondary soft–soft interactions to covalent bonds, 14

Hypervalent organotin(IV) derivatives

Hypervalent organotin(IV) derivatives containing [2-(Me₂NCH₂)-C₆H₄]Sn moieties. Competition between nitrogen and chalcogen atoms for coordination to the metal centre, 161

Iodine

Tuning selenium–iodine contacts: from secondary soft–soft interactions to covalent bonds, 14

Ion exchange

Preparation of CdTe coatings using the chemical deposition method, 81

Iron

Synthesis of double and multiple butterfly Fe/E (E = S, Se, Te) cluster complexes via reactions of complex anions [(μ -RE)(μ -CO)Fe₂(CO)₆][−] (E = S, Se, Te). Crystal structures of (μ -*p*-MeC₆H₄Se)(μ -MeS)[Fe₂(CO)₆]₂(μ -S) and {(μ -*t*-BuS)[Fe₂(CO)₆]₂(μ -S)}₂(μ -*p*-SCH₂)₂C₆H₄], 56

Macrocyclic

Synthesis of some macrocycles/bicycles from bis(*o*-formylphenyl) selenide: X-ray crystal structure of bis(*o*-formylphenyl) selenide and the first 28-membered selenium containing macrocyclic ligand, 87

Mechanism

Formation of rhodium carbonyl thiolate dimers via elimination of dihydrogen; crystal and molecular structure of [Rh₂(CO)₄(μ -SC₆H₄CH₃)₂], 65

Mercury complexes

Schiff bases derived from mercury(II)–aminothiolate complexes as metalloligands for transition metals, 137

Metal centre

Hypervalent organotin(IV) derivatives containing [2-(Me₂NCH₂)-C₆H₄]Sn moieties. Competition between nitrogen and chalcogen atoms for coordination to the metal centre, 161

Microwave heating

Synthesis of α -phenylseleno- α,β -unsaturated esters by Wittig-type reactions. Studies on the Diels–Alder reaction, 131

Monothiocarbamates

Synthesis, spectroscopic characterization and structural studies of dialkyl dithiophosphinate and *N,N*-dialkyl dithio- and monothio-carbamate derivatives of 1-iodo-1,1,2,3,4,5-hexahydrotellurophene, 74

Nanoparticulate metal sulfides

Transition metal complexes of 2,4,6-trimercapto-1,3,5-triazine (TMT): potential precursors to nanoparticulate metal sulfides, 185

New dithio ligands

Tellurium complexes with new, large-bite dithio ligands. The crystal structures of *cis*- and *trans*- [Te{Ph₂P(S)-N-P(S)(OPh)₂}₂] and [(4-MeOC₆H₄TeCl₃)₂{ μ -^{*i*}Pr₂P(S)-Fc-P(S)Pr₂}], 124

Nitriles

A comparison of the coordination chemistry of Ph₂SNH and Ph₂SNCH₂CH₂CN; the preparation and X-ray crystal structures of [PPh₄][PdBr₃(Ph₂SNCH₂CH₂CN)] and *trans*-[PdBr₂(Ph₂SNCH₂CH₂CN)₂], the first fully characterised complexes of a N-substituted sulfimide, 120

Nitrogen

Hypervalent organotin(IV) derivatives containing [2-(Me₂NCH₂)-C₆H₄]Sn moieties. Competition between nitrogen and chalcogen atoms for coordination to the metal centre, 161

NMR

Synthesis and characterization of benzylselenolate complexes of palladium(II) and platinum(II), 48

NMR spectroscopy

Electrochemical and NMR spectroscopic studies of selenium- and tellurium-substituted ferrocenes I: ferrocenyl alkyl chalcogenides [Fe(η -C₅H₅)(η -C₅H₄ER)], 101

Schiff bases derived from mercury(II)–aminothiolate complexes as metalloligands for transition metals, 137

- The oxidative addition of diphenyl diselenide and ditelluride to tetrakis(triphenylphosphine)palladium, 168
- Non-octahedral six-coordinate tris(dithiolene)
Chemical applications of topology and group theory: Part 35. Non-octahedral six-coordinate tris(dithiolene) and related complexes of the early transition metals, 95
- Organotellurium compounds
Carbon–carbon bond formation by means of organotellurium compounds, 43
- Osmabenzene
Reaction between the thiocarbonyl complex, Os(CS)(CO)(PPh₃)₃, and propyne: crystal structure of a new sulfur-substituted osmabenzene, 109
- Osmium
Reaction between the thiocarbonyl complex, Os(CS)(CO)(PPh₃)₃, and propyne: crystal structure of a new sulfur-substituted osmabenzene, 109
Synthesis and characterization of new thioether derivatives of [Os₃(CO)₁₂] and [H₂Os₃(CO)₁₀]; crystal and molecular structures of [Os₃(CO)₁₁(L)] [Et₂S, Pr₂S, S(CH₂)₅] and [Os₃(CO)₉(μ-H)(μ-X){SMe(Bu^t)}] (X = H, OH), 191
- Palladium
A comparison of the coordination chemistry of Ph₂SNH and Ph₂SNCH₂CH₂CN; the preparation and X-ray crystal structures of [PPh₄][PdBr₃(Ph₂SNCH₂CH₂CN)] and *trans*-[PdBr₂(Ph₂SNCH₂CH₂CN)₂], the first fully characterised complexes of a N-substituted sulfimide, 120
Synthesis and characterization of benzylselenolate complexes of palladium(II) and platinum(II), 48
- Palladium complexes
The oxidative addition of diphenyl diselenide and ditelluride to tetrakis(triphenylphosphine)palladium, 168
- Phosphorus
Bridge cleavage of [{PhP(Se)(μ-Se)}₂] by 1,2-C₆H₄(EH)(E'H) (E, E' = O or NH). X-ray crystal structure of Ph-P(Se)(NHC₆H₄NH-1,2), 116
- Platinum
Synthesis and characterization of benzylselenolate complexes of palladium(II) and platinum(II), 48
- Polyamine
Synthesis of some macrocycles/bicycles from bis(*o*-formylphenyl) selenide: X-ray crystal structure of bis(*o*-formylphenyl) selenide and the first 28-membered selenium containing macrocyclic ligand, 87
- Propynyl complex
Reaction between the thiocarbonyl complex, Os(CS)(CO)(PPh₃)₃, and propyne: crystal structure of a new sulfur-substituted osmabenzene, 109
- Reaction schemes
The general course of reactions and structural correlations in S–N chemistry discussed in terms of a carbon and sulfur(IV) equivalence, 3
- Rhodium
Formation of rhodium carbonyl thiolate dimers via elimination of dihydrogen; crystal and molecular structure of [Rh₂(CO)₄(μ-SC₆H₄CH₃)₂], 65
The reaction of ethynylferrocene with Cp*Rh half-sandwich complexes containing a chelating 1,2-dicarba-*closo*-dodecaborane(12)-1,2-dichalcogenolato ligand, Cp*Rh[E₂C₂-(B₁₀H₁₀)], E = S, Se, 149
- Rongalite
Preparation of CdTe coatings using the chemical deposition method, 81
- Schiff base
Schiff bases derived from mercury(II)–aminothiolate complexes as metalloligands for transition metals, 137
- S₂Cl₂-reaction
The general course of reactions and structural correlations in S–N chemistry discussed in terms of a carbon and sulfur(IV) equivalence, 3
- Selenaazamacrocyclic
Synthesis of some macrocycles/bicycles from bis(*o*-formylphenyl) selenide: X-ray crystal structure of bis(*o*-formylphenyl) selenide and the first 28-membered selenium containing macrocyclic ligand, 87
- Selenide
Synthesis of some macrocycles/bicycles from bis(*o*-formylphenyl) selenide: X-ray crystal structure of bis(*o*-formylphenyl) selenide and the first 28-membered selenium containing macrocyclic ligand, 87
- Selenium
Bridge cleavage of [{PhP(Se)(μ-Se)}₂] by 1,2-C₆H₄(EH)(E'H) (E, E' = O or NH). X-ray crystal structure of Ph-P(Se)(NHC₆H₄NH-1,2), 116
Electrochemical and NMR spectroscopic studies of selenium- and tellurium-substituted ferrocenes I: ferrocenyl alkyl chalcogenides [Fe(η-C₅H₅)(η-C₅H₄ER)], 101
Synthesis of double and multiple butterfly Fe/E (E = S, Se, Te) cluster complexes via reactions of complex anions [(μ-RE)(μ-CO)Fe₂(CO)₆][−] (E = S, Se, Te). Crystal structures of (μ-*p*-MeC₆H₄Se)(μ-MeS)[Fe₂(CO)₆]₂(μ₄-S) and {(μ-*t*-BuS)[Fe₂(CO)₆]₂(μ₄-S)}₂[(μ-*p*-SCH₂)₂C₆H₄], 56
The reaction of ethynylferrocene with Cp*Rh half-sandwich complexes containing a chelating 1,2-dicarba-*closo*-dodecaborane(12)-1,2-dichalcogenolato ligand, Cp*Rh[E₂C₂-(B₁₀H₁₀)], E = S, Se, 149
Tuning selenium–iodine contacts: from secondary soft–soft interactions to covalent bonds, 14
- Selenium and compounds
Synthesis of α-phenylseleno-α,β-unsaturated esters by Wittig-type reactions. Studies on the Diels–Alder reaction, 131
- Self-assembly
Thiophosphorus and related ligands in coordination, organometallic and supramolecular chemistry. A personal account, 29
- Semiconductors
Preparation of CdTe coatings using the chemical deposition method, 81
- S₄N₄-synthesis
The general course of reactions and structural correlations in S–N chemistry discussed in terms of a carbon and sulfur(IV) equivalence, 3
- Soft–soft interactions
Tuning selenium–iodine contacts: from secondary soft–soft interactions to covalent bonds, 14
- Structure
Synthesis, spectroscopic characterization and structural studies of dialkyl dithiophosphinate and *N,N*-dialkyl dithio- and monothio-carbamate derivatives of 1-iodo-1,1,2,3,4,5-hexahydro-tellurophene, 74
- Sulfimides
A comparison of the coordination chemistry of Ph₂SNH and Ph₂SNCH₂CH₂CN; the preparation and X-ray crystal structures of [PPh₄][PdBr₃(Ph₂SNCH₂CH₂CN)] and *trans*-[PdBr₂(Ph₂SNCH₂CH₂CN)₂], the first fully characterised complexes of a N-substituted sulfimide, 120
- Sulfur
Synthesis of double and multiple butterfly Fe/E (E = S, Se, Te) cluster complexes via reactions of complex anions [(μ-RE)(μ-CO)Fe₂(CO)₆][−] (E = S, Se, Te). Crystal structures of (μ-*p*-MeC₆H₄Se)(μ-MeS)[Fe₂(CO)₆]₂(μ₄-S) and {(μ-*t*-BuS)[Fe₂(CO)₆]₂(μ₄-S)}₂[(μ-*p*-SCH₂)₂C₆H₄], 56

- The key role of sulfur in thiosemicarbazone compounds. Crystal and molecular structure of $[\text{Pd}\{4\text{-MeOC}_6\text{H}_4\text{C}(\text{Me})=\text{NN}=\text{C}(\text{S})\text{NHPH}\}_2]$, 176
- The reaction of ethynylferrocene with Cp^*Rh half-sandwich complexes containing a chelating 1,2-dicarba-*closo*-dodecaborane(12)-1,2-dichalcogenolato ligand, $\text{Cp}^*\text{Rh}[\text{E}_2\text{C}_2\text{-}(\text{B}_{10}\text{H}_{10})]$, $\text{E} = \text{S}, \text{Se}$, 149
- Supramolecular
- Thiophosphorus and related ligands in coordination, organometallic and supramolecular chemistry. A personal account, 29
- Supramolecular structures
- Tuning selenium–iodine contacts: from secondary soft–soft interactions to covalent bonds, 14
- Tellurium
- Electrochemical and NMR spectroscopic studies of selenium- and tellurium-substituted ferrocenes I: ferrocenyl alkyl chalcogenides $[\text{Fe}(\eta\text{-C}_5\text{H}_5)(\eta\text{-C}_5\text{H}_4\text{ER})]$, 101
- Synthesis of double and multiple butterfly Fe/E ($\text{E} = \text{S}, \text{Se}, \text{Te}$) cluster complexes via reactions of complex anions $[(\mu\text{-RE})(\mu\text{-CO})\text{Fe}_2(\text{CO})_6]^-$ ($\text{E} = \text{S}, \text{Se}, \text{Te}$). Crystal structures of $(\mu\text{-}p\text{-MeC}_6\text{H}_4\text{Se})(\mu\text{-MeS})[\text{Fe}_2(\text{CO})_6]_2(\mu_4\text{-S})$ and $\{(\mu\text{-}t\text{-BuS})[\text{Fe}_2(\text{CO})_6]_2(\mu_4\text{-S})\}_2[(\mu\text{-}p\text{-SCH}_2)_2\text{C}_6\text{H}_4]$, 56
- Tellurium(II) and (IV) complexes
- Tellurium complexes with new, large-bite dithio ligands. The crystal structures of *cis*- and *trans*- $[\text{Te}\{\text{Ph}_2\text{P}(\text{S})\text{-N-P}(\text{S})(\text{OPh})_2\}_2]$ and $[(4\text{-MeOC}_6\text{H}_4\text{TeCl}_3)_2\{\mu\text{-}^i\text{Pr}_2\text{P}(\text{S})\text{-Fc-P}(\text{S})\text{Pr}_2\}]$, 124
- Tellurium(IV)
- Synthesis, spectroscopic characterization and structural studies of dialkyl dithiophosphinate and *N,N*-dialkyl dithio- and monothio-carbamate derivatives of 1-iodo-1,1,2,3,4,5-hexahydro-tellurophene, 74
- Tertiaryphosphines
- Synthesis and characterization of benzylselenolate complexes of palladium(II) and platinum(II), 48
- Tetrabromide salt
- Synthesis of some macrocycles/bicycles from bis(*o*-formylphenyl) selenide: X-ray crystal structure of bis(*o*-formylphenyl) selenide and the first 28-membered selenium containing macrocyclic ligand, 87
- Thioether
- Synthesis and characterization of new thioether derivatives of $[\text{Os}_3(\text{CO})_{12}]$ and $[\text{H}_2\text{Os}_3(\text{CO})_{10}]$; crystal and molecular structures of $[\text{Os}_3(\text{CO})_{11}(\text{L})]$ [$\text{Et}_2\text{S}, \text{Pr}_2\text{S}, \text{S}(\text{CH}_2)_5$] and $[\text{Os}_3(\text{CO})_9(\mu\text{-H})(\mu\text{-X})\{\text{SMe}(\text{Bu}')\}]$ ($\text{X} = \text{H}, \text{OH}$), 191
- Thiol
- Formation of rhodium carbonyl thiolate dimers via elimination of dihydrogen; crystal and molecular structure of $[\text{Rh}_2(\text{CO})_4(\mu\text{-SC}_6\text{H}_4\text{CH}_3)_2]$, 65
- Thiosemicarbazone compounds
- The key role of sulfur in thiosemicarbazone compounds. Crystal and molecular structure of $[\text{Pd}\{4\text{-MeOC}_6\text{H}_4\text{C}(\text{Me})=\text{NN}=\text{C}(\text{S})\text{NHPH}\}_2]$, 176
- TMT ligand
- Transition metal complexes of 2,4,6-trimercapto-1,3,5-triazine (TMT): potential precursors to nanoparticulate metal sulfides, 185
- Topology and group theory
- Chemical applications of topology and group theory: Part 35. Non-octahedral six-coordinate tris(dithiolene) and related complexes of the early transition metals, 95
- Transmetalation
- Carbon–carbon bond formation by means of organotellurium compounds, 43
- Voltammetry
- Electrochemical and NMR spectroscopic studies of selenium- and tellurium-substituted ferrocenes I: ferrocenyl alkyl chalcogenides $[\text{Fe}(\eta\text{-C}_5\text{H}_5)(\eta\text{-C}_5\text{H}_4\text{ER})]$, 101
- Wittig reaction
- Synthesis of α -phenylseleno- α,β -unsaturated esters by Wittig-type reactions. Studies on the Diels–Alder reaction, 131
- X-ray crystallography
- Schiff bases derived from mercury(II)–aminothiolate complexes as metalloligands for transition metals, 137
- The oxidative addition of diphenyl diselenide and ditelluride to tetrakis(triphenylphosphine)palladium, 168
- X-ray crystal structure
- Bridge cleavage of $\{[\text{PhP}(\text{Se})(\mu\text{-Se})]_2\}$ by 1,2- $\text{C}_6\text{H}_4(\text{EH})(\text{E}'\text{H})$ ($\text{E}, \text{E}' = \text{O}$ or NH). X-ray crystal structure of $\text{Ph-P}(\text{Se})(\text{NHC}_6\text{H}_4\text{NH-1,2})$, 116
- X-ray structures
- Synthesis of double and multiple butterfly Fe/E ($\text{E} = \text{S}, \text{Se}, \text{Te}$) cluster complexes via reactions of complex anions $[(\mu\text{-RE})(\mu\text{-CO})\text{Fe}_2(\text{CO})_6]^-$ ($\text{E} = \text{S}, \text{Se}, \text{Te}$). Crystal structures of $(\mu\text{-}p\text{-MeC}_6\text{H}_4\text{Se})(\mu\text{-MeS})[\text{Fe}_2(\text{CO})_6]_2(\mu_4\text{-S})$ and $\{(\mu\text{-}t\text{-BuS})[\text{Fe}_2(\text{CO})_6]_2(\mu_4\text{-S})\}_2[(\mu\text{-}p\text{-SCH}_2)_2\text{C}_6\text{H}_4]$, 56