

Subject Index of Volume 626

- Alcohol**
P-chiral β -aminophosphine oxides vs. β -aminophosphines as auxiliaries for ruthenium catalysed enantioselective transfer hydrogenation of arylketones, 157
- Aminocarbene**
Synthesis of chromium aminocarbene complexes of diterpenoids, 199
- Arene complexes**
Regio- and diastereoselective formation of some (η^6 -2-methylindoline)tricarbonylchromium(0) complex derivatives, 233
- Arylisocyanates**
Transition-metal-mediated activation of arylisocyanates in supercritical carbon dioxide, 227
- Asymmetric synthesis**
Regio- and diastereoselective formation of some (η^6 -2-methylindoline)tricarbonylchromium(0) complex derivatives, 233
- Bisphosphine**
Synthesis of new rhodium complexes with a hemilabile nitrogen-containing bis(phosphinite) or bis(phosphine) ligand. Application to hydroformylation of styrene, 221
- Bisphosphinite**
Synthesis of new rhodium complexes with a hemilabile nitrogen-containing bis(phosphinite) or bis(phosphine) ligand. Application to hydroformylation of styrene, 221
- Bis(pyrazol-1-yl)acetate**
Bis(pyrazol-1-yl)acetates as tripodal “scorpionate” ligands in transition metal carbonyl chemistry: syntheses, structures and reactivity of manganese and rhenium carbonyl complexes of the type $[\text{LM}(\text{CO})_3]$ (L = bpza, bdmpza), 16
- Bromomethyl**
Preparation, crystal structure determination, and properties of adducts of halogenomethyl compounds of indium with Group 16 donors, 68
- Bromopentacarbonylmanganese**
A new ligating mode for the anions of the type $[\text{R}_2\text{P}(\text{S})\text{NP}(\text{S})\text{R}_2]^-$: study of the interaction of $[\text{K}\{\text{Me}_2\text{P}(\text{S})\text{NP}(\text{S})\text{Me}_2\}]$ with the early transition halometalcarbonyl, $[\text{BrMn}(\text{CO})_5]$, 113
- Calcium**
Formation and crystal structures of a monomer–dimer pair of mono(cyclopentadienyl)calcium complexes, $[\text{C}_5(t\text{-Bu})_3\text{H}_2]_2\text{CaI}(\text{THF})_2$ and $\{[\text{C}_5(t\text{-Bu})_3\text{H}_2]_2\text{CaI}(\text{THF})\}_2$, 43
- ϵ -Caprolactone**
Syntheses of $\{(\text{MeC}_5\text{H}_4)_2\text{Ln}[\overline{\text{O}-\text{C}-\text{N}(\text{CH}_2)_4\text{CH}_2}]\}_2$ (Ln = Yb, Y, Er) and the X-ray crystal structure of the ytterbium complex, 176
- Carbene ligands**
A simple access to palladium complexes of functionalized heterocyclic carbenes, 100
- Carbonyl**
Bis(pyrazol-1-yl)acetates as tripodal “scorpionate” ligands in transition metal carbonyl chemistry: syntheses, structures and reactivity of manganese and rhenium carbonyl complexes of the type $[\text{LM}(\text{CO})_3]$ (L = bpza, bdmpza), 16
- Chirality**
P-chiral β -aminophosphine oxides vs. β -aminophosphines as auxiliaries for ruthenium catalysed enantioselective transfer hydrogenation of arylketones, 157
Synthesis and structure of $[\text{Sm}(\eta^5\text{-CpPPh}_2)(\kappa^3\text{-Tp}^{\text{Me}_2})(\text{I})(\text{THF})]$; a samarium complex with four different ligands, 76
- Chiral ligands**
Chiral dioxomolybdenum(VI) complexes for enantioselective alkene epoxidation, 1
- Chromium**
Regio- and diastereoselective formation of some (η^6 -2-methylindoline)tricarbonylchromium(0) complex derivatives, 233
Synthesis of chromium aminocarbene complexes of diterpenoids, 199
- Cluster**
Synthesis, characterisation, crystal structures and reactivity of a series of ruthenium–cobalt mixed-metal nitrido carbonyl clusters with iodide ligands, 125
- Co complexes**
Transition-metal-mediated activation of arylisocyanates in supercritical carbon dioxide, 227
- Conjugation**
Radical isomerization of 9,10-dihydro-9,10-disilaanthracene derivatives: stabilization factor of silyl radicals, 32
- 18-Crown-6 ether**
The first example of a dinuclear anionic lanthanoidocene complex: $[\text{K}(18\text{-crown-6})\{(\text{C}_{13}\text{H}_8)\text{CPh}_2(\text{C}_5\text{H}_4)\text{Nd}(\text{BH}_4)_2\}]_2\cdot\text{C}_4\text{H}_8\text{O}_2$, 171
- Crystal structure**
Reactions of metal–metal triply bonded complexes $[(\eta^5\text{-RC}_5\text{H}_4)_2\text{M}_2(\text{CO})_4]$ (M = Mo, W) with diphenyl ditelluride. Crystal structures of $[(\eta^5\text{-MeC}_5\text{H}_4)\text{Mo}_2(\text{CO})_4(\mu\text{-TePh}_3)]$ and $[(\eta^5\text{-EtO}_2\text{CC}_5\text{H}_4)_2\text{W}_2(\text{CO})_4(\mu\text{-TePh}_3)]$, 192
The synthesis of new complexes of rhenium(I) with heterotridentate [P,N,O] ligands, 181
- Crystal structure analysis**
On the chemistry of gallium: Part 19. Synthesis and crystal structure analysis of novel complexes containing Ga–FeCp(CO)₂-fragments, 82
- Crystal structures**
Formation and crystal structures of a monomer–dimer pair of mono(cyclopentadienyl)calcium complexes, $[\text{C}_5(t\text{-Bu})_3\text{H}_2]_2\text{CaI}(\text{THF})_2$ and $\{[\text{C}_5(t\text{-Bu})_3\text{H}_2]_2\text{CaI}(\text{THF})\}_2$, 43
Novel complexes *cis,cis*- $[\text{Rh}(\text{R})_2(\text{I})(\text{CO})(\text{dmb})]$ (R = Me, ^tPr; dmb = 4,4'-dimethyl-2,2'-bipyridine): synthesis, structure and photoreactivity, 118
Preparation, crystal structure determination, and properties of adducts of halogenomethyl compounds of indium with Group 16 donors, 68
The first example of a dinuclear anionic lanthanoidocene complex: $[\text{K}(18\text{-crown-6})\{(\text{C}_{13}\text{H}_8)\text{CPh}_2(\text{C}_5\text{H}_4)\text{Nd}(\text{BH}_4)_2\}]_2\cdot\text{C}_4\text{H}_8\text{O}_2$, 171
- Cyclisations**
2,2-Dihydroxy-methylcyclosiloxanes and other 2,2-difunctional methylcyclosiloxanes, 53
- Cyclopentadienyl**
Formation and crystal structures of a monomer–dimer pair of mono(cyclopentadienyl)calcium complexes, $[\text{C}_5(t\text{-Bu})_3\text{H}_2]_2\text{CaI}(\text{THF})_2$ and $\{[\text{C}_5(t\text{-Bu})_3\text{H}_2]_2\text{CaI}(\text{THF})\}_2$, 43
Synthesis and structures of doubly bridged bis(cyclopentadienyl) tetracarbonyl diiron complexes, 186

Cyclosiloxanes

2,2-Dihydroxy-methylcyclosiloxanes and other 2,2-difunctional methylcyclosiloxanes, 53

Cyclotrimerisation

Transition-metal-mediated activation of arylisocyanates in supercritical carbon dioxide, 227

1,4-Diaza-1,3-diene complexes

1,4-Diaza-1,3-diene (DAD) complexes of early transition elements. Syntheses, structures and molecular dynamics of mono- and bis(η^5 -cyclopentadienyl)titanium-, zirconium- and hafnium(DAD) complexes. Crystal- and molecular structures of CpTi(DAD)CH₂Ph, [CpTi(DAD)]₂O, CpZr[(DAD)(N \curvearrowright O)] and Cp₂Hf(DAD), 243

Dichalcogenoimidodiphosphinate anions

A new ligating mode for the anions of the type [R₂P(S)NP(S)R₂]⁻: study of the interaction of [K{Me₂P(S)NP(S)Me₂}] with the early transition halometalcarbonyl, [BrMn(CO)₅], 113

Dimeric titanium ketone complex

[(*t*-Bu-DAD)Ti(μ , η^2 -OCPh₂)₂Mg(μ -Cl)₂(THF): synthesis and molecular structure of a dimeric titanium ketone complex, 168

Dinuclear anionic lanthanocene compounds

The first example of a dinuclear anionic lanthanoidocene complex: [K(18-crown-6){(C₁₃H₈)CPh₂(C₅H₄)Nd(BH₄)₂}]₂·C₄H₈O₂, 171

Diphenylphosphinocyclopentadienyl

Synthesis and structure of [Sm(η^5 -CpPPh₂)(κ^3 -Tp^{Me2})(I)(THF)]; a samarium complex with four different ligands, 76

Diphenylphosphino groups

Chelating phosphines with low coordinated phosphorus atoms: Part I. Phosphaalkenes associated with phosphine moieties — X-ray structure of *cis*-(CO)₄Mo[*ortho*-Ph₂P-C₆H₄-CH=P(*t*-Bu₃C₆H₂)]·C₇H₁₆, 106

9,10-Disilaanthracenes

Radical isomerization of 9,10-dihydro-9,10-disilaanthracene derivatives: stabilization factor of silyl radicals, 32

Diterpenoid

Synthesis of chromium aminocarbene complexes of diterpenoids, 199

Doubly bridged complexes

Synthesis and structures of doubly bridged bis(cyclopentadienyl)tetracarbonyl diiron complexes, 186

Early transition metal

A new ligating mode for the anions of the type [R₂P(S)NP(S)R₂]⁻: study of the interaction of [K{Me₂P(S)NP(S)Me₂}] with the early transition halometalcarbonyl, [BrMn(CO)₅], 113

Electrochemistry

Oxidative reactions of “half-sandwich” ruthenium compounds: formation of cationic nitrosyl ruthenium(II) derivatives, 145

Enantioselective epoxidations

Chiral dioxomolybdenum(VI) complexes for enantioselective alkene epoxidation, 1

EPR spectra

Novel complexes *cis,cis*-[Rh(R)₂(I)(CO)(dmb)] (R = Me, ^{*i*}Pr; dmb = 4,4'-dimethyl-2,2'-bipyridine): synthesis, structure and photoreactivity, 118

EPR spectroscopy

Oxidative reactions of “half-sandwich” ruthenium compounds: formation of cationic nitrosyl ruthenium(II) derivatives, 145

Erbium

Syntheses of {(MeC₃H₄)₂Ln[$\overline{\text{O}-\text{C}-\text{N}(\text{CH}_2)_4\text{C}_2$]}₂ (Ln = Yb, Y, Er) and the X-ray crystal structure of the ytterbium complex, 176

Fe–Fe bond

Synthesis and structures of doubly bridged bis(cyclopentadienyl)tetracarbonyl diiron complexes, 186

Formamidines

Reactions of InMe₃ with isocyanides in the presence of amines: chemical and mass spectrometric evidence of unprecedented insertion into In–N bonds, 11

Gallium

On the chemistry of gallium: Part 19. Synthesis and crystal structure analysis of novel complexes containing Ga–FeCp(CO)₂-fragments, 82

Gallium cyclopentadienide derivatives

Chemistry of (tetramethylcyclopentadienyl)gallium(III) compounds, 59

Gallium–iron bond

On the chemistry of gallium: Part 19. Synthesis and crystal structure analysis of novel complexes containing Ga–FeCp(CO)₂-fragments, 82

Gas-phase chemistry

Entropy effects in gas phase ion-molecule association reactions, 24

Group 2

Formation and crystal structures of a monomer–dimer pair of mono(cyclopentadienyl)calcium complexes, [C₅(*t*-Bu)₃H₂]₂CaI(THF)₂ and {[C₅(*t*-Bu)₃H₂]₂CaI(THF)}₂, 43

Half-sandwich complexes

1,4-Diaza-1,3-diene (DAD) complexes of early transition elements. Syntheses, structures and molecular dynamics of mono- and bis(η^5 -cyclopentadienyl)titanium-, zirconium- and hafnium(DAD) complexes. Crystal- and molecular structures of CpTi(DAD)CH₂Ph, [CpTi(DAD)]₂O, CpZr[(DAD)(N \curvearrowright O)] and Cp₂Hf(DAD), 243

Heck reaction

Rigid rod molecules containing two Si(OMe)₃ or SiMe(OMe)₂ groups for hybrid materials synthesis, 92

Hemilabile amphiphilic phosphines

Palladium(II) complexes with hemilabile amphiphilic phosphines, 139

Hemilabile ligand

Synthesis of new rhodium complexes with a hemilabile nitrogen-containing bis(phosphinite) or bis(phosphine) ligand. Application to hydroformylation of styrene, 221

Heterogeneous catalysis

Chiral dioxomolybdenum(VI) complexes for enantioselective alkene epoxidation, 1

Heterotridentate

The synthesis of new complexes of rhenium(I) with heterotridentate [P,N,O] ligands, 181

Hydroformylation

Synthesis of new rhodium complexes with a hemilabile nitrogen-containing bis(phosphinite) or bis(phosphine) ligand. Application to hydroformylation of styrene, 221

Hydrogen bonding

Crystal and molecular structure of H₂C(SnPh₂OMe)₂·MeOH, 49

Hydrotris(3,5-dimethylpyrazolyl)borate

Synthesis and structure of [Sm(η^5 -CpPPh₂)(κ^3 -Tp^{Me2})(I)(THF)]; a samarium complex with four different ligands, 76

Imidazoles

A simple access to palladium complexes of functionalized heterocyclic carbenes, 100

Indium

Preparation, crystal structure determination, and properties of adducts of halogenomethyl compounds of indium with Group 16 donors, 68

Indolines

Regio- and diastereoselective formation of some (η^6 -2-methylindoline)tricarbonylchromium(0) complex derivatives, 233

Iodide

Formation and crystal structures of a monomer–dimer pair of mono(cyclopentadienyl)calcium complexes, $[\text{C}_5(t\text{-Bu})_3\text{H}_2]_2\text{CaI}(\text{THF})_2$ and $\{[\text{C}_5(t\text{-Bu})_3\text{H}_2]_2\text{CaI}(\text{THF})\}_2$, 43

Synthesis, characterisation, crystal structures and reactivity of a series of ruthenium–cobalt mixed-metal nitrido carbonyl clusters with iodide ligands, 125

Ion-molecule reactions

Entropy effects in gas phase ion-molecule association reactions, 24

Iron

On the chemistry of gallium: Part 19. Synthesis and crystal structure analysis of novel complexes containing Ga–FeCp(CO)₂-fragments, 82

Isocyanides

Reactions of InMe₃ with isocyanides in the presence of amines: chemical and mass spectrometric evidence of unprecedented insertion into In–N bonds, 11

Ketone

P-chiral β -aminophosphine oxides vs. β -aminophosphines as auxiliaries for ruthenium catalysed enantioselective transfer hydrogenation of arylketones, 157

Kinetics

Entropy effects in gas phase ion-molecule association reactions, 24

Ligand properties

Chelating phosphines with low coordinated phosphorus atoms: Part I. Phosphaalkenes associated with phosphine moieties — X-ray structure of *cis*-(CO)₄Mo[*ortho*-Ph₂P–C₆H₄–CH=P(*t*-Bu₃C₆H₂)]·C₇H₁₆, 106

Low-valent titanium

$[(t\text{-Bu-DAD})\text{Ti}(\mu, \eta^2\text{-OCPh}_2)_2]\text{Mg}(\mu\text{-Cl})_2(\text{THF})$: synthesis and molecular structure of a dimeric titanium ketone complex, 168

Manganese

Bis(pyrazol-1-yl)acetates as tripodal “scorpionate” ligands in transition metal carbonyl chemistry: syntheses, structures and reactivity of manganese and rhenium carbonyl complexes of the type $[\text{LM}(\text{CO})_3]$ (L = bpza, bdmppza), 16

Mass spectrometry

Entropy effects in gas phase ion-molecule association reactions, 24

Mesoporous materials

Chiral dioxomolybdenum(VI) complexes for enantioselective alkene epoxidation, 1

Metal carbonyl

Synthesis and structures of doubly bridged bis(cyclopentadienyl) tetracarbonyl diiron complexes, 186

Metallocenes

Entropy effects in gas phase ion-molecule association reactions, 24

Metalloligand

Synthesis and structure of $[\text{Sm}(\eta^5\text{-CpPPh}_2)(\kappa^3\text{-Tp}^{\text{Me}_2})(\text{I})(\text{THF})]$; a samarium complex with four different ligands, 76

Metal–metal triple bond

Reactions of metal–metal triply bonded complexes $[(\eta^5\text{-RC}_5\text{H}_4)_2\text{M}_2(\text{CO})_4]$ (M = Mo, W) with diphenyl ditelluride. Crystal structures of $[(\eta^5\text{-MeC}_5\text{H}_4)\text{Mo}_2(\text{CO})_4(\mu\text{-TePh})_3]$ and $[(\eta^5\text{-EtO}_2\text{CC}_5\text{H}_4)_2\text{W}_2(\text{CO})_4(\mu\text{-TePh})_2]$, 192

Methyl dimethoxysilanes

Rigid rod molecules containing two Si(OMe)₃ or SiMe(OMe)₂ groups for hybrid materials synthesis, 92

Methylenepyran carbene complexes

Synthesis of electron donor–acceptor polyunsaturated methyl-

enepyran Fischer type carbene complexes: dynamic ¹H-NMR study and solvatochromic properties, 37

Molybdenum complexes

Chiral dioxomolybdenum(VI) complexes for enantioselective alkene epoxidation, 1

Monoanionic ligands

The synthesis of new complexes of rhenium(I) with heterotridentate [P,N,O] ligands, 181

Mo(0), Pd(II), Pt(II) complexes

Chelating phosphines with low coordinated phosphorus atoms: Part I. Phosphaalkenes associated with phosphine moieties — X-ray structure of *cis*-(CO)₄Mo[*ortho*-Ph₂P–C₆H₄–CH=P(*t*-Bu₃C₆H₂)]·C₇H₁₆, 106

Mo₂Te₂ complex

Reactions of metal–metal triply bonded complexes $[(\eta^5\text{-RC}_5\text{H}_4)_2\text{M}_2(\text{CO})_4]$ (M = Mo, W) with diphenyl ditelluride. Crystal structures of $[(\eta^5\text{-MeC}_5\text{H}_4)\text{Mo}_2(\text{CO})_4(\mu\text{-TePh})_3]$ and $[(\eta^5\text{-EtO}_2\text{CC}_5\text{H}_4)_2\text{W}_2(\text{CO})_4(\mu\text{-TePh})_2]$, 192

Mo₂Te₃ complex

Reactions of metal–metal triply bonded complexes $[(\eta^5\text{-RC}_5\text{H}_4)_2\text{M}_2(\text{CO})_4]$ (M = Mo, W) with diphenyl ditelluride. Crystal structures of $[(\eta^5\text{-MeC}_5\text{H}_4)\text{Mo}_2(\text{CO})_4(\mu\text{-TePh})_3]$ and $[(\eta^5\text{-EtO}_2\text{CC}_5\text{H}_4)_2\text{W}_2(\text{CO})_4(\mu\text{-TePh})_2]$, 192

Ni complexes

Transition-metal-mediated activation of arylisocyanates in supercritical carbon dioxide, 227

Nitrido

Synthesis, characterisation, crystal structures and reactivity of a series of ruthenium–cobalt mixed-metal nitrido carbonyl clusters with iodide ligands, 125

Nitrosyls

Oxidative reactions of “half-sandwich” ruthenium compounds: formation of cationic nitrosyl ruthenium(II) derivatives, 145

NMR spectroscopy

The synthesis of new complexes of rhenium(I) with heterotridentate [P,N,O] ligands, 181

¹⁵N-NMR spectroscopy

Synthesis, characterisation, crystal structures and reactivity of a series of ruthenium–cobalt mixed-metal nitrido carbonyl clusters with iodide ligands, 125

Organogallium compounds

Chemistry of (tetramethylcyclopentadienyl)gallium(III) compounds, 59

Organotin compounds

The behaviour of polydentate hydrazonic ligands derived from 2-acetylpyridine towards organotin compounds: Part II. The monoorganotin(IV) complexes, 161

Palladium

A simple access to palladium complexes of functionalized heterocyclic carbenes, 100

Palladium(II) complexes

Palladium(II) complexes with hemilabile amphiphilic phosphines, 139

Pentacoordination

Crystal and molecular structure of H₂C(SnPh₂OMe)₂·MeOH, 49

Phosphaalkenes

Chelating phosphines with low coordinated phosphorus atoms: Part I. Phosphaalkenes associated with phosphine moieties — X-ray structure of *cis*-(CO)₄Mo[*ortho*-Ph₂P–C₆H₄–CH=P(*t*-Bu₃C₆H₂)]·C₇H₁₆, 106

Phosphine oxide

P-chiral β -aminophosphine oxides vs. β -aminophosphines as auxiliaries for ruthenium catalysed enantioselective transfer hydrogenation of arylketones, 157

Photochemistry

Novel complexes *cis,cis*-[Rh(R)₂(I)(CO)(dmb)] (R = Me, ⁱPr; dmb = 4,4'-dimethyl-2,2'-bipyridine): synthesis, structure and photoreactivity, 118

Ph₂Te₂

Reactions of metal–metal triply bonded complexes [(η⁵-RC₅H₄)₂M₂(CO)₄] (M = Mo, W) with diphenyl ditelluride. Crystal structures of [(η⁵-MeC₅H₄)Mo₂(CO)₄(μ-TePh)₃] and [(η⁵-EtO₂CC₅H₄)₂W₂(CO)₄(μ-TePh)₂], 192

³¹P-NMR spectroscopy

Palladium(II) complexes with hemilabile amphiphilic phosphines, 139

P,N,P-Ligand

Synthesis of new rhodium complexes with a hemilabile nitrogen-containing bis(phosphinite) or bis(phosphine) ligand. Application to hydroformylation of styrene, 221

Polydentate ligands

The behaviour of polydentate hydrazonic ligands derived from 2-acetylpyridine towards organotin compounds: Part II. The monoorganotin(IV) complexes, 161

Polymerization

Syntheses of {(MeC₅H₄)₂Ln[O₂C₂N(CH₂)₄CH₂]}₂ (Ln = Yb, Y, Er) and the X-ray crystal structure of the ytterbium complex, 176

'Push pull' molecules

Synthesis of electron donor–acceptor polyunsaturated methylenepyrans Fischer type carbene complexes: dynamic ¹H-NMR study and solvatochromic properties, 37

Pyridyl ligands

The behaviour of polydentate hydrazonic ligands derived from 2-acetylpyridine towards organotin compounds: Part II. The monoorganotin(IV) complexes, 161

Pyrrolidine

Reactions of InMe₃ with isocyanides in the presence of amines: chemical and mass spectrometric evidence of unprecedented insertion into In–N bonds, 11

Radical isomerization

Radical isomerization of 9,10-dihydro-9,10-disilaanthracene derivatives: stabilization factor of silyl radicals, 32

Reductive coupling

[(*t*-Bu-DAD)Ti(μ,η²-OCPh₂)₂Mg(μ-Cl)₂(THF): synthesis and molecular structure of a dimeric titanium ketone complex, 168

Resonance Raman spectra

Novel complexes *cis,cis*-[Rh(R)₂(I)(CO)(dmb)] (R = Me, ⁱPr; dmb = 4,4'-dimethyl-2,2'-bipyridine): synthesis, structure and photoreactivity, 118

Rhenium

Bis(pyrazol-1-yl)acetates as tripodal "scorpionate" ligands in transition metal carbonyl chemistry: syntheses, structures and reactivity of manganese and rhenium carbonyl complexes of the type [LM(CO)₃] (L = bpza, bdmpza), 16

The synthesis of new complexes of rhenium(I) with heterotridentate [P,N,O] ligands, 181

Rhodium–alkyl complexes

Novel complexes *cis,cis*-[Rh(R)₂(I)(CO)(dmb)] (R = Me, ⁱPr; dmb = 4,4'-dimethyl-2,2'-bipyridine): synthesis, structure and photoreactivity, 118

Rhodium complex

Synthesis of new rhodium complexes with a hemilabile nitrogen-containing bis(phosphinite) or bis(phosphine) ligand. Application to hydroformylation of styrene, 221

Ruthenium

P-chiral β-aminophosphine oxides vs. β-aminophosphines as auxiliaries for ruthenium catalysed enantioselective transfer hydrogenation of arylketones, 157

Ruthenium–carbon bond cleavage

Oxidative reactions of "half-sandwich" ruthenium compounds: formation of cationic nitrosyl ruthenium(II) derivatives, 145

Ruthenium–cobalt

Synthesis, characterisation, crystal structures and reactivity of a series of ruthenium–cobalt mixed-metal nitrido carbonyl clusters with iodide ligands, 125

Samarium

Synthesis and structure of [Sm(η⁵-CpPPh₂)(κ³-Tp^{Me2})(I)(THF)]; a samarium complex with four different ligands, 76

Scorpionate ligands

Bis(pyrazol-1-yl)acetates as tripodal "scorpionate" ligands in transition metal carbonyl chemistry: syntheses, structures and reactivity of manganese and rhenium carbonyl complexes of the type [LM(CO)₃] (L = bpza, bdmpza), 16

Silanols

2,2-Dihydroxy-methylcyclosiloxanes and other 2,2-difunctional methylcyclosiloxanes, 53

Silyl radical

Radical isomerization of 9,10-dihydro-9,10-disilaanthracene derivatives: stabilization factor of silyl radicals, 32

Spirosiloxanes

2,2-Dihydroxy-methylcyclosiloxanes and other 2,2-difunctional methylcyclosiloxanes, 53

Structure

Synthesis and structure of [Sm(η⁵-CpPPh₂)(κ³-Tp^{Me2})(I)(THF)]; a samarium complex with four different ligands, 76

Sulfur ylid

Preparation, crystal structure determination, and properties of adducts of halogenomethyl compounds of indium with Group 16 donors, 68

Sulphur ligands

The behaviour of polydentate hydrazonic ligands derived from 2-acetylpyridine towards organotin compounds: Part II. The monoorganotin(IV) complexes, 161

Supercritical CO₂

Transition-metal-mediated activation of arylisocyanates in supercritical carbon dioxide, 227

supine Conformation

1,4-Diaza-1,3-diene (DAD) complexes of early transition elements. Syntheses, structures and molecular dynamics of mono- and bis(η⁵-cyclopentadienyl)titanium-, zirconium- and hafnium(DAD) complexes. Crystal- and molecular structures of CpTi(DAD)CH₂Ph, [CpTi(DAD)]₂O, CpZr[(DAD)(N⁺O)] and Cp₂Hf(DAD), 243

Supported catalysts

Chiral dioxomolybdenum(VI) complexes for enantioselective alkene epoxidation, 1

Synthesis

Rigid rod molecules containing two Si(OMe)₃ or SiMe(OMe)₂ groups for hybrid materials synthesis, 92

Syntheses of {(MeC₅H₄)₂Ln[O₂C₂N(CH₂)₄CH₂]}₂ (Ln = Yb, Y, Er) and the X-ray crystal structure of the ytterbium complex, 176

Tetramethylcyclopentadienide derivatives

Chemistry of (tetramethylcyclopentadienyl)gallium(III) compounds, 59

Tin

Crystal and molecular structure of H₂C(SnPh₂OMe)₂·MeOH, 49

Transfer hydrogenation

P-chiral β-aminophosphine oxides vs. β-aminophosphines as auxiliaries for ruthenium catalysed enantioselective transfer hydrogenation of arylketones, 157

Transition metal complexes

A simple access to palladium complexes of functionalized heterocyclic carbenes, 100

Trimethoxysilanes

Rigid rod molecules containing two Si(OMe)₃ or SiMe(OMe)₂ groups for hybrid materials synthesis, 92

Triphenylphosphine oxide complex

Preparation, crystal structure determination, and properties of adducts of halogenomethyl compounds of indium with Group 16 donors, 68

Triphenylphosphonium-sulfur methylide

Preparation, crystal structure determination, and properties of adducts of halogenomethyl compounds of indium with Group 16 donors, 68

Tungsten and chromium Fischer type carbenes

Synthesis of electron donor–acceptor polyunsaturated methylenepyran Fischer type carbene complexes: dynamic ¹H-NMR study and solvatochromic properties, 37

W₂Te₂ complex

Reactions of metal–metal triply bonded complexes [(η⁵-RC₅H₄)₂M₂(CO)₄] (M = Mo, W) with diphenyl ditelluride. Crystal structures of [(η⁵-MeC₅H₄)Mo₂(CO)₄(μ-TePh)₃] and [(η⁵-EtO₂CC₅H₄)₂W₂(CO)₄(μ-TePh)₂], 192

X-ray analysis

A new ligating mode for the anions of the type [R₂P(S)NP(S)R₂][−]:

study of the interaction of [K{Me₂P(S)NP(S)Me₂}] with the early transition halometalcarbonyl, [BrMn(CO)₅], 113

X-ray crystal structures

2,2-Dihydroxy-methylcyclosiloxanes and other 2,2-difunctional methylcyclosiloxanes, 53

X-ray structural study

Chemistry of (tetramethylcyclopentadienyl)gallium(III) compounds, 59

X-ray structure

Chelating phosphines with low coordinated phosphorus atoms: Part I. Phosphaalkenes associated with phosphine moieties — X-ray structure of *cis*-(CO)₄Mo[*ortho*-Ph₂P-C₆H₄-CH=P(*t*-Bu₃C₆H₂)]·C₇H₁₆, 106

Crystal and molecular structure of H₂C(SnPh₂OMe)₂·MeOH, 49

Syntheses of {(MeC₅H₄)₂Ln[$\overline{\text{O}-\text{C}-\text{N}(\text{CH}_2\text{C}_4\text{CH}_2)_2}$]}₂ (Ln = Yb, Y, Er) and the X-ray crystal structure of the ytterbium complex, 176

Ytterbium

Syntheses of {(MeC₅H₄)₂Ln[$\overline{\text{O}-\text{C}-\text{N}(\text{CH}_2)_4\text{CH}_2}$]}₂ (Ln = Yb, Y, Er) and the X-ray crystal structure of the ytterbium complex, 176

Yttrium

Syntheses of {(MeC₅H₄)₂Ln[$\overline{\text{O}-\text{C}-\text{N}(\text{CH}_2)_4\text{CH}_2}$]}₂ (Ln = Yb, Y, Er) and the X-ray crystal structure of the ytterbium complex, 176