

Subject Index of Volumes 637–639

- Acetal
Synthesis of controlled π -extended conjugate nanostructures of 1,1'-ferrocene, 230
- Acetylene dicobalthexacarbonyl adducts
Semimasked 1,1'-diethynylferrocenes: synthetic concepts, preparations, and reactions, 558
- Acetylenes
Synthesis of controlled π -extended conjugate nanostructures of 1,1'-ferrocene, 230
- 1-Acetyl-1'-enonylferrocenes
Preparation of ferrocenyl mono- and dienone derivatives through aldol condensation of 1,1'-diacetylferrocene with aromatic aldehydes in dry conditions, 782
- Acyl thiourea
Synthesis and molecular structure of a novel ferrocene-containing macrocyclic acyl thiourea derivative, 204
- Alcohols
Synthesis of ferrocenylketyl radicals by chromium(II) complexes, 813
- Aldol condensation
Preparation of ferrocenyl mono- and dienone derivatives through aldol condensation of 1,1'-diacetylferrocene with aromatic aldehydes in dry conditions, 782
- Alkanethiol SAM on gold
Anion effect on the electrochemical characteristics of a gold electrode modified with a self-assembled monolayer of ferrocenylhexanethiol in aqueous and dichloromethane solutions, 841
- Alkylation
Novel ferrocene modified *P,N*-ligands for enantioselective palladium-catalyzed allylic substitution reactions, 762
Palladium-catalyzed asymmetric allylic alkylations of cycloalkenyl acetates with planar chiral phosphino-ferrocene carboxylic acids, 845
- Alkyl ferrocenes
Enzymatic chemistry of ferrocenes: micellar tuning of the glucose oxidase reactivity toward solubilized electrochemically generated *n*-alkylferrocenium cations, 469
- Alkylidyne
A comparison of ferrocenyl carbenium/non-carbenium structures in isomeric osmium cluster complexes obtained from the reaction of bis-ferrocenylbutadiyne with $\text{Os}_3(\text{CO})_{10}(\mu\text{-py})(\mu\text{-H})$, 514
- Alkynes
Internal ferrocenylalkynes—a comparative electrochemical and mass spectrometric study, 291
- Alkynyl complexes
Synthesis, structures and electrochemistry of bis(alkynylferrocene) complexes with fluorene spacers, 159
- Allenyl carbene complexes
The reaction of ferrocenyl acetylene with $[\text{RuCp}(\text{PR}_3)(\text{CH}_3\text{CN})_2]\text{-PF}_6$ (R = Me, Ph, Cy). Formation of the first allenyl carbene complexes, 70
- Amide coupling
Synthesis and dihydrogen phosphate binding properties of pyrrole containing *ansa*-ferrocenes, 343
- Amidates
Synthesis, reactivity, and crystal structures of ferrocene-substituted amidate derivatives, 521
- Amination
Novel ferrocene modified *P,N*-ligands for enantioselective palladium-catalyzed allylic substitution reactions, 762
- Amino acid derivatives
Solution phase peptide synthesis with ferrocenyl amino acid derivatives, 349
- Aminoborane
New ferrocenyl amine derivatives: *N*-silyl, *N*-stannyl and *N*-boryl ferrocenyl amines, 727
- Aminosilane
New ferrocenyl amine derivatives: *N*-silyl, *N*-stannyl and *N*-boryl ferrocenyl amines, 727
- Aminostannane
New ferrocenyl amine derivatives: *N*-silyl, *N*-stannyl and *N*-boryl ferrocenyl amines, 727
- β -Amino sulphides
Synthesis and reactivity of achiral and of a novel planar chiral thioferrocenylsilanes, 407
- Anion complexation
Complexation and electrochemical sensing of anions by amide-substituted ferrocenyl ligands, 356
- Anion effect
Anion effect on the electrochemical characteristics of a gold electrode modified with a self-assembled monolayer of ferrocenylhexanethiol in aqueous and dichloromethane solutions, 841
- Anion sensing
An electrochemical study in acetonitrile of macrocyclic or open-chain ferrocene-containing oxa-aza or polyaza receptors in the presence of protons, metal cations and anions, 151
- Antiestrogen
Studies on organometallic selective estrogen receptor modulators. (SERMs) Dual activity in the hydroxy-ferrocifen series, 500
- Applications
Studies on the cyclometallation of ferrocenylimines, 27
- Arene stacking
Conformational control of intramolecular arene stacking in ferrocene complexes bearing *tert*-butyl and pentafluorophenyl substituents, 107
- Arrangement
1,1',2,2'-Tetrakis(diphenylphosphino)-4,4'-di-*tert*-butylferrocene, a new *cisoid* arrangement of phosphino groups, 126
- Arsonic acid
Synthesis and characterisation of ferrocenyl-phosphonic and -arsonic acids, 216
- Arylene-alkynylene
A new phthalocyanine-ferrocene conjugated dyad, 751

Asymmetric catalysis

Palladium-catalyzed asymmetric allylic alkylations of cycloalkenyl acetates with planar chiral phosphino-ferrocene carboxylic acids, 845

Asymmetric synthesis

Asymmetric synthesis of planar chiral 2-mono- and 2,2'-disubstituted 1,1'-bisbenzoylferrocenes, 698

Baylis–Hillman adducts

Ferrocenyl substituted chlorostilbenes and butadienes, 616

Benzoferrocenes

Ferrocenes derived from cyclopenta[*l*]phenanthrene: dibenzindene–metal complexes that resist haptotropic shifts, 372

Biferrocene

Improved electrochemistry of multi-ferrocenyl compounds: investigation of biferrocene, terferrocene, bis(fulvalene)diiron and diferrocenylethane in dichloromethane using [NBu₄][B(C₆F₅)₄] as supporting electrolyte, 823

Binaphthyl

Study on the synthesis of nonracemic C₂-symmetric 1,1'-binaphthyl-2,2'-diyl bridged ferrocene. Stereochemical result of the cross-coupling reactions controlled by Pd(II) or Pd(IV) complex intermediacy, 318

Binding constants

Synthesis and dihydrogen phosphate binding properties of pyrrole containing *ansa*-ferrocenes, 343

Binuclear ferrocene derivatives

Control of the symmetry of binuclear ferrocene derivative by using chiral substituent and its mixed-valence state, 820

Bioelectrocatalysis

Enzymatic chemistry of ferrocenes: micellar tuning of the glucose oxidase reactivity toward solubilized electrochemically generated *n*-alkylferrocenium cations, 469

Bioorganometallic

1,1'-Ferrocenyl–oligoprolines. A synthetic, structural and electrochemical study, 335

Bioorganometallic chemistry

Structural characterization and complexation behavior of ferrocene bearing dipeptide chain (-L-Ala-L-Pro-NHPy), 75

Bio-organometallic chemistry

Solution phase peptide synthesis with ferrocenyl amino acid derivatives, 349

Bioorganometallic chemistry

Synthesis and characterization of ferrocene-labeled oligodeoxynucleotides, 398

Biosensors

Solution phase peptide synthesis with ferrocenyl amino acid derivatives, 349

Biphenyl

Synthesis of controlled π -extended conjugate nanostructures of 1,1'-ferrocene, 230

2,2'-Bipyridylboronium cations

Multistep redox properties of 2,2'-bipyridylboronium substituted ferrocenes, 390

Bis(allyl) complexes

Reactivity of the bis(allyl)-ruthenium(IV) dimer [$\{\text{Ru}(\eta^3\text{:}\eta^3\text{-C}_{10}\text{H}_{16})(\mu\text{-Cl})\text{Cl}\}_2$] (C₁₀H₁₆ = 2,7-dimethylocta-2,6-diene-1,8-diyl) towards dppf (dppf = [Fe($\eta^5\text{-C}_5\text{H}_4\text{PPh}_2$)₂]) and dppfO (dppfO = [Fe($\eta^5\text{-C}_5\text{H}_4\text{PPh}_2$)($\eta^5\text{-C}_5\text{H}_4\text{P(=O)Ph}_2$)]), 767

1,1'-bis(Diphenylphosphino)

-2-[1-((boronato-functionalised)amino)ethyl]ferrocenes

Redox behavior of boronato-functionalized 1,1'-bis(diphenylphosphino)ferrocenes, 800

1,1'-Bis(diphenylphosphino)ferrocene

Reactivity of the bis(allyl)-ruthenium(IV) dimer [$\{\text{Ru}(\eta^3\text{:}\eta^3\text{-$

C₁₀H_{16})($\mu\text{-Cl})\text{Cl}\}_2$] (C₁₀H₁₆ = 2,7-dimethylocta-2,6-diene-1,8-diyl) towards dppf (dppf = [Fe($\eta^5\text{-C}_5\text{H}_4\text{PPh}_2$)₂]) and dppfO (dppfO = [Fe($\eta^5\text{-C}_5\text{H}_4\text{PPh}_2$)($\eta^5\text{-C}_5\text{H}_4\text{P(=O)Ph}_2$)]), 767}

Synthesis, X-ray, spectroscopic and a preliminary Suzuki coupling screening studies of a complete series of dppfMX₂ (M = Pt, Pd; X = Cl, Br, I), 691

Bis(ferrocenes)

Characterization and electrochemical study of bis(ferrocenes) with a furan spacer and ferrocenophanes prepared from α -bromoacetyl substituted ferrocenes, 259

bis-(Ferrocenyl)butadiyne

A comparison of ferrocenyl carbenium/non-carbenium structures in isomeric osmium cluster complexes obtained from the reaction of bis-ferrocenylbutadiyne with Os₃(CO)₁₀($\mu\text{-py}$)($\mu\text{-H}$), 514

Bis(ferrocenyl)mercury

Bis(ferrocenyl)mercury as a source of ferrocenyl moiety in Pd-catalyzed reactions of carbon–carbon bond formation, 653

Bis(fulvalene)diiron

Improved electrochemistry of multi-ferrocenyl compounds: investigation of biferrocene, terferrocene, bis(fulvalene)diiron and diferrocenylethane in dichloromethane using [NBu₄][B(C₆F₅)₄] as supporting electrolyte, 823

Borane

Highly efficient reduction of ferrocenyl derivatives by borane, 364

 α -Bromoacetyl substituted ferrocenes

Characterization and electrochemical study of bis(ferrocenes) with a furan spacer and ferrocenophanes prepared from α -bromoacetyl substituted ferrocenes, 259

Bulky cyclopentadienyl ligand

Hindering the formation of ferrocenes: mono(cyclopentadienyl)halo iron complexes [Fe(C₅R₅)X] containing a sterically bulky cyclopentadienyl ligand, 786

Butatrienylidene

Allylferrocenylselenide and the synthesis of the first seleno-substituted allenylidene complex: synthesis, spectroscopy, electrochemistry and the effect of electron transfer from the ferrocenylselenyl subunit, 240

Carbon chains

Syntheses and some reactions of complexes containing carbon chains capped by ferrocenyl and W(CO)₃Cp groups, 484

Carbon nucleophile

($\eta^5\text{-C}_5\text{H}_5$)-Ring alkylation reaction with the C₅H₅ anion: towards the construction of tri-Fe complex Fe{ $[\mu, \eta^5\text{:}\eta^4\text{-5-}exo\text{-}(1'\text{-C}_5\text{H}_4)\text{C}_5\text{H}_5]\text{Fe}(\text{CO})_2(\text{PPh}_3)$]₂, 549

Catalysis

ortho-Silylation of 2,2'-bis(oxazolonyl)-1,1'-bis(diphenylphosphino)ferrocenes and remarkable effect of the silyl groups on the enantioselectivity in Pd-catalyzed asymmetric allylic alkylation, 99

Catalytic hydrogenation

Formation and structural characterization of amino-substituted [3]ferrocenophanes derived from intramolecular Mannich-type coupling reactions at the metallocene framework, 621

Cation sensing

An electrochemical study in acetonitrile of macrocyclic or open-chain ferrocene-containing oxa-aza or polyaza receptors in the presence of protons, metal cations and anions, 151

Cation symmetry

Control of the symmetry of binuclear ferrocene derivative by using chiral substituent and its mixed-valence state, 820

C–C bond formation

The reaction of ferrocenyl acetylene with [RuCp(PR₃)-(CH₃CN)₂]PF₆ (R = Me, Ph, Cy). Formation of the first allenyl carbene complexes, 70

- Charge transfer
 Charge transfer spectra and photoreactivity of (*E*)-1-ferrocenyl-2-(1-methyl-4-pyridiniumyl)ethylene cation, 777
- Charge-transfer complexes
 Short syntheses of 8-substituted 8'-[1-(1'-phenylthio)ferrocenyl]-1,1'-binaphthyls from Suzuki coupling reactions. A strategy for generating new chiral ligands and charge-transfer complexes, 832
- Chelate
 Synthesis and characterisation of novel multidentate ferrocene ligands and their Re(I) and Pt(II) complexes, 418
- Chiral
 Chiral ferrocenylthiazolidines, new ligands for palladium complexes, 116
- Chirality
 (Ferrocenylmethyl)trialkyl ammoniums as template cations in optically active two-dimensional oxalate bridged [Cr–Mn] and [Cr–Ni] molecule-based magnets: synthesis and magnetic properties, 182
 New chiral ferrocenyl-pyridinium salts for non-linear optics, 531
- Chiral ligand
 Octahedral ruthenium(II) complexes containing the chiral ligand (4*S*)-2-[(*S_p*)-2-(diphenylphosphino)ferrocenyl]-4-(isopropyl)oxazoline (FcPN). X-Ray crystal structures of *fac*-[RuCl₂(PMe₃)₂(FcPN)] and *fac*-[RuCl₂(dppm)(FcPN)] (dppm = bis(diphenylphosphino)methane), 463
- Chiral phosphines
 Short syntheses of 8-substituted 8'-[1-(1'-phenylthio)ferrocenyl]-1,1'-binaphthyls from Suzuki coupling reactions. A strategy for generating new chiral ligands and charge-transfer complexes, 832
- Chromium
 (Ferrocenylmethyl)trialkyl ammoniums as template cations in optically active two-dimensional oxalate bridged [Cr–Mn] and [Cr–Ni] molecule-based magnets: synthesis and magnetic properties, 182
- Chromium complexes
 Synthesis of ferrocenylketyl radicals by chromium(II) complexes, 813
- Circular dichroism
 (Ferrocenylmethyl)trialkyl ammoniums as template cations in optically active two-dimensional oxalate bridged [Cr–Mn] and [Cr–Ni] molecule-based magnets: synthesis and magnetic properties, 182
- Cisoid*
 1,1',2,2'-Tetrakis(diphenylphosphino)-4,4'-di-*tert*-butylferrocene, a new *cisoid* arrangement of phosphino groups, 126
- Cluster
 A comparison of ferrocenyl carbenium/non-carbenium structures in isomeric osmium cluster complexes obtained from the reaction of bis-ferrocenylbutadiyne with Os₃(CO)₁₀(μ-py)(μ-H), 514
 Synthesis, characterisation, crystal structures and electrochemistry of triosmium nitrite carbonyl clusters containing ferrocenyl phosphines, 276
- Cobalt
 Synthesis, reactivity, and crystal structures of ferrocene-substituted amidinate derivatives, 521
- Complexes
 The effect of d-block metal complexation on the spectroscopic and redox properties of ferrocene derivatives containing pyridine ligands, 304
- Conformational analysis
 Conformational control of intramolecular arene stacking in ferrocene complexes bearing *tert*-butyl and pentafluorophenyl substituents, 107
- π-Conjugated polymers
 Electrochemical determination of diffusion coefficient of π-conjugated polymers containing ferrocene unit, 809
- π-Conjugation
 4-Ethynyl-benzonitrile-ferrocenes bridged by a Pd(PPh₃)₂ unit; the solid-state structure of (η⁵-C₅H₅)Fe(η⁵-C₅H₄C≡CC₆H₄C≡N-1,4), 198
- Coordination
 Properties of some ferrocene macrocyclic dioxetetraamines: the roles of aromatic side-arms, 327
- Copper(I) complexes
 Copper(I) complexes containing 1,1'-bis(diphenylphosphino) ferrocene (dppf) as a chelate ligand. X-ray crystal structures of [Cu₂(μ-Cl)₂(κ²-*P,P*-dppf)₂] and [Cu(CN^tBu)₂(κ²-*P,P*-dppf)][BF₄], 677
- Coupling reactions
 Synthesis of controlled π-extended conjugate nanostructures of 1,1'-ferrocene, 230
- Cp-ring alkylation
 (η⁵-C₅H₅)-Ring alkylation reaction with the C₅H₅ anion: towards the construction of tri-Fe complex Fe{[μ,η²:η⁴-5-*exo*-(1'-C₅H₄)C₅H₅]Fe(CO)₂(PPh₃)₂}, 549
- Cross-coupling
 Study on the synthesis of nonracemic C₂-symmetric 1,1'-binaphthyl-2,2'-diyl bridged ferrocene. Stereochemical result of the cross-coupling reactions controlled by Pd(II) or Pd(IV) complex intermediacy, 318
 Synthesis and characterization of ferrocene-labeled oligodeoxynucleotides, 398
- Cross-coupling reactions
 Bis(ferrocenyl)mercury as a source of ferrocenyl moiety in Pd-catalyzed reactions of carbon–carbon bond formation, 653
- Crown thioethers
 Synthetic, electrochemical, and structural studies on heterobimetallic crown thioether complexes with Group 10 metals: the crystal structures of [Pt(9S3)(dppf)](PF₆)₂·CH₃NO₂ and [Pd(9S3)(dppf)](PF₆)₂·CH₃NO₂, 683
- Crystal engineering
 Conformational control of intramolecular arene stacking in ferrocene complexes bearing *tert*-butyl and pentafluorophenyl substituents, 107
 Organometallic building blocks for crystal engineering. Synthesis, structure and hydrogen bonding interactions in [Fe(η⁵-C₅H₄-CH₂(CH₃)OH)₂], [Fe(η⁵-C₅H₃(CH₃)COOH)₂], [Fe(η⁵-C₅H₄CH(CH₃)NH(η⁵-C₅H₄CH(CH₃)))] and in the diaminecyclohexane salt [Fe(η⁵-C₅H₄COO)₂]²⁻·[(1*S*,2*S*)-(NH₃)₂C₆H₁₀]²⁺·2[H₂O], 609
- Crystal packing
 New chiral ferrocenyl-pyridinium salts for non-linear optics, 531
 Structural characterization and complexation behavior of ferrocene bearing dipeptide chain (-L-Ala-L-Pro-NHPy), 75
- Crystal structures
 A re-examination of the reaction between 1,1'-bis(diphenylphosphino)ferrocene (dppf) diselenide and [Ru₃(CO)₁₂]. Electrochemical behaviour of the isomeric *nido*-clusters [Ru₃(μ₃-Se)₂(dppf)(CO)₇] and [Ru₃(μ₃-Se)₂(μ-dppf)(CO)₇] and crystal structure of [Ru₃Se{μ-P(Ph)C₅H₄FeC₅H₄-PPh₂}(μ-OCPh)(CO)₆], 586
 Bis[bis(trimethylsilyl)cyclopentadienyl]selenane as a ligand: seleno-2,2',4,4'-tetrakis(trimethylsilyl)-[1]ferrocenophane and 1,2,3-triseleno-2,2',4,4'-tetrakis(trimethylsilyl)-[3]ferrocenophane, 745
 Ferrocene substituted nitronyl nitroxide and imino nitroxide radicals. Synthesis, X-ray structure and magnetic properties, 507

- 1,1'-Ferrocenyl-oligoprolines. A synthetic, structural and electrochemical study, 335
- Iron(II) versus osmium(II) oxidation in 1,1'-bis(diorganophosphino)ferrocene-osmium(II) complexes, 167
- Synthesis and characterisation of ferrocenyl-phosphonic and -arsonic acids, 216
- Synthesis and characterisation of rhodium(0) and rhodium(-1) complexes stabilised by 1,1'-bis(diphenylphosphino)ferrocene (dppf). Crystal structures of [Rh(dppf)₂] and [Na(THF)₅][Rh(dppf)₂]:THF, 710
- Synthesis and molecular structure of a novel ferrocene-containing macrocyclic acyl thiourea derivative, 204
- Synthesis, structures and electrochemical properties of Group 6 metal carbonyl complexes containing ferrocenylpyrazole ligands, 209
- Synthesis, structures and electrochemistry of bis(alkynylferrocene) complexes with fluorene spacers, 159
- Synthetic, electrochemical, and structural studies on heterobimetallic crown thioether complexes with Group 10 metals: the crystal structures of [Pt(9S3)(dppf)](PF₆)₂·CH₃NO₂ and [Pd(9S3)(dppf)](PF₆)₂·CH₃NO₂, 683
- Technetium and rhenium heterocomplexes containing the diphenylphosphinoferrocenyl fragment, 772
- The linear and nonlinear optical properties of organometallic chromophores derived from ferrocene, [Fe₂(η⁵-C₅H₅)₂(CO)₂(μ-CO)(μ-C-CH₃)]⁺[BF₄]⁻ and terthienyl spacers. Crystal structure of 2-[(E)-2-ferrocenylethenyl]-5-(2-thienyl)thiophene, 435
- The reaction of ferrocene with mercuric chloride revisited: 'Cp₂Fe*2HgCl₂' = [(C₅H₅)Fe(C₅H₄HgCl)]₂[Hg₂Cl₆], comparison of its molecular structure with that of chloromercuriferrocene, 796
- Cyclic voltammetry
[2 + 2 + 2] Cyclotrimerisation of bisaryl acetylene bearing ferrocenyl units with planar chirality: synthesis of enantiopure conjugated polyferrocene complexes, 84
- Cyclic voltammetry
Allylferrocenylselenide and the synthesis of the first seleno-substituted allenylidene complex: synthesis, spectroscopy, electrochemistry and the effect of electron transfer from the ferrocenylselenyl subunit, 240
- Electron acceptors of the fluorene series: Part 12. 9-(Metalocenylidene)nitrofluorene derivatives of Fc-NF, NF-Fc-NF, and NF-Rc-NF types, and the vinylogues Fc-π-NF: synthesis, characterisation, intramolecular charge transfer, redox properties and X-ray structures for three fluorene-ferrocene derivatives, 445
- 4-Ethynyl-benzonitrile-ferrocenes bridged by a Pd(PPh₃)₂ unit; the solid-state structure of (η⁵-C₅H₅)Fe(η⁵-C₅H₄C≡CC₆H₄C≡N-1,4), 198
- Improved electrochemistry of multi-ferrocenyl compounds: investigation of biferrocene, terferrocene, bis(fulvalene)diiron and diferrocenylethane in dichloromethane using [NBu₄][B(C₆F₅)₄] as supporting electrolyte, 823
- Multistep redox properties of 2,2'-bipyridylboronium substituted ferrocenes, 390
- Properties of some ferrocene macrocyclic dioxotetraamines: the roles of aromatic side-arms, 327
- Ruthenium(II) complexes containing 4-ferrocenylphenylisocyanide ligands. Crystal structure of *trans, trans, trans*-[RuCl₂-(POMe-*P*)₂(FcC₆H₄NC)] (POMe = PPh₂C₆H₄OCH₃; Fc = ferrocenyl), 145
- Synthesis and crystal structure of 1,1'-bis[4-(2,2':6',2''-terpyridin-4'-yl)phenyl]octamethylferrocene, 733
- Synthesis, structures and electrochemical properties of Group 6 metal carbonyl complexes containing ferrocenylpyrazole ligands, 209
- Synthesis, structures and electrochemistry of bis(alkynylferrocene) complexes with fluorene spacers, 159
- The effect of d-block metal complexation on the spectroscopic and redox properties of ferrocene derivatives containing pyridine ligands, 304
- Cycloaddition
Syntheses and some reactions of complexes containing carbon chains capped by ferrocenyl and W(CO)₃Cp groups, 484
- Cyclometallation
Cyclopalladated ferrocenylimines: highly active catalysts for Heck reactions, 793
- Novel cyclopalladated ferrocenyl Schiff base compounds with bridging and chelating diphosphines. Crystal and molecular structure of [{Pd}[(η⁵-C₅H₅)Fe(η⁵-C₅H₅)C(H)=N-2,4,6-Me₃C₆H₂)]{Ph₂P(CH₂)_nPPh₂-*P, P*}] [PF₆] (n = 1, 2), 577
- Studies on the cyclometallation of ferrocenylimines, 27
- Cyclopentadienes
Synthesis and crystal structure of 1,1'-bis[4-(2,2':6',2''-terpyridin-4'-yl)phenyl]octamethylferrocene, 733
- Cyclopentadienyl
Bis[bis(trimethylsilyl)cyclopentadienyl]selenane as a ligand: seleno-2,2',4,4'-tetrakis(trimethylsilyl)-[1]ferrocenophane and 1,2,3-triseleno-2,2',4,4'-tetrakis(trimethylsilyl)-[3]ferrocenophane, 745
- Cyclopenta[*l*]phenanthrene
Ferrocenes derived from cyclopenta[*l*]phenanthrene: dibenzindene-metal complexes that resist haptotropic shifts, 372
- Cyclophosphazene
Synthesis and characterization of ferrocenylalcohol derivatives of hexachlorocyclotriphosphazene. X-ray crystal structure of N₃P₃Cl₅OCH₂CH₂C₅H₄FeCp, 284
- Cyclotrimerisation
[2 + 2 + 2] Cyclotrimerisation of bisaryl acetylene bearing ferrocenyl units with planar chirality: synthesis of enantiopure conjugated polyferrocene complexes, 84
- Cyclotrimerisation
Complexation and electrochemical sensing of anions by amide-substituted ferrocenyl ligands, 356
- Cytotoxicity
Studies on organometallic selective estrogen receptor modulators. (SERMs) Dual activity in the hydroxy-ferrocifen series, 500
- Dendrimers
Functionalization via hydrosilylation of linear and cyclic siloxanes with appendent first generation dendrons containing electronically communicated ferrocenyl units, 642
- Water soluble ferrocenyl and polyferrocenyl compounds: synthesis and electrochemistry, 595
- 1,1'-Diacylferrocene
Preparation of ferrocenyl mono- and dienone derivatives through aldol condensation of 1,1'-diacylferrocene with aromatic aldehydes in dry conditions, 782
- Diacylhydrazine
Synthesis, spectroscopy and biological activity of novel acylhydrazines containing ferrocenyl moiety, 94
- Diels-Alder reaction
Synthesis and reactivity of achiral and of a novel planar chiral thioferrocenoylsilanes, 407
- 1,1'-Dienonylferrocenes
Preparation of ferrocenyl mono- and dienone derivatives through aldol condensation of 1,1'-diacylferrocene with aromatic aldehydes in dry conditions, 782
- Diffusion coefficient
Electrochemical determination of diffusion coefficient of π-conjugated polymers containing ferrocene unit, 809

Dipeptide

Structural characterization and complexation behavior of ferrocene bearing dipeptide chain (-L-Ala-L-Pro-NHPy), 75

Diphenylphosphinoferrrocene

Techetium and rhenium heterocomplexes containing the diphenylphosphinoferrrocenyl fragment, 772

Diphosphine

Palladium(II) phosphine thio-carboxylates. Structures of *cis*-Pd(κ S-SOCMe)₂(dppf) [dppf = 1,1'-bis(diphenylphosphino)ferrocene] and *trans*-Pd(κ S-SOCMe)₂(PPh₃)₂, 757

Diphosphine ligands

Synthetic, electrochemical, and structural studies on heterobimetallic crown thioether complexes with Group 10 metals: the crystal structures of [Pt(9S3)(dppf)](PF₆)₂·CH₃NO₂ and [Pd(9S3)(dppf)](PF₆)₂·CH₃NO₂, 683

Diyne

Synthesis of controlled π -extended conjugate nanostructures of 1,1'-ferrocene, 230

DNA-RNA hetero duplex

Ferrocenyl naphthalene diimide can bind to DNA-RNA hetero duplex: potential use in an electrochemical detection of mRNA expression, 476

dppf

Copper(I) complexes containing 1,1'-bis(diphenylphosphino) ferrocene (dppf) as a chelate ligand. X-ray crystal structures of [Cu₂(μ -Cl)₂(κ^2 -P,P-dppf)₂] and [Cu(CN^tBu)₂(κ^2 -P,P-dppf)](BF₄), 677

(E)- and (Z)-isomers

Synthesis of controlled π -extended conjugate nanostructures of 1,1'-ferrocene, 230

EHMO calculations

Ferrocenes derived from cyclopenta[*l*]phenanthrene: dibenzindene-metal complexes that resist haptotropic shifts, 372

Electrochemical behaviour

A re-examination of the reaction between 1,1'-bis(diphenylphosphino)ferrocene (dppf) diselenide and [Ru₃(CO)₁₂]. Electrochemical behaviour of the isomeric *nido*-clusters [Ru₃(μ_3 -Se)₂(dppf)(CO)₇] and [Ru₃(μ_3 -Se)₂(μ -dppf)(CO)₇] and crystal structure of [Ru₃Se{ μ -P(Ph)C₅H₄FeC₅H₄-PPh₂}(μ -OCPh)(CO)₆], 586

Electrochemical detection

Ferrocenyl naphthalene diimide can bind to DNA-RNA hetero duplex: potential use in an electrochemical detection of mRNA expression, 476

Electrochemical recognition

Complexation and electrochemical sensing of anions by amide-substituted ferrocenyl ligands, 356

Electrochemical sensor

Synthesis and dihydrogen phosphate binding properties of pyrrole containing *ansa*-ferrocenes, 343

Electrochemistry

An electrochemical study in acetonitrile of macrocyclic or open-chain ferrocene-containing oxa-aza or polyaza receptors in the presence of protons, metal cations and anions, 151

Characterization and electrochemical study of bis(ferrocenes) with a furan spacer and ferrocenophanes prepared from α -bromoacetyl substituted ferrocenes, 259

Diferrocenyltriphosphines 2. Reversible phosphine deligation in the chemistry of diferrocenyltriphosphine Ru(II) dichloride complexes with nitriles and pyridines: towards a pH-switchable catalyst?, 538

Facile synthesis of bidimensional ferrocenyl-based branched oligomers by palladium-catalyzed coupling reactions, 191

Functionalization via hydrosilylation of linear and cyclic siloxanes with appendent first generation dendrons containing electronically communicated ferrocenyl units, 642

Improved electrochemistry of multi-ferrocenyl compounds: investigation of biferrocene, terferrocene, bis(fulvalene)diiron and diferrocenylethane in dichloromethane using [NBu₄][B(C₆F₅)₄] as supporting electrolyte, 823

Internal ferrocenylalkynes—a comparative electrochemical and mass spectrometric study, 291

Iron(II) versus osmium(II) oxidation in 1,1'-bis(diorganophosphino)ferrocene-osmium(II) complexes, 167

Redox behavior of boronato-functionalized 1,1'-bis(diphenylphosphino)ferrocenes, 800

Solution phase peptide synthesis with ferrocenyl amino acid derivatives, 349

Star-shaped polyferrocenes based on thiophene and triphenylamine: synthesis, spectroscopy and electrochemistry, 139

Synthesis, characterisation, crystal structures and electrochemistry of triosmium nitrite carbonyl clusters containing ferrocenyl phosphines, 276

Synthetic, electrochemical, and structural studies on heterobimetallic crown thioether complexes with Group 10 metals: the crystal structures of [Pt(9S3)(dppf)](PF₆)₂·CH₃NO₂ and [Pd(9S3)(dppf)](PF₆)₂·CH₃NO₂, 683

Voltammetric metal cation sensors based on ferrocene derivatives with oxazoline and imine substituents, 134

Water soluble ferrocenyl and polyferrocenyl compounds: synthesis and electrochemistry, 595

Electron acceptors

Electron acceptors of the fluorene series: Part 12. 9-(Metallocenylidene)nitrofluorene derivatives of Fc-NF, NF-Fc-NF, and NF-Rc-NF types, and the vinyllogues Fc- π -NF: synthesis, characterisation, intramolecular charge transfer, redox properties and X-ray structures for three fluorene-ferrocene derivatives, 445

Electronic communication

Functionalization via hydrosilylation of linear and cyclic siloxanes with appendent first generation dendrons containing electronically communicated ferrocenyl units, 642

Electronic interaction

Synthesis of a new bis(ferrocenyl)ruthenacyclopentatriene compound with a significant inter-metal electronic communication, 80

Electronic spectra

Charge transfer spectra and photoreactivity of (*E*)-1-ferrocenyl-2-(1-methyl-4-pyridiniumyl)ethylene cation, 777

Electron transfer

Enzymatic chemistry of ferrocenes: micellar tuning of the glucose oxidase reactivity toward solubilized electrochemically generated *n*-alkylferricenium cations, 469

Electron-transfer chain catalysis

(η^5 -C₅H₅)-Ring alkylation reaction with the C₅H₅ anion: towards the construction of tri-Fe complex Fe{[μ , η^5 : η^4 -5-*exo*-(1'-C₅H₄)C₅H₅]Fe(CO)₂(PPh₃)₂}, 549

Electrospray mass spectrometry

Synthesis and characterisation of ferrocenyl-phosphonic and -arsonic acids, 216

Enantioselectivity

ortho-Silylation of 2,2'-bis(oxazolonyl)-1,1'-bis(diphenylphosphino)ferrocenes and remarkable effect of the silyl groups on the enantioselectivity in Pd-catalyzed asymmetric allylic alkylation, 99

Epoxy resins

Synthesis of new ferrocene containing diamines and their use in epoxy resins, 837

EPR spectroscopy

Iron(II) versus osmium(II) oxidation in 1,1'-bis(diorganophosphino)ferrocene-osmium(II) complexes, 167

ESR

Ferrocene as a ferromagnetic coupler. Synthesis and characterization of a ferrocene bridged polychlorotriphenylmethyl diradical, 251

Ethylnylferrocenes

Semimasked 1,1'-diethynylferrocenes: synthetic concepts, preparations, and reactions, 558

⁵⁷Fe Mössbauer spectroscopy

Control of the symmetry of binuclear ferrocene derivative by using chiral substituent and its mixed-valence state, 820

Ferricenium

The reaction of ferrocene with mercuric chloride revisited: 'Cp₂Fe*2HgCl₂' = [(C₅H₅)Fe(C₅H₄HgCl)]₂[Hg₂Cl₆], comparison of its molecular structure with that of chloromercuriferrocene, 796

Ferricenium ions

Enzymatic chemistry of ferrocenes: micellar tuning of the glucose oxidase reactivity toward solubilized electrochemically generated *n*-alkylferricenium cations, 469

Ferrocene

Allylferrocenylselenide and the synthesis of the first seleno-substituted allenylidene complex: synthesis, spectroscopy, electrochemistry and the effect of electron transfer from the ferrocenylselenyl subunit, 240

An electrochemical study in acetonitrile of macrocyclic or open-chain ferrocene-containing oxa-aza or polyaza receptors in the presence of protons, metal cations and anions, 151

A new phthalocyanine-ferrocene conjugated dyad, 751

Anion effect on the electrochemical characteristics of a gold electrode modified with a self-assembled monolayer of ferrocenylhexanethiol in aqueous and dichloromethane solutions, 841

Chiral ferrocenylthiazolidines, new ligands for palladium complexes, 116

Complexation and electrochemical sensing of anions by amide-substituted ferrocenyl ligands, 356

Conformational control of intramolecular arene stacking in ferrocene complexes bearing *tert*-butyl and pentafluorophenyl substituents, 107

Diferrocenyltriphosphines 2. Reversible phosphine deligation in the chemistry of diferrocenyltriphosphine Ru(II) dichloride complexes with nitriles and pyridines: towards a pH-switchable catalyst?, 538

Electron acceptors of the fluorene series: Part 12. 9-(Metalloenylidene)nitrofluorene derivatives of Fc-NF, NF-Fc-NF, and NF-Rc-NF types, and the vinylogues Fc- π -NF: synthesis, characterisation, intramolecular charge transfer, redox properties and X-ray structures for three fluorene-ferrocene derivatives, 445

Electrostatic field effects manifested in ferrocenyl metal complexes and the crystal structure of [Fe(η^5 -C₅H₅)(η^5 -C₅H₄CH=NNHC₃H₄N)]·HCl, 311

Enzymatic chemistry of ferrocenes: micellar tuning of the glucose oxidase reactivity toward solubilized electrochemically generated *n*-alkylferricenium cations, 469

4-Ethynyl-benzonitrile-ferrocenes bridged by a Pd(PPh₃)₂ unit; the solid-state structure of (η^5 -C₅H₅)Fe(η^5 -C₅H₄C≡CC₆H₄C≡N-1,4), 198

Ferrocene as a ferromagnetic coupler. Synthesis and characterization of a ferrocene bridged polychlorotriphenylmethyl diradical, 251

Ferrocene derivatives (I). Synthesis and X-ray structure determination of *N*-*o*-methylphenylferrocenesulfonamide, 738

Ferrocene polymers: current architectures, syntheses and utility, 47

Ferrocene substituted nitronyl nitroxide and imino nitroxide radicals. Synthesis, X-ray structure and magnetic properties, 507

1,1'-Ferrocenyl-oligoprolines. A synthetic, structural and electrochemical study, 335

Formation and structural characterization of amino-substituted [3]ferrocenophanes derived from intramolecular Mannich-type coupling reactions at the metallocene framework, 621

Functionalization via hydrosilylation of linear and cyclic siloxanes with appendent first generation dendrons containing electronically communicated ferrocenyl units, 642

Highly efficient reduction of ferrocenyl derivatives by borane, 364

Hindering the formation of ferrocenes: mono(cyclopentadienyl)halo iron complexes [Fe(C₅R₅)X] containing a sterically bulky cyclopentadienyl ligand, 786

Internal ferrocenylalkynes—a comparative electrochemical and mass spectrometric study, 291

Multistep redox properties of 2,2'-bipyridylboronium substituted ferrocenes, 390

New chiral ferrocenyl-pyridinium salts for non-linear optics, 531

New ferrocenyl amine derivatives: *N*-silyl, *N*-stannyl and *N*-boryl ferrocenyl amines, 727

ortho-Silylation of 2,2'-bis(oxazolonyl)-1,1'-bis(diphenylphosphino)ferrocenes and remarkable effect of the silyl groups on the enantioselectivity in Pd-catalyzed asymmetric allylic alkylation, 99

Palladium-catalyzed asymmetric allylic alkylations of cycloalkenyl acetates with planar chiral phosphino-ferrocene carboxylic acids, 845

Preparation of 1,5-diketone derivatives containing ferrocenyl by Michael reaction under solvent-free condition, 719

Properties of some ferrocene macrocyclic dioxetetraamines: the roles of aromatic side-arms, 327

Reactions of ferrocene-derived bis(hydroxymethyl) phosphine sulfides FcCH(R)P(S)(CH₂OH)₂ (R = H, CH₃) with cyclic thionylphosphazenes: crystal structures of FcCH₂P(S)-(CH₂O)₂PN(NPCl₂)[NS(O)Ph] and FcCH₂P(S)(CH₂O)₂-PN₂P[N(Me)CH₂]₂[NS(O)Ph] (Fc = ferrocenyl), 382

Ruthenium(II) complexes containing 4-ferrocenylphenylisocyanide ligands. Crystal structure of *trans*, *trans*-[RuCl₂(POMe-*P*)₂(FcC₆H₄NC)₂] (POMe = PPh₂C₆H₄-OCH₃; Fc = ferrocenyl), 145

Second harmonic generation in ferrocene based hydrogen bonded assemblies, 827

Solution phase peptide synthesis with ferrocenyl amino acid derivatives, 349

Star-shaped polyferrocenes based on thiophene and triphenylamine: synthesis, spectroscopy and electrochemistry, 139

Structural characterization and complexation behavior of ferrocene bearing dipeptide chain (-L-Ala-L-Pro-NHPy), 75

Studies on organometallic selective estrogen receptor modulators. (SERMs) Dual activity in the hydroxy-ferrocifen series, 500

Study on the synthesis of nonracemic C₂-symmetric 1,1'-binaphthyl-2,2'-diyl bridged ferrocene. Stereochemical result of the cross-coupling reactions controlled by Pd(II) or Pd(IV) complex intermediacy, 318

Synthesis and characterisation of ferrocenyl-phosphonic and -arsonic acids, 216

Synthesis and characterization of ferrocene-labeled oligodeoxynucleotides, 398

Synthesis and dihydrogen phosphate binding properties of pyrrole containing *ansa*-ferrocenes, 343

Synthesis and reactivity of achiral and of a novel planar chiral thioferrocenoylsilanes, 407

Synthesis and structure of an alkylgallium bridged [1,1]ferrocenophane, 300

- Synthesis, characterisation, crystal structures and electrochemistry of trisium nitrite carbonyl clusters containing ferrocenyl phosphines, 276
- Synthesis of a new bis(ferrocenyl)ruthenacyclopentatriene compound with a significant inter-metal electronic communication, 80
- Synthesis of 1,1'-bis(heterocyclyl)biferrocene compounds, 723
- Synthesis of new ferrocene containing diamines and their use in epoxy resins, 837
- Synthesis, spectroscopy and biological activity of novel acylhydrazines containing ferrocenyl moiety, 94
- Synthesis, structures and electrochemistry of bis(alkynylferrocene) complexes with fluorene spacers, 159
- The effect of d-block metal complexation on the spectroscopic and redox properties of ferrocene derivatives containing pyridine ligands, 304
- The linear and nonlinear optical properties of organometallic chromophores derived from ferrocene, $[\text{Fe}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_2(\mu\text{-CO})(\mu\text{-C-CH}_3)]^+ [\text{BF}_4]^-$ and terthienyl spacers. Crystal structure of 2-[(*E*)-2-ferrocenylethenyl]-5-(2-thienyl)thiophene, 435
- The reaction of ferrocene with mercuric chloride revisited: ' $\text{Cp}_2\text{Fe}^*\text{2HgCl}_2$ ' = $[(\text{C}_5\text{H}_5)\text{Fe}(\text{C}_5\text{H}_4\text{HgCl})_2][\text{Hg}_2\text{Cl}_6]$, comparison of its molecular structure with that of chloromercuriferrocene, 796
- Voltammetric metal cation sensors based on ferrocene derivatives with oxazoline and imine substituents, 134
- Ferrocenecarboxaldehydes**
Knoevenagel condensation of ferrocenyl substituted carboxaldehydes with methylene active compounds on inorganic supports and the NLO properties of the resulting push-pull alkenes, 426
- Ferrocenecarboxylic anhydride**
Synthesis, spectroscopy and biological activity of novel acylhydrazines containing ferrocenyl moiety, 94
- Ferrocene compounds**
Charge transfer spectra and photoreactivity of (*E*)-1-ferrocenyl-2-(1-methyl-4-pyridiniumyl)ethylene cation, 777
Iron(II) versus osmium(II) oxidation in 1,1'-bis(diorganophosphino)ferrocene-osmium(II) complexes, 167
- Ferrocene-containing polymers**
Electrochemical determination of diffusion coefficient of π -conjugated polymers containing ferrocene unit, 809
- Ferrocene derivatives**
Synthesis and molecular structure of a novel ferrocene-containing macrocyclic acyl thiourea derivative, 204
- Ferrocenediyl-bridged tri-Fe complex**
 $(\eta^5\text{-C}_5\text{H}_5)$ -Ring alkylation reaction with the C_5H_5 anion: towards the construction of tri-Fe complex $\text{Fe}\{[\mu, \eta^5: \eta^4\text{-5-}exo\text{-}(1'\text{-C}_5\text{H}_4)\text{C}_5\text{H}_5]\text{Fe}(\text{CO})_2(\text{PPh}_3)\}_2$, 549
- Ferrocene half-sandwich**
Bonding in 'closed,' open, and half-open ferrocenes: new insight from structural and Mössbauer spectroscopic studies, 172
- Ferrocene ligands**
Synthesis, reactivity, and crystal structures of ferrocene-substituted amidinate derivatives, 521
- Ferrocene metal cluster**
Syntheses and some reactions of complexes containing carbon chains capped by ferrocenyl and $\text{W}(\text{CO})_3\text{Cp}$ groups, 484
- Ferrocenes**
A rapid approach to ferrocenophanes via ring-closing metathesis, 669
Bis(ferrocenyl)mercury as a source of ferrocenyl moiety in Pd-catalyzed reactions of carbon-carbon bond formation, 653
Facile synthesis of bidimensional ferrocenyl-based branched oligomers by palladium-catalyzed coupling reactions, 191
Synthesis of controlled π -extended conjugate nanostructures of 1,1'-ferrocene, 230
Synthesis of ferrocenylketyl radicals by chromium(II) complexes, 813
- Ferrocenesulfonamide**
Ferrocene derivatives (I). Synthesis and X-ray structure determination of *N*-*o*-methylphenylferrocenesulfonamide, 738
- Ferrocenesulfonyl chloride**
Ferrocene derivatives (I). Synthesis and X-ray structure determination of *N*-*o*-methylphenylferrocenesulfonamide, 738
- Ferrocenium ion rearrangement**
Redox behavior of boronato-functionalized 1,1'-bis(diphenylphosphino)ferrocenes, 800
- Ferrocenophane**
Synthesis and structure of an alkylgallium bridged [1,1]ferrocenophane, 300
- [1]Ferrocenophane**
Bis[bis(trimethylsilyl)cyclopentadienyl]selena-2,2',4,4'-tetrakis(trimethylsilyl)-[1]ferrocenophane and 1,2,3-triselena-2,2',4,4'-tetrakis(trimethylsilyl)-[3]ferrocenophane, 745
- [3]Ferrocenophane**
Bis[bis(trimethylsilyl)cyclopentadienyl]selena-2,2',4,4'-tetrakis(trimethylsilyl)-[1]ferrocenophane and 1,2,3-triselena-2,2',4,4'-tetrakis(trimethylsilyl)-[3]ferrocenophane, 745
- Ferrocenophanes**
Characterization and electrochemical study of bis(ferrocenes) with a furan spacer and ferrocenophanes prepared from α -bromoacetyl substituted ferrocenes, 259
- [1]Ferrocenophanes**
Synthesis and characterization of new germylferrocenyl, germylferrocenophane and polymeric germyleneferrocenylene systems, 664
- Ferrocenophanes**
A rapid approach to ferrocenophanes via ring-closing metathesis, 669
Formation and structural characterization of amino-substituted [3]ferrocenophanes derived from intramolecular Mannich-type coupling reactions at the metallocene framework, 621
- Ferrocenoyl equivalent**
An investigation of the chemistry of ferrocenoyl derivatives. The synthesis and reactions of ferrocenoyl imidazolide and its derivatives, 266
- Ferrocenoyl imidazolide**
An investigation of the chemistry of ferrocenoyl derivatives. The synthesis and reactions of ferrocenoyl imidazolide and its derivatives, 266
- Ferrocenyl**
(Ferrocenylmethyl)trialkyl ammoniums as template cations in optically active two-dimensional oxalate bridged [Cr-Mn] and [Cr-Ni] molecule-based magnets: synthesis and magnetic properties, 182
- Ferrocenylalcohols**
Synthesis and characterization of ferrocenylalcohol derivatives of hexachlorocyclotriphosphazene. X-ray crystal structure of $\text{N}_3\text{P}_3\text{Cl}_5\text{OCH}_2\text{CH}_2\text{C}_5\text{H}_4\text{FeCp}$, 284
- Ferrocenyl amines**
Metal-directed assembly of polyferrocenyl transition metal dithiocarbamate macrocyclic molecular boxes, 89
- Ferrocenyl-carbenium**
A comparison of ferrocenyl carbenium/non-carbenium structures in isomeric osmium cluster complexes obtained from the reaction of bis-ferrocenylbutadiyne with $\text{Os}_3(\text{CO})_{10}(\mu\text{-py})(\mu\text{-H})$, 514

- Ferrocenyl complexes
[2 + 2 + 2] Cyclotrimerisation of bisaryl acetylene bearing ferrocenyl units with planar chirality: synthesis of enantiopure conjugated polyferrocene complexes, 84
- Ferrocenyl compounds
Water soluble ferrocenyl and polyferrocenyl compounds: synthesis and electrochemistry, 595
- Ferrocenyl-derived ligands
Novel ferrocene modified *P,N*-ligands for enantioselective palladium-catalyzed allylic substitution reactions, 762
- 1,1'-Ferrocenyl diketones
Asymmetric synthesis of planar chiral 2-mono- and 2,2'-disubstituted 1,1'-bisbenzoylferrocenes, 698
- Ferrocenylene phosphine
Palladium(II) phosphine thio-carboxylates. Structures of *cis*-Pd(κ S-SOCMe)₂(dppf) [dppf = 1,1'-bis(diphenylphosphino)ferrocene] and *trans*-Pd(κ S-SOCMe)₂(PPh₃)₂, 757
- Ferrocenylimine
Cyclopalladated ferrocenylimines: highly active catalysts for Heck reactions, 793
Studies on the cyclometallation of ferrocenylimines, 27
- Ferrocenyl naphthalene diimide
Ferrocenyl naphthalene diimide can bind to DNA-RNA heteroduplex: potential use in an electrochemical detection of mRNA expression, 476
- Ferrocenyloxazoline
Octahedral ruthenium(II) complexes containing the chiral ligand (4*S*)-2-[(*S_p*)-2-(diphenylphosphino)ferrocenyl]-4-(isopropyl)oxazoline (FcPN). X-Ray crystal structures of *fac*-[RuCl₂(PMe₃)₂(FcPN)] and *fac*-[RuCl₂(dppm)(FcPN)] (dppm = bis(diphenylphosphino)methane), 463
- Ferrocenylphosphine
Octahedral ruthenium(II) complexes containing the chiral ligand (4*S*)-2-[(*S_p*)-2-(diphenylphosphino)ferrocenyl]-4-(isopropyl)oxazoline (FcPN). X-Ray crystal structures of *fac*-[RuCl₂(PMe₃)₂(FcPN)] and *fac*-[RuCl₂(dppm)(FcPN)] (dppm = bis(diphenylphosphino)methane), 463
- Ferrocenyl phosphine
Synthesis and characterisation of rhodium(0) and rhodium(-1) complexes stabilised by 1,1'-bis(diphenylphosphino)ferrocene (dppf). Crystal structures of [Rh(dppf)₂] and [Na(THF)₅][Rh(dppf)₂]:THF, 710
- Ferrocenylpyrazole
Synthesis, structures and electrochemical properties of Group 6 metal carbonyl complexes containing ferrocenylpyrazole ligands, 209
- Ferrocenyl substituted olefins
Ferrocenyl substituted chlorostilbenes and butadienes, 616
- 4-Ferrocenyl-2-thiazolamine
Ferrocene derivatives (II). Synthesis and reactions of 4-ferrocenyl-2-thiazolamine, 742
- Ferromagnetism
Ferrocene as a ferromagnetic coupler. Synthesis and characterization of a ferrocene bridged polychlorotriphenylmethyl diradical, 251
- Fluorene
Synthesis, structures and electrochemistry of bis(alkynylferrocene) complexes with fluorene spacers, 159
- Gallium
Synthesis and structure of an alkylgallium bridged [1,1]ferrocenophane, 300
- Germyleneferrocenylene systems
Synthesis and characterization of new germlyferrocenyl, germlyferrocenophane and polymeric germyleneferrocenylene systems, 664
- Germlyferrocenophane
Synthesis and characterization of new germlyferrocenyl, germlyferrocenophane and polymeric germyleneferrocenylene systems, 664
- Glucose oxidase
Enzymatic chemistry of ferrocenes: micellar tuning of the glucose oxidase reactivity toward solubilized electrochemically generated *n*-alkylferrocenium cations, 469
- Group IX and X complexes
1,1',2,2'-Tetrakis(diphenylphosphino)-4,4'-di-*tert*-butylferrocene, a new *cisoid* arrangement of phosphino groups, 126
- Half-sandwich complexes
Hindering the formation of ferrocenes: mono(cyclopentadienyl)halo iron complexes [Fe(C₅R₅)X] containing a sterically bulky cyclopentadienyl ligand, 786
- Haptotropic shifts
Ferrocenes derived from cyclopenta[*l*]phenanthrene: dibenzindene-metal complexes that resist haptotropic shifts, 372
- Heck reaction
Cyclopalladated ferrocenylimines: highly active catalysts for Heck reactions, 793
- Heterocyclylbiferrocene
Synthesis of 1',1''-bis(heterocyclyl)biferrocene compounds, 723
- Homocoupling
Synthesis of 1',1''-bis(heterocyclyl)biferrocene compounds, 723
- Horner-Wadsworth-Emmons reaction
Ferrocenyl substituted chlorostilbenes and butadienes, 616
- Hydrazones
Asymmetric synthesis of planar chiral 2-mono- and 2,2'-disubstituted 1,1'-bisbenzoylferrocenes, 698
- Hydride abstraction
(η^5 -C₅H₅)-Ring alkylation reaction with the C₅H₅ anion: towards the construction of tri-Fe complex Fe{[μ , η^5 : η^4 -5-*exo*-(1'-C₅H₄)C₅H₅]Fe(CO)₂(PPh₃)₂}, 549
- Hydrogen bond
Structural characterization and complexation behavior of ferrocene bearing dipeptide chain (-L-Ala-L-Pro-NHPy), 75
- Hydrogen bonding
Second harmonic generation in ferrocene based hydrogen bonded assemblies, 827
Synthesis and molecular structure of a novel ferrocene-containing macrocyclic acyl thiourea derivative, 204
- Hydrogen bonds
Organometallic building blocks for crystal engineering. Synthesis, structure and hydrogen bonding interactions in [Fe(η^5 -C₅H₄-CH₂(CH₃)OH)₂], [Fe(η^5 -C₅H₃(CH₃)COOH)₂], [Fe(η^5 -C₅H₄CH(CH₃)NH(η^5 -C₅H₄CH(CH₃)))] and in the diaminecyclohexane salt [Fe(η^5 -C₅H₄COO)₂]²⁻[(1*S*,2*S*)-(NH₃)₂C₆H₁₀]²⁺·2[H₂O], 609
- Hydrosilylation
Functionalization via hydrosilylation of linear and cyclic siloxanes with appended first generation dendrons containing electronically communicated ferrocenyl units, 642
- Hydroxymethyl phosphine sulfide
Reactions of ferrocene-derived bis(hydroxymethyl) phosphine sulfides FeCH(R)P(S)(CH₂OH)₂ (R = H, CH₃) with cyclic thionylphosphazenes: crystal structures of FeCH₂P(S)-(CH₂O)₂PN(NPCl₂)[NS(O)Ph] and FeCH₂P(S)(CH₂O)₂-PN₂P[N(Me)CH₂]₂[NS(O)Ph] (Fc = ferrocenyl), 382
- β-Hydroxy sulphides
Synthesis and reactivity of achiral and of a novel planar chiral thioferrocenoylsilanes, 407
- Hyper-Rayleigh scattering
Knoevenagel condensation of ferrocenyl substituted carboxaldehydes with methylene active compounds on inorganic supports and the NLO properties of the resulting push-pull alkenes, 426

- Imine**
 Novel cyclopalladated ferrocenyl Schiff base compounds with bridging and chelating diphosphines. Crystal and molecular structure of $\{[\text{Pd}[(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\eta^5\text{-C}_5\text{H}_5)\text{C}(\text{H})=\text{N}-2,4,6\text{-Me}_3\text{C}_6\text{H}_2]\{\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2\text{-}P,P\}][\text{PF}_6]\}$ ($n = 1, 2$), 577
- Insect growth regulators**
 Synthesis, spectroscopy and biological activity of novel acylhydrazines containing ferrocenyl moiety, 94
- Intramolecular charge transfer**
 Electron acceptors of the fluorene series: Part 12. 9-(Metallo-cene-ylidene)nitrofluorene derivatives of Fc–NF, NF–Fc–NF, and NF–Rc–NF types, and the vinylogues Fc– π –NF: synthesis, characterisation, intramolecular charge transfer, redox properties and X-ray structures for three fluorene–ferrocene derivatives, 445
- Intramolecular coordination**
 Studies on the cyclometallation of ferrocenylimines, 27
- Iron**
 Highly efficient reduction of ferrocenyl derivatives by borane, 364
 New chiral ferrocenyl-pyridinium salts for non-linear optics, 531
 Synthesis and crystal structure of 1,1'-bis[4-(2,2':6',2''-terpyridin-4'-yl)phenyl]octamethylferrocene, 733
 Synthesis, reactivity, and crystal structures of ferrocene-substituted amidinate derivatives, 521
- Isocyanide**
 Ruthenium(II) complexes containing 4-ferrocenylphenylisocyanide ligands. Crystal structure of *trans*, *trans*, *trans*- $[\text{RuCl}_2(\text{POMe}-P)_2(\text{FcC}_6\text{H}_4\text{NC})_2]$ (POMe = $\text{PPh}_2\text{C}_6\text{H}_4\text{-OCH}_3$; Fc = ferrocenyl), 145
- Isocyanide complexes**
 Copper(I) complexes containing 1,1'-bis(diphenylphosphino) ferrocene (dppf) as a chelate ligand. X-ray crystal structures of $[\text{Cu}_2(\mu\text{-Cl})_2(\kappa^2\text{-}P,P\text{-dppf})_2]$ and $[\text{Cu}(\text{CN}'\text{Bu})_2(\kappa^2\text{-}P,P\text{-dppf})][\text{BF}_4]$, 677
- Isomeric nido-clusters**
 A re-examination of the reaction between 1,1'-bis(diphenylphosphino)ferrocene (dppf) diselenide and $[\text{Ru}_3(\text{CO})_{12}]$. Electrochemical behaviour of the isomeric *nido*-clusters $[\text{Ru}_3(\mu_3\text{-Se})_2(\text{dppf})(\text{CO})_7]$ and $[\text{Ru}_3(\mu_3\text{-Se})_2(\mu\text{-dppf})(\text{CO})_7]$ and crystal structure of $[\text{Ru}_3\text{Se}\{\mu\text{-P}(\text{Ph})\text{C}_5\text{H}_4\text{FeC}_5\text{H}_4\text{-PPh}_2\}(\mu\text{-OCPh})(\text{CO})_6]$, 586
- Ketone**
 Preparation of 1,5-diketone derivatives containing ferrocenyl by Michael reaction under solvent-free condition, 719
- Ketones**
 Synthesis of ferrocenylketyl radicals by chromium(II) complexes, 813
- Kinetics**
 Enzymatic chemistry of ferrocenes: micellar tuning of the glucose oxidase reactivity toward solubilized electrochemically generated *n*-alkylferricenium cations, 469
- Knoevenagel condensation**
 Knoevenagel condensation of ferrocenyl substituted carboxaldehydes with methylene active compounds on inorganic supports and the NLO properties of the resulting push–pull alkenes, 426
- Lethal larval molt**
 Synthesis, spectroscopy and biological activity of novel acylhydrazines containing ferrocenyl moiety, 94
- Macrocyclic molecular boxes**
 Metal-directed assembly of polyferrocenyl transition metal dithiocarbamate macrocyclic molecular boxes, 89
- Macrocyclic polyamine**
 Properties of some ferrocene macrocyclic dioxetraamines: the roles of aromatic side-arms, 327
- Magnetochemistry**
 Ferrocene substituted nitronyl nitroxide and imino nitroxide radicals. Synthesis, X-ray structure and magnetic properties, 507
- Mass spectrometry**
 Internal ferrocenylalkynes—a comparative electrochemical and mass spectrometric study, 291
 Redox behavior of boronato-functionalized 1,1'-bis(diphenylphosphino)ferrocenes, 800
- MCF7**
 Studies on organometallic selective estrogen receptor modulators. (SERMs) Dual activity in the hydroxy-ferrocifen series, 500
- Mechanism**
 Study on the synthesis of nonracemic C_2 -symmetric 1,1'-binaphthyl-2,2'-diyl bridged ferrocene. Stereochemical result of the cross-coupling reactions controlled by Pd(II) or Pd(IV) complex intermediacy, 318
- Mercury**
 The reaction of ferrocene with mercuric chloride revisited: ' $\text{Cp}_2\text{Fe}^*\text{2HgCl}_2$ ' = $[(\text{C}_5\text{H}_5)\text{Fe}(\text{C}_5\text{H}_4\text{HgCl})_2][\text{Hg}_2\text{Cl}_6]$, comparison of its molecular structure with that of chloromercuriferrocene, 796
- Merocyanine**
 The linear and nonlinear optical properties of organometallic chromophores derived from ferrocene, $[\text{Fe}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_2(\mu\text{-CO})(\mu\text{-C-CH}_3)]^+[\text{BF}_4]^-$ and terthienyl spacers. Crystal structure of 2-[(*E*)-2-ferrocenylethenyl]-5-(2-thienyl)thiophene, 435
- Metal carbonyl complexes**
 Synthesis, structures and electrochemical properties of Group 6 metal carbonyl complexes containing ferrocenylpyrazole ligands, 209
- Metal-containing dendrimers**
 Facile synthesis of bidimensional ferrocenyl-based branched oligomers by palladium-catalyzed coupling reactions, 191
- Metal-directed assembly**
 Metal-directed assembly of polyferrocenyl transition metal dithiocarbamate macrocyclic molecular boxes, 89
- Metallacycle**
 Synthesis of a new bis(ferrocenyl)ruthenacyclopentatriene compound with a significant inter-metal electronic communication, 80
- Metalocene**
 Ferrocene polymers: current architectures, syntheses and utility, 47
- Metalloenes**
 Bonding in 'closed,' open, and half-open ferrocenes: new insight from structural and Mössbauer spectroscopic studies, 172
- Metal–metal interaction**
 4-Ethynyl-benzonitrile-ferrocenes bridged by a $\text{Pd}(\text{PPh}_3)_2$ unit; the solid-state structure of $(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\eta^5\text{-C}_5\text{H}_4\text{C}\equiv\text{CC}_6\text{H}_4\text{C}\equiv\text{N-1,4})$, 198
- Micelles**
 Enzymatic chemistry of ferrocenes: micellar tuning of the glucose oxidase reactivity toward solubilized electrochemically generated *n*-alkylferricenium cations, 469
- Michael reaction**
 Preparation of 1,5-diketone derivatives containing ferrocenyl by Michael reaction under solvent-free condition, 719
- Mixed ligand complexes**
 Technetium and rhenium heterocomplexes containing the diphenylphosphinoferrocenyl fragment, 772

Mixed-valence state

Control of the symmetry of binuclear ferrocene derivative by using chiral substituent and its mixed-valence state, 820

Modified electrodes

Functionalization via hydrosilylation of linear and cyclic siloxanes with appendent first generation dendrons containing electronically communicated ferrocenyl units, 642

Molecular recognition

Voltammetric metal cation sensors based on ferrocene derivatives with oxazoline and imine substituents, 134

Molecule-based magnet

(Ferrocenylmethyl)trialkyl ammoniums as template cations in optically active two-dimensional oxalate bridged [Cr–Mn] and [Cr–Ni] molecule-based magnets: synthesis and magnetic properties, 182

Mössbauer

Electrostatic field effects manifested in ferrocenyl metal complexes and the crystal structure of $[\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\eta^5\text{-C}_5\text{H}_4\text{CH}=\text{NNHC}_5\text{H}_4\text{N})]\cdot\text{HCl}$, 311

Mössbauer spectroscopy

Bonding in ‘closed,’ open, and half-open ferrocenes: new insight from structural and Mössbauer spectroscopic studies, 172

Nitrido complexes

Technetium and rhenium heterocomplexes containing the diphenylphosphinoferrrocenyl fragment, 772

Nitrite

Synthesis, characterisation, crystal structures and electrochemistry of triosmium nitrite carbonyl clusters containing ferrocenyl phosphines, 276

NLO properties

Knoevenagel condensation of ferrocenyl substituted carboxaldehydes with methylene active compounds on inorganic supports and the NLO properties of the resulting push–pull alkenes, 426

NMR

1,1'-Ferrocenyl–oligoprolines. A synthetic, structural and electrochemical study, 335

New ferrocenyl amine derivatives: *N*-silyl, *N*-stannyl and *N*-boryl ferrocenyl amines, 727

Synthesis, X-ray, spectroscopic and a preliminary Suzuki coupling screening studies of a complete series of dppfMX_2 ($M = \text{Pt}$, Pd ; $X = \text{Cl}$, Br , I), 691

NMR spectroscopy

Palladium(II) complexes containing symmetrical diphosphine ligands based on ferrocene. Sterically enforced palladium–iron bonds, 805

Nondestructive imaging

Synthesis of new ferrocene containing diamines and their use in epoxy resins, 837

Nonlinear optics

Second harmonic generation in ferrocene based hydrogen bonded assemblies, 827

The linear and nonlinear optical properties of organometallic chromophores derived from ferrocene, $[\text{Fe}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_2(\mu\text{-CO})(\mu\text{-C-CH}_3)]^+[\text{BF}_4]^-$ and terthienyl spacers. Crystal structure of 2-[(*E*)-2-ferrocenylethenyl]-5-(2-thienyl)thiophene, 435

Non-linear optics

New chiral ferrocenyl-pyridinium salts for non-linear optics, 531

 $\eta^3\text{:}\eta^3$ -Octadienediyl complexes

Reactivity of the bis(allyl)-ruthenium(IV) dimer $[\{\text{Ru}(\eta^3\text{:}\eta^3\text{-C}_{10}\text{H}_{16})(\mu\text{-Cl})\text{Cl}\}_2]$ ($\text{C}_{10}\text{H}_{16} = 2,7$ -dimethylocta-2,6-diene-1,8-diyl) towards dppf ($\text{dppf} = [\text{Fe}(\eta^5\text{-C}_5\text{H}_4\text{PPh}_2)_2]$) and dppfO ($\text{dppfO} = [\text{Fe}(\eta^5\text{-C}_5\text{H}_4\text{PPh}_2)(\eta^5\text{-C}_5\text{H}_4\text{P}(=\text{O})\text{Ph}_2)]$), 767

Oligodeoxynucleotides

Synthesis and characterization of ferrocene-labeled oligodeoxynucleotides, 398

Oligonuclear complexes

Multistep redox properties of 2,2'-bipyridylboronium substituted ferrocenes, 390

Oligopeptide

1,1'-Ferrocenyl–oligoprolines. A synthetic, structural and electrochemical study, 335

Oligopyridine ligands

Synthesis and crystal structure of 1,1'-bis[4-(2,2':6',2''-terpyridin-4'-yl)phenyl]octamethylferrocene, 733

On-column

Synthesis and characterization of ferrocene-labeled oligodeoxynucleotides, 398

Organic radical

Ferrocene as a ferromagnetic coupler. Synthesis and characterization of a ferrocene bridged polychlorotriphenylmethyl diradical, 251

Ferrocene substituted nitronyl nitroxide and imino nitroxide radicals. Synthesis, X-ray structure and magnetic properties, 507

Organic solvent

Anion effect on the electrochemical characteristics of a gold electrode modified with a self-assembled monolayer of ferrocenylhexanethiol in aqueous and dichloromethane solutions, 841

Organometallic complexes

Organometallic building blocks for crystal engineering. Synthesis, structure and hydrogen bonding interactions in $[\text{Fe}(\eta^5\text{-C}_5\text{H}_4\text{-CH}_2(\text{CH}_3)\text{OH})_2]$, $[\text{Fe}(\eta^5\text{-C}_5\text{H}_3(\text{CH}_3)\text{COOH})_2]$, $[\text{Fe}(\eta^5\text{-C}_5\text{H}_4\text{CH}(\text{CH}_3)\text{NH}(\eta^5\text{-C}_5\text{H}_4\text{CH}(\text{CH}_3)))]$ and in the diaminecyclohexane salt $[\text{Fe}(\eta^5\text{-C}_5\text{H}_4\text{COO})_2]^{2-}[(1S,2S)\text{-}(\text{NH}_3)_2\text{C}_6\text{H}_{10}]^{2+}\cdot 2[\text{H}_2\text{O}]$, 609

Organometallic hormone

Studies on organometallic selective estrogen receptor modulators. (SERMs) Dual activity in the hydroxy-ferrocifen series, 500

ortho-Metalation

Asymmetric synthesis of planar chiral 2-mono- and 2,2'-disubstituted 1,1'-bisbenzoylferrocenes, 698

Osmium

A comparison of ferrocenyl carbenium/non-carbenium structures in isomeric osmium cluster complexes obtained from the reaction of bis-ferrocenylbutadiyne with $\text{Os}_3(\text{CO})_{10}(\mu\text{-py})(\mu\text{-H})$, 514

Synthesis, characterisation, crystal structures and electrochemistry of triosmium nitrite carbonyl clusters containing ferrocenyl phosphines, 276

Osmium complexes

Iron(II) versus osmium(II) oxidation in 1,1'-bis(diorganophosphino)ferrocene–osmium(II) complexes, 167

Oxalate bridge

(Ferrocenylmethyl)trialkyl ammoniums as template cations in optically active two-dimensional oxalate bridged [Cr–Mn] and [Cr–Ni] molecule-based magnets: synthesis and magnetic properties, 182

Oxazoline

Voltammetric metal cation sensors based on ferrocene derivatives with oxazoline and imine substituents, 134

Palladium

Characterization and electrochemical study of bis(ferrocenes) with a furan spacer and ferrocenophanes prepared from α -bromoacetyl substituted ferrocenes, 259

- Chiral ferrocenylthiazolidines, new ligands for palladium complexes, 116
- Novel cyclopalladated ferrocenyl Schiff base compounds with bridging and chelating diphosphines. Crystal and molecular structure of $[\{\text{Pd}[(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\eta^5\text{-C}_5\text{H}_5)\text{C}(\text{H})=\text{N}-2,4,6\text{-Me}_3\text{C}_6\text{H}_2]\}\{\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2\text{-}P,P\}][\text{PF}_6^-]$ ($n = 1, 2$), 577
- ortho*-Silylation of 2,2'-bis(oxazolanyl)-1,1'-bis(diphenylphosphino)ferrocenes and remarkable effect of the silyl groups on the enantioselectivity in Pd-catalyzed asymmetric allylic alkylation, 99
- Palladium(II) phosphine thiocarboxylates. Structures of *cis*-Pd($\kappa\text{S-SOCMe}$)₂(dppf) [dppf = 1,1'-bis(diphenylphosphino)ferrocene] and *trans*-Pd($\kappa\text{S-SOCMe}$)₂(PPh₃)₂, 757
- Palladium complex
- Structural characterization and complexation behavior of ferrocene bearing dipeptide chain (-L-Ala-L-Pro-NHPy), 75
- Palladium complexes
- Study on the synthesis of nonracemic C₂-symmetric 1,1'-binaphthyl-2,2'-diyl bridged ferrocene. Stereochemical result of the cross-coupling reactions controlled by Pd(II) or Pd(IV) complex intermediacy, 318
- Palladium–iron bonds
- Palladium(II) complexes containing symmetrical diphosphine ligands based on ferrocene. Sterically enforced palladium–iron bonds, 805
- Pd(0)-catalysed cross-coupling
- A new phthalocyanine–ferrocene conjugated dyad, 751
- Pentafluorophenyl
- Conformational control of intramolecular arene stacking in ferrocene complexes bearing *tert*-butyl and pentafluorophenyl substituents, 107
- Peptide synthesis
- Solution phase peptide synthesis with ferrocenyl amino acid derivatives, 349
- Perfluorophenyl borate
- Improved electrochemistry of multi-ferrocenyl compounds: investigation of biferrocene, terferrocene, bis(fulvalene)diiron and diferrocenylethane in dichloromethane using [NBu₄][B(C₆F₅)₄] as supporting electrolyte, 823
- Phosphate binding
- Synthesis and dihydrogen phosphate binding properties of pyrrole containing *ansa*-ferrocenes, 343
- Phosphine ligand
- Diferrocenyltriphosphines 2. Reversible phosphine deligation in the chemistry of diferrocenyltriphosphine Ru(II) dichloride complexes with nitriles and pyridines: towards a pH-switchable catalyst?, 538
- Phosphine ligands
- Palladium(II) complexes containing symmetrical diphosphine ligands based on ferrocene. Sterically enforced palladium–iron bonds, 805
- Phosphinoferrocene
- 1,1',2,2'-Tetrakis(diphenylphosphino)-4,4'-di-*tert*-butylferrocene, a new *cisoid* arrangement of phosphino groups, 126
- Phosphonic acid
- Synthesis and characterisation of ferrocenyl-phosphonic and -arsonic acids, 216
- Photochemistry
- Charge transfer spectra and photoreactivity of (*E*)-1-ferrocenyl-2-(1-methyl-4-pyridiniumyl)ethylene cation, 777
- Phthalocyanines
- A new phthalocyanine–ferrocene conjugated dyad, 751
- Planar chirality
- Asymmetric synthesis of planar chiral 2-mono- and 2,2'-disubstituted 1,1'-bisbenzoylferrocenes, 698
- [2 + 2 + 2] Cyclootrimerisation of bisaryl acetylene bearing ferrocenyl units with planar chirality: synthesis of enantiopure conjugated polyferrocene complexes, 84
- Novel ferrocene modified *P,N*-ligands for enantioselective palladium-catalyzed allylic substitution reactions, 762
- Palladium-catalyzed asymmetric allylic alkylations of cycloalkenyl acetates with planar chiral phosphino-ferrocene carboxylic acids, 845
- Synthesis and reactivity of achiral and of a novel planar chiral thioferrocenoylsilanes, 407
- Platinum
- Synthesis and characterisation of novel multidentate ferrocene ligands and their Re(I) and Pt(II) complexes, 418
- Polycarboxylic acids
- Organometallic building blocks for crystal engineering. Synthesis, structure and hydrogen bonding interactions in [Fe($\eta^5\text{-C}_5\text{H}_4\text{-CH}_2(\text{CH}_3)\text{OH}$)₂], [Fe($\eta^5\text{-C}_5\text{H}_3(\text{CH}_3)\text{COOH}$)₂], [Fe($\eta^5\text{-C}_5\text{H}_4\text{CH}(\text{CH}_3)\text{NH}(\eta^5\text{-C}_5\text{H}_4\text{CH}(\text{CH}_3))$)] and in the diaminecyclohexane salt [Fe($\eta^5\text{-C}_5\text{H}_4\text{COO}$)₂]²⁻[(1*S*,2*S*)-(NH₃)₂C₆H₁₀]²⁺·2[H₂O], 609
- Polymer
- Ferrocene polymers: current architectures, syntheses and utility, 47
- Polysiloxanes
- Functionalization via hydrosilylation of linear and cyclic siloxanes with appendent first generation dendrons containing electronically communicated ferrocenyl units, 642
- Proline
- 1,1'-Ferrocenyl–oligoproline. A synthetic, structural and electrochemical study, 335
- Pt and Pd complexes of dppf
- Synthesis, X-ray, spectroscopic and a preliminary Suzuki coupling screening studies of a complete series of dppfMX₂ (M = Pt, Pd; X = Cl, Br, I), 691
- Push–pull compounds
- Electron acceptors of the fluorene series: Part 12. 9-(Metallocenylidene)nitrofluorene derivatives of Fc–NF, NF–Fc–NF, and NF–Rc–NF types, and the vinylogues Fc– π –NF: synthesis, characterisation, intramolecular charge transfer, redox properties and X-ray structures for three fluorene–ferrocene derivatives, 445
- Pyridinium
- New chiral ferrocenyl-pyridinium salts for non-linear optics, 531
- Pyrrole
- Synthesis and dihydrogen phosphate binding properties of pyrrole containing *ansa*-ferrocenes, 343
- Radicals
- Synthesis of ferrocenylketyl radicals by chromium(II) complexes, 813
- Redox
- Synthesis and characterisation of novel multidentate ferrocene ligands and their Re(I) and Pt(II) complexes, 418
- Synthesis of a new bis(ferrocenyl)ruthenacyclopentatriene compound with a significant inter-metal electronic communication, 80
- Redox-active ligands
- Complexation and electrochemical sensing of anions by amide-substituted ferrocenyl ligands, 356
- Redox behavior
- 1,1'-Ferrocenyl–oligoproline. A synthetic, structural and electrochemical study, 335
- Reduction
- Highly efficient reduction of ferrocenyl derivatives by borane, 364
- Rhenium
- Synthesis and characterisation of novel multidentate ferrocene ligands and their Re(I) and Pt(II) complexes, 418

- Technetium and rhenium heterocomplexes containing the diphenylphosphinoferrocenyl fragment, 772
- Rhodium
Synthesis, reactivity, and crystal structures of ferrocene-substituted amidinate derivatives, 521
- Rhodium and iridium ESR
Synthesis and characterisation of rhodium(0) and rhodium(–1) complexes stabilised by 1,1'-bis(diphenylphosphino)ferrocene (dppf). Crystal structures of [Rh(dppf)₂] and [Na(THF)₅][Rh(dppf)₂].THF, 710
- Rhodium complexes
Synthesis and characterisation of rhodium(0) and rhodium(–1) complexes stabilised by 1,1'-bis(diphenylphosphino)ferrocene (dppf). Crystal structures of [Rh(dppf)₂] and [Na(THF)₅][Rh(dppf)₂].THF, 710
- Ring-closing metathesis
A rapid approach to ferrocenophanes via ring-closing metathesis, 669
- Ring-opening polymerisation
Ferrocene polymers: current architectures, syntheses and utility, 47
- Ruthenium
Allylferrocenylselenide and the synthesis of the first seleno-substituted allenylidene complex: synthesis, spectroscopy, electrochemistry and the effect of electron transfer from the ferrocenylselenyl subunit, 240
- Ruthenium(II) complexes containing 4-ferrocenylphenylisocyanide ligands. Crystal structure of *trans*, *trans*, *trans*-[RuCl₂(POMe-*P*)₂(FcC₆H₄NC)₂] (POMe = PPh₂C₆H₄-OCH₃; Fc = ferrocenyl), 145
- Synthesis of a new bis(ferrocenyl)ruthenacyclopentatriene compound with a significant inter-metal electronic communication, 80
- The reaction of ferrocenyl acetylene with [RuCp(PR₃)(CH₃CN)₂]PF₆ (R = Me, Ph, Cy). Formation of the first allenyl carbene complexes, 70
- Ruthenium complexes
Diferrocenyltriphosphines 2. Reversible phosphine deligation in the chemistry of diferrocenyltriphosphine Ru(II) dichloride complexes with nitriles and pyridines: towards a pH-switchable catalyst?, 538
- Ruthenium(II)
Octahedral ruthenium(II) complexes containing the chiral ligand (4*S*)-2-[(*S_p*)-2-(diphenylphosphino)ferrocenyl]-4-(isopropyl)oxazoline (FcPN). X-Ray crystal structures of *fac*-[RuCl₂(PMe₃)₂(FcPN)] and *fac*-[RuCl₂(dppm)(FcPN)] (dppm = bis(diphenylphosphino)methane), 463
- Ruthenium(IV) complexes
Reactivity of the bis(allyl)-ruthenium(IV) dimer [{Ru(η³:η³-C₁₀H₁₆)(μ-Cl)Cl₂}] (C₁₀H₁₆ = 2,7-dimethylocta-2,6-diene-1,8-diyl) towards dppf (dppf = [Fe(η⁵-C₅H₄PPh₂)₂]) and dppfO (dppfO = [Fe(η⁵-C₅H₄PPh₂)(η⁵-C₅H₄P(=O)Ph₂)]), 767
- S-donor
Chiral ferrocenylthiazolidines, new ligands for palladium complexes, 116
- Selenium
Allylferrocenylselenide and the synthesis of the first seleno-substituted allenylidene complex: synthesis, spectroscopy, electrochemistry and the effect of electron transfer from the ferrocenylselenyl subunit, 240
- Bis[bis(trimethylsilyl)cyclopentadienyl]selene as a ligand: selena-2,2',4,4'-tetrakis(trimethylsilyl)-[1]ferrocenophane and 1,2,3-triseleno-2,2',4,4'-tetrakis(trimethylsilyl)-[3]ferrocenophane, 745
- Self-assembled monolayer
Anion effect on the electrochemical characteristics of a gold electrode modified with a self-assembled monolayer of ferrocenylhexanethiol in aqueous and dichloromethane solutions, 841
- Sensor
The effect of d-block metal complexation on the spectroscopic and redox properties of ferrocene derivatives containing pyridine ligands, 304
- Sensors
Voltammetric metal cation sensors based on ferrocene derivatives with oxazoline and imine substituents, 134
- Silylation
ortho-Silylation of 2,2'-bis(oxazolyl)-1,1'-bis(diphenylphosphino)ferrocenes and remarkable effect of the silyl groups on the enantioselectivity in Pd-catalyzed asymmetric allylic alkylation, 99
- Solid-phase
Synthesis and characterization of ferrocene-labeled oligodeoxynucleotides, 398
- Solid-state reaction
Preparation of 1,5-diketone derivatives containing ferrocenyl by Michael reaction under solvent-free condition, 719
- Preparation of ferrocenyl mono- and dienone derivatives through aldol condensation of 1,1'-diacetylferrocene with aromatic aldehydes in dry conditions, 782
- Solvent-free
Preparation of 1,5-diketone derivatives containing ferrocenyl by Michael reaction under solvent-free condition, 719
- Sonogashira coupling
Star-shaped polyferrocenes based on thiophene and triphenylamine: synthesis, spectroscopy and electrochemistry, 139
- Spectro-electrochemistry
Allylferrocenylselenide and the synthesis of the first seleno-substituted allenylidene complex: synthesis, spectroscopy, electrochemistry and the effect of electron transfer from the ferrocenylselenyl subunit, 240
- Spectroscopy
The effect of d-block metal complexation on the spectroscopic and redox properties of ferrocene derivatives containing pyridine ligands, 304
- Spirocycles
Reactions of ferrocene-derived bis(hydroxymethyl) phosphine sulfides FcCH(R)P(S)(CH₂OH)₂ (R = H, CH₃) with cyclic thionylphosphazenes: crystal structures of FcCH₂P(S)(CH₂O)₂PN(PCl₂)[NS(O)Ph] and FcCH₂P(S)(CH₂O)₂PN₂P[N(Me)CH₂]₂[NS(O)Ph] (Fc = ferrocenyl), 382
- π-Stacking
Synthesis and crystal structure of 1,1'-bis[4-(2,2':6',2''-terpyridin-4'-yl)phenyl]octamethylferrocene, 733
- Star-shaped complexes
Facile synthesis of bidimensional ferrocenyl-based branched oligomers by palladium-catalyzed coupling reactions, 191
- Stereochemistry
Formation and structural characterization of amino-substituted [3]ferrocenophanes derived from intramolecular Mannich-type coupling reactions at the metallocene framework, 621
- Studies on the cyclometallation of ferrocenylimines, 27
- Stereoconservative
Study on the synthesis of nonracemic C₂-symmetric 1,1'-binaphthyl-2,2'-diyl bridged ferrocene. Stereochemical result of the cross-coupling reactions controlled by Pd(II) or Pd(IV) complex intermediacy, 318
- Stereospecific substitution
A rapid approach to ferrocenophanes via ring-closing metathesis, 669

- Stille coupling
 Synthesis of 1',1''-bis(heterocyclyl)biferrocene compounds, 723
- Structure–property relationships
 Studies on the cyclometallation of ferrocenylimines, 27
- Supporting electrolyte
 Improved electrochemistry of multi-ferrocenyl compounds: investigation of biferrocene, terferrocene, bis(fulvalene)diiron and diferrocenylolethane in dichloromethane using [NBu₄][B(C₆F₅)₄] as supporting electrolyte, 823
- Supramolecular assembly
 Second harmonic generation in ferrocene based hydrogen bonded assemblies, 827
- Surfactants
 Enzymatic chemistry of ferrocenes: micellar tuning of the glucose oxidase reactivity toward solubilized electrochemically generated *n*-alkylferrocenium cations, 469
- Suzuki coupling
 Short syntheses of 8-substituted 8'-[1-(1'-phenylthio)ferrocenyl]-1,1'-binaphthyls from Suzuki coupling reactions. A strategy for generating new chiral ligands and charge-transfer complexes, 832
 Synthesis, X-ray, spectroscopic and a preliminary Suzuki coupling screening studies of a complete series of dppfMX₂ (M = Pt, Pd; X = Cl, Br, I), 691
- Synthesis
 Ferrocene derivatives (II). Synthesis and reactions of 4-ferrocenyl-2-thiazolamine, 742
 Internal ferrocenylalkynes—a comparative electrochemical and mass spectrometric study, 291
 Properties of some ferrocene macrocyclic dioxotetraamines: the roles of aromatic side-arms, 327
 Synthesis and characterization of ferrocene-labeled oligodeoxynucleotides, 398
- Tamoxifen
 Studies on organometallic selective estrogen receptor modulators. (SERMs) Dual activity in the hydroxy-ferrocifen series, 500
- Technetium
 Technetium and rhenium heterocomplexes containing the diphenylphosphinoferrrocenyl fragment, 772
- Thiazolidines
 Chiral ferrocenylthiazolidines, new ligands for palladium complexes, 116
- Thiazolidone
 Ferrocene derivatives (II). Synthesis and reactions of 4-ferrocenyl-2-thiazolamine, 742
- Thiazolines
 Ferrocene derivatives (II). Synthesis and reactions of 4-ferrocenyl-2-thiazolamine, 742
- Thioacetate
 Palladium(II) phosphine thiocarboxylates. Structures of *cis*-Pd(κ S-SOCMe)₂(dppf) [dppf = 1,1'-bis(diphenylphosphino)ferrocene] and *trans*-Pd(κ S-SOCMe)₂(PPh₃)₂, 757
- Thioacylsilanes
 Synthesis and reactivity of achiral and of a novel planar chiral thioferrocenoylsilanes, 407
- Thioether
 Synthesis and characterisation of novel multidentate ferrocene ligands and their Re(I) and Pt(II) complexes, 418
- Thionylphosphazene
 Reactions of ferrocene-derived bis(hydroxymethyl) phosphine sulfides FcCH(R)P(S)(CH₂OH)₂ (R = H, CH₃) with cyclic thionylphosphazenes: crystal structures of FcCH₂P(S)-(CH₂O)₂PN(NPCl₂)₂[NS(O)Ph] and FcCH₂P(S)(CH₂O)₂-PN₂P[N(Me)CH₂]₂[NS(O)Ph] (Fc = ferrocenyl), 382
- Thiophene
 Star-shaped polyferrocenes based on thiophene and triphenylamine: synthesis, spectroscopy and electrochemistry, 139
 The linear and nonlinear optical properties of organometallic chromophores derived from ferrocene, [Fe₂(η ⁵-C₅H₅)₂(CO)₂(μ -CO)(μ -C-CH₃)]⁺[BF₄]⁻ and terthienyl spacers. Crystal structure of 2-[(*E*)-2-ferrocenylethenyl]-5-(2-thienyl)thiophene, 435
- Thiophilic addition
 Synthesis and reactivity of achiral and of a novel planar chiral thioferrocenoylsilanes, 407
- Three-component reaction
 (η ⁵-C₅H₅)-Ring alkylation reaction with the C₅H₅ anion: towards the construction of tri-Fe complex Fe{[μ , η ⁵: η ⁴-5-*exo*-(1'-C₃H₄)C₅H₅]Fe(CO)₂(PPh₃)₂}, 549
- Transmetalation
 Studies on the cyclometallation of ferrocenylimines, 27
- Triarylamine
 Star-shaped polyferrocenes based on thiophene and triphenylamine: synthesis, spectroscopy and electrochemistry, 139
- Triferrocenyl derivatives
 An investigation of the chemistry of ferrocenoyl derivatives. The synthesis and reactions of ferrocenoyl imidazolide and its derivatives, 266
- Tungsten
 Syntheses and some reactions of complexes containing carbon chains capped by ferrocenyl and W(CO)₃Cp groups, 484
- Unsymmetrical 8,8'-disubstituted 1,1'-binaphthyls
 Short syntheses of 8-substituted 8'-[1-(1'-phenylthio)ferrocenyl]-1,1'-binaphthyls from Suzuki coupling reactions. A strategy for generating new chiral ligands and charge-transfer complexes, 832
- UV–vis spectrometry
 Properties of some ferrocene macrocyclic dioxotetraamines: the roles of aromatic side-arms, 327
- Vinylidene complexes
 The reaction of ferrocenyl acetylene with [RuCp(PR₃)-(CH₃CN)₂]PF₆ (R = Me, Ph, Cy). Formation of the first allenyl carbene complexes, 70
- Water solubility
 Water soluble ferrocenyl and polyferrocenyl compounds: synthesis and electrochemistry, 595
- X-ray
 Synthesis of new ferrocene containing diamines and their use in epoxy resins, 837
 Synthesis, X-ray, spectroscopic and a preliminary Suzuki coupling screening studies of a complete series of dppfMX₂ (M = Pt, Pd; X = Cl, Br, I), 691
- X-ray crystal structure analysis
 Multistep redox properties of 2,2'-bipyridylboronium substituted ferrocenes, 390
- X-ray diffraction
 Electron acceptors of the fluorene series: Part 12. 9-(Metalloenylidene)nitrofluorene derivatives of Fc-NF, NF-Fc-NF, and NF-Rc-NF types, and the vinylogues Fc- π -NF: synthesis, characterisation, intramolecular charge transfer, redox properties and X-ray structures for three fluorene-ferrocene derivatives, 445
- X-ray structure
 Ferrocene derivatives (I). Synthesis and X-ray structure determination of *N*-*o*-methylphenylferrocenesulfonamide, 738
- X-ray structure analysis
 4-Ethynyl-benzonitrile-ferrocenes bridged by a Pd(PPh₃)₂ unit; the

- solid-state structure of $(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\eta^5\text{-C}_5\text{H}_4\text{C}\equiv\text{CC}_6\text{H}_4\text{C}\equiv\text{N-1,4})$, 198
- X-Ray structures
- Semimasked 1,1'-diethynylferrocenes: synthetic concepts, preparations, and reactions, 558
- Zero-valent complexes
- Synthesis and characterisation of rhodium(0) and rhodium(–1) complexes stabilised by 1,1'-bis(diphenylphosphino)ferrocene (dppf). Crystal structures of $[\text{Rh}(\text{dppf})_2]$ and $[\text{Na}(\text{THF})_5][\text{Rh}(\text{dppf})_2]\cdot\text{THF}$, 710