

Subject Index of Volume 635

Alkane hydroxylation

Methane inversion on transition metal ions: a possible mechanism for stereochemical scrambling in metal-catalyzed alkane hydroxylations, 100

Alkene complexes

Olefin strain energies and platinum complexes of highly pyramidalised alkenes, 142

Amide

Bis-amide and amine bis-amide ligands in M(IV) (M = V, Cr, Mn) based olefin polymerization catalysts: a theoretical study, 153

Amine

Bis-amide and amine bis-amide ligands in M(IV) (M = V, Cr, Mn) based olefin polymerization catalysts: a theoretical study, 153

Back bonding

Back bonding without σ -bonding: a unique π -complex of dinitrogen with uranium, 69

Backdonation

Metal–dihydrogen and σ -bond coordination: the consummate extension of the Dewar–Chatt–Duncanson model for metal–olefin π bonding, 37

σ -Bond coordination

Metal–dihydrogen and σ -bond coordination: the consummate extension of the Dewar–Chatt–Duncanson model for metal–olefin π bonding, 37

Bonding in π -complexes

A historical perspective on Dewar's landmark contribution to organometallic chemistry, 1

Bond theory

New types of non-classical interligand interactions involving silicon based ligands, 24

Boranes

Analogies between the chemical bonding topologies in metal–olefin complexes and in metallaboranes: the role of three-center two-electron bonding, 75

Bridging alkynes

Reactions of the μ -alkyne-dicobalt complexes $[\text{Co}_2(\text{CO})_6(\mu\text{-CF}_3\text{-C}\equiv\text{C-R})]$ (R = CF_3 , H) with $[\text{Co}_2\text{Cp}_2(\mu\text{-SMe})_2]$: substitution and insertion leading to novel thiolato-alkyne tetra- and tricobalt clusters, 110

Butadiyne complexes

Structure and bonding of metallocyclocumulenes, radialenes, butadiyne complexes and their possible interconversion: a theoretical study, 204

Carbonyl clusters

Reactions of $[\text{Os}_3(\text{CO})_{10}(\text{NCMe})_2]$ with symmetric diynes and their reactivity towards $[\text{Co}_2(\text{CO})_8]$, 119

Carbonyls

Understanding the nature of the bonding in transition metal complexes: from Dewar's molecular orbital model to an energy partitioning analysis of the metal–ligand bond, 9

Catalysis

Hybrid QM/MM study of propene insertion into the Rh–H bond of $\text{HRh}(\text{PPh}_3)_2(\text{CO})(\eta^2\text{-CH}_2=\text{CHCH}_3)$: the role of the olefin adduct in determining product selectivity, 132

Chemical bonding topology

Analogies between the chemical bonding topologies in metal–olefin complexes and in metallaboranes: the role of three-center two-electron bonding, 75

Chromium

Bis-amide and amine bis-amide ligands in M(IV) (M = V, Cr, Mn) based olefin polymerization catalysts: a theoretical study, 153

Cobalt

Bis(indenyl) complexes of Fe, Co, and Ni: electronic structure and preferences, 197

Metal vapour synthesis of phospho-organometallic compounds via reaction of cobalt atoms with the phospho-alkyne $\text{t-BuC}\equiv\text{P}$: synthesis and structural characterisation of the sandwich compounds $[\text{Co}(\eta^5\text{-P}_3\text{C}_2\text{Bu}_2)(\eta^4\text{-P}_2\text{C}_2\text{Bu}_2)]$, $[\text{Co}(\eta^5\text{-P}_2\text{C}_3\text{Bu}_3)(\eta^4\text{-P}_2\text{C}_2\text{Bu}_2)]$ and the protonated tetraphospha-barrele complex $[\text{Co}(\eta^4\text{-P}_4\text{C}_4\text{Bu}_4\text{H})(\eta^4\text{-P}_2\text{C}_2\text{Bu}_2)]$, the latter as its $[\text{W}(\text{CO})_5]$ adduct, 212

Reactions of $[\text{Os}_3(\text{CO})_{10}(\text{NCMe})_2]$ with symmetric diynes and their reactivity towards $[\text{Co}_2(\text{CO})_8]$, 119

Reactions of the μ -alkyne-dicobalt complexes $[\text{Co}_2(\text{CO})_6(\mu\text{-CF}_3\text{-C}\equiv\text{C-R})]$ (R = CF_3 , H) with $[\text{Co}_2\text{Cp}_2(\mu\text{-SMe})_2]$: substitution and insertion leading to novel thiolato-alkyne tetra- and tricobalt clusters, 110

Competition for π -backdonation

Structural preference in complexes containing both double-face and single-face π -acceptor ligands, 84

π -Complexes

Back bonding without σ -bonding: a unique π -complex of dinitrogen with uranium, 69

Coordinate link

A historical perspective on Dewar's landmark contribution to organometallic chemistry, 1

Dative bond

A historical perspective on Dewar's landmark contribution to organometallic chemistry, 1

Density functional

Olefin strain energies and platinum complexes of highly pyramidalised alkenes, 142

Density functional theory

A density functional study of the structures, vibrations and bond energies of dinitrogen phosphine complexes of the first transition series, 165

Dewar–Chatt–Duncanson

A historical perspective on Dewar's landmark contribution to organometallic chemistry, 1

Dewar–Chatt–Duncanson model

Metal–dihydrogen and σ -bond coordination: the consummate extension of the Dewar–Chatt–Duncanson model for metal–olefin π bonding, 37

- DFT**
 Bis-amide and amine bis-amide ligands in M(IV) (M = V, Cr, Mn) based olefin polymerization catalysts: a theoretical study, 153
 Hybrid QM/MM study of propene insertion into the Rh–H bond of $\text{HRh}(\text{PPh}_3)_2(\text{CO})(\eta^2\text{-CH}_2=\text{CHCH}_3)$: the role of the olefin adduct in determining product selectivity, 132
 M_2E_2 four-member ring structure, $\text{M}_2(\mu\text{-EH}_2)_2(\text{P}2)_2$ (M = Pd or Pt; E = Si or Ge; P2 = $(\text{PH}_3)_2$ or $\text{H}_2\text{PCH}_2\text{CH}_2\text{PH}_2$) versus μ -disilene and μ -digermene-bridged structures, $\text{M}_2(\mu\text{-E}_2\text{H}_4)(\text{P}2)_2$. A theoretical study, 173
 Understanding the nature of the bonding in transition metal complexes: from Dewar's molecular orbital model to an energy partitioning analysis of the metal–ligand bond, 9
- DFT calculations**
 Bis(indenyl) complexes of Fe, Co, and Ni: electronic structure and preferences, 197
- Dinitrogen**
 Back bonding without σ -bonding: a unique π -complex of dinitrogen with uranium, 69
- Dinitrogen complexes**
 A density functional study of the structures, vibrations and bond energies of dinitrogen phosphine complexes of the first transition series, 165
- Diyne**
 Reactions of $[\text{Os}_3(\text{CO})_{10}(\text{NCMe})_2]$ with symmetric diynes and their reactivity towards $[\text{Co}_2(\text{CO})_8]$, 119
- Electronic structures**
 A density functional study of the structures, vibrations and bond energies of dinitrogen phosphine complexes of the first transition series, 165
- Electron population analysis**
 Electronic structure and spectra of ruthenium diimine complexes by density functional theory and INDO/S. Comparison of the two methods, 187
- μ -Germene**
 M_2E_2 four-member ring structure, $\text{M}_2(\mu\text{-EH}_2)_2(\text{P}2)_2$ (M = Pd or Pt; E = Si or Ge; P2 = $(\text{PH}_3)_2$ or $\text{H}_2\text{PCH}_2\text{CH}_2\text{PH}_2$) versus μ -disilene and μ -digermene-bridged structures, $\text{M}_2(\mu\text{-E}_2\text{H}_4)(\text{P}2)_2$. A theoretical study, 173
- Germylene**
 M_2E_2 four-member ring structure, $\text{M}_2(\mu\text{-EH}_2)_2(\text{P}2)_2$ (M = Pd or Pt; E = Si or Ge; P2 = $(\text{PH}_3)_2$ or $\text{H}_2\text{PCH}_2\text{CH}_2\text{PH}_2$) versus μ -disilene and μ -digermene-bridged structures, $\text{M}_2(\mu\text{-E}_2\text{H}_4)(\text{P}2)_2$. A theoretical study, 173
- Group 13-diyl**
 Understanding the nature of the bonding in transition metal complexes: from Dewar's molecular orbital model to an energy partitioning analysis of the metal–ligand bond, 9
- Hyrido complexes**
 New types of non-classical interligand interactions involving silicon based ligands, 24
- Hydroformylation**
 Hybrid QM/MM study of propene insertion into the Rh–H bond of $\text{HRh}(\text{PPh}_3)_2(\text{CO})(\eta^2\text{-CH}_2=\text{CHCH}_3)$: the role of the olefin adduct in determining product selectivity, 132
- Hypervalent compounds**
 New types of non-classical interligand interactions involving silicon based ligands, 24
- Indenyl effect**
 Bis(indenyl) complexes of Fe, Co, and Ni: electronic structure and preferences, 197
- Iron**
 Bis(indenyl) complexes of Fe, Co, and Ni: electronic structure and preferences, 197
- Manganese**
 Bis-amide and amine bis-amide ligands in M(IV) (M = V, Cr, Mn) based olefin polymerization catalysts: a theoretical study, 153
- Metal–alkene complexes**
 Structural preference in complexes containing both double-face and single-face π -acceptor ligands, 84
- Metal–alkenyl complexes**
 Structural preference in complexes containing both double-face and single-face π -acceptor ligands, 84
- Metal–borane complex**
 Metal–dihydrogen and σ -bond coordination: the consummate extension of the Dewar–Chatt–Duncanson model for metal–olefin π bonding, 37
- Metal–boryl complexes**
 Structural preference in complexes containing both double-face and single-face π -acceptor ligands, 84
- Metal–dihydrogen complex**
 Metal–dihydrogen and σ -bond coordination: the consummate extension of the Dewar–Chatt–Duncanson model for metal–olefin π bonding, 37
- Metallaboranes**
 The relevance of the Dewar model to neighboring p-block element analogs of metal hydrocarbon π -complexes, 92
- Metallacyclocumulenes**
 Structure and bonding of metallacyclocumulenes, radialenes, butadiene complexes and their possible interconversion: a theoretical study, 204
- Metallaphosphanes**
 The relevance of the Dewar model to neighboring p-block element analogs of metal hydrocarbon π -complexes, 92
- Metallsilanes**
 The relevance of the Dewar model to neighboring p-block element analogs of metal hydrocarbon π -complexes, 92
- Metal–ligand covalency**
 Electronic structure and spectra of ruthenium diimine complexes by density functional theory and INDO/S. Comparison of the two methods, 187
- Metallaboranes**
 Analogies between the chemical bonding topologies in metal–olefin complexes and in metallaboranes: the role of three-center two-electron bonding, 75
- Metallocenes**
 Bis(indenyl) complexes of Fe, Co, and Ni: electronic structure and preferences, 197
- Metal–olefin π bonding**
 Metal–dihydrogen and σ -bond coordination: the consummate extension of the Dewar–Chatt–Duncanson model for metal–olefin π bonding, 37
- Metal–silane complex**
 Metal–dihydrogen and σ -bond coordination: the consummate extension of the Dewar–Chatt–Duncanson model for metal–olefin π bonding, 37
- Metal–silane complexes**
 Structural preference in complexes containing both double-face and single-face π -acceptor ligands, 84
- Metal vapour synthesis**
 Metal vapour synthesis of phospho-organometallic compounds via reaction of cobalt atoms with the phospho-alkyne $\text{tBuC}\equiv\text{P}$: synthesis and structural characterisation of the sandwich compounds $[\text{Co}(\eta^5\text{-P}_3\text{C}_2\text{Bu}_2)(\eta^4\text{-P}_2\text{C}_2\text{Bu}_2)]$, $[\text{Co}(\eta^5\text{-P}_3\text{C}_2\text{Bu}_3)(\eta^4\text{-P}_2\text{C}_2\text{Bu}_2)]$ and the protonated tetraphospha-barrelene complex $[\text{Co}(\eta^4\text{-P}_4\text{C}_4\text{Bu}_4\text{H})(\eta^4\text{-P}_2\text{C}_2\text{Bu}_2)]$, the latter as its $[\text{W}(\text{CO})_5]$ adduct, 212

Methane activation

Methane inversion on transition metal ions: a possible mechanism for stereochemical scrambling in metal-catalyzed alkane hydroxylations, 100

Molecular orbital

Olefin strain energies and platinum complexes of highly pyramidalised alkenes, 142

Molecular orbital calculation

Bis-amide and amine bis-amide ligands in M(IV) (M = V, Cr, Mn) based olefin polymerization catalysts: a theoretical study, 153

Molecular orbital calculations

Understanding the nature of the bonding in transition metal complexes: from Dewar's molecular orbital model to an energy partitioning analysis of the metal–ligand bond, 9

Molecular orbital theory

Methane inversion on transition metal ions: a possible mechanism for stereochemical scrambling in metal-catalyzed alkane hydroxylations, 100

Molecules orbital calculations

M₂E₂ four-member ring structure, M₂(μ-EH₂)₂(P₂)₂ (M = Pd or Pt; E = Si or Ge; P₂ = (PH₃)₂ or H₂PCH₂CH₂PH₂) versus μ-disilene and μ-digermene-bridged structures, M₂(μ-E₂H₄)(P₂)₂. A theoretical study, 173

Nickel

Bis(indenyl) complexes of Fe, Co, and Ni: electronic structure and preferences, 197

Nonclassical bonding

Metal–dihydrogen and σ-bond coordination: the consummate extension of the Dewar–Chatt–Duncanson model for metal–olefin π bonding, 37

Non-classical interactions

New types of non-classical interligand interactions involving silicon based ligands, 24

Olefin polymerization

Bis-amide and amine bis-amide ligands in M(IV) (M = V, Cr, Mn) based olefin polymerization catalysts: a theoretical study, 153

Osmium

Reactions of [Os₃(CO)₁₀(NCMe)₂] with symmetric diynes and their reactivity towards [Co₂(CO)₈], 119

Oxidative addition

Metal–dihydrogen and σ-bond coordination: the consummate extension of the Dewar–Chatt–Duncanson model for metal–olefin π bonding, 37

Palladium

M₂E₂ four-member ring structure, M₂(μ-EH₂)₂(P₂)₂ (M = Pd or Pt; E = Si or Ge; P₂ = (PH₃)₂ or H₂PCH₂CH₂PH₂) versus μ-disilene and μ-digermene-bridged structures, M₂(μ-E₂H₄)(P₂)₂. A theoretical study, 173

Phospha-alkyne

Metal vapour synthesis of phospha-organometallic compounds via reaction of cobalt atoms with the phospha-alkyne 'BuC≡P: synthesis and structural characterisation of the sandwich compounds [Co(η⁵-P₃C₂Bu₂)(η⁴-P₂C₂Bu₂)], [Co(η⁵-P₂C₃Bu₃)(η⁴-P₂C₂Bu₂)] and the protonated tetraphospha-barrele complex [Co(η⁴-P₄C₄Bu₄H)(η⁴-P₂C₂Bu₂)], the latter as its [W(CO)₅] adduct, 212

Platinum

M₂E₂ four-member ring structure, M₂(μ-EH₂)₂(P₂)₂ (M = Pd or Pt; E = Si or Ge; P₂ = (PH₃)₂ or H₂PCH₂CH₂PH₂) versus μ-disilene and μ-digermene-bridged structures, M₂(μ-E₂H₄)(P₂)₂. A theoretical study, 173

QM/MM

Hybrid QM/MM study of propene insertion into the Rh–H bond of HRh(PPh₃)₂(CO)(η²-CH₂=CHCH₃): the role of the olefin adduct in determining product selectivity, 132

Radialenes

Structure and bonding of metallacyclocumulenes, radialenes, butadiene complexes and their possible interconversion: a theoretical study, 204

Reactivity

A density functional study of the structures, vibrations and bond energies of dinitrogen phosphine complexes of the first transition series, 165

Regioselectivity

Hybrid QM/MM study of propene insertion into the Rh–H bond of HRh(PPh₃)₂(CO)(η²-CH₂=CHCH₃): the role of the olefin adduct in determining product selectivity, 132

Rhodium

Hybrid QM/MM study of propene insertion into the Rh–H bond of HRh(PPh₃)₂(CO)(η²-CH₂=CHCH₃): the role of the olefin adduct in determining product selectivity, 132

Sandwich compounds

Metal vapour synthesis of phospha-organometallic compounds via reaction of cobalt atoms with the phospha-alkyne 'BuC≡P: synthesis and structural characterisation of the sandwich compounds [Co(η⁵-P₃C₂Bu₂)(η⁴-P₂C₂Bu₂)], [Co(η⁵-P₂C₃Bu₃)(η⁴-P₂C₂Bu₂)] and the protonated tetraphospha-barrele complex [Co(η⁴-P₄C₄Bu₄H)(η⁴-P₂C₂Bu₂)], the latter as its [W(CO)₅] adduct, 212

μ-Silene

M₂E₂ four-member ring structure, M₂(μ-EH₂)₂(P₂)₂ (M = Pd or Pt; E = Si or Ge; P₂ = (PH₃)₂ or H₂PCH₂CH₂PH₂) versus μ-disilene and μ-digermene-bridged structures, M₂(μ-E₂H₄)(P₂)₂. A theoretical study, 173

Silicon

New types of non-classical interligand interactions involving silicon based ligands, 24

Silylene

M₂E₂ four-member ring structure, M₂(μ-EH₂)₂(P₂)₂ (M = Pd or Pt; E = Si or Ge; P₂ = (PH₃)₂ or H₂PCH₂CH₂PH₂) versus μ-disilene and μ-digermene-bridged structures, M₂(μ-E₂H₄)(P₂)₂. A theoretical study, 173

Stereochemical scrambling

Methane inversion on transition metal ions: a possible mechanism for stereochemical scrambling in metal-catalyzed alkane hydroxylations, 100

Strain energies

Olefin strain energies and platinum complexes of highly pyramidalised alkenes, 142

Synergic bonding

A historical perspective on Dewar's landmark contribution to organometallic chemistry, 1

Understanding the nature of the bonding in transition metal complexes: from Dewar's molecular orbital model to an energy partitioning analysis of the metal–ligand bond, 9

Three-center bonds

Analogies between the chemical bonding topologies in metal–olefin complexes and in metallaboranes: the role of three-center two-electron bonding, 75

Time-dependent density-functional response theory

Electronic structure and spectra of ruthenium diimine complexes by density functional theory and INDO/S. Comparison of the two methods, 187

Titanocene

Structure and bonding of metallacyclocumulenes, radialenes, butadiene complexes and their possible interconversion: a theoretical study, 204

Trans-effect

Metal–dihydrogen and σ -bond coordination: the consummate extension of the Dewar–Chatt–Duncanson model for metal–olefin π bonding, 37

Triangular clusters

Reactions of the μ -alkyne-dicobalt complexes $[\text{Co}_2(\text{CO})_6(\mu\text{-CF}_3\text{-C}\equiv\text{C-R})]$ ($\text{R} = \text{CF}_3, \text{H}$) with $[\text{Co}_2\text{Cp}_2(\mu\text{-SMe})_2]$: substitution and insertion leading to novel thiolato-alkyne tetra- and tricobalt clusters, 110

Uranium

Back bonding without σ -bonding: a unique π -complex of dinitrogen with uranium, 69

Vanadium

Bis-amide and amine bis-amide ligands in M(IV) ($\text{M} = \text{V}, \text{Cr}, \text{Mn}$) based olefin polymerization catalysts: a theoretical study, 153

X-ray crystal structures

Reactions of $[\text{Os}_3(\text{CO})_{10}(\text{NCMe})_2]$ with symmetric diynes and their reactivity towards $[\text{Co}_2(\text{CO})_8]$, 119

X-ray structures

Reactions of the μ -alkyne-dicobalt complexes $[\text{Co}_2(\text{CO})_6(\mu\text{-CF}_3\text{-C}\equiv\text{C-R})]$ ($\text{R} = \text{CF}_3, \text{H}$) with $[\text{Co}_2\text{Cp}_2(\mu\text{-SMe})_2]$: substitution and insertion leading to novel thiolato-alkyne tetra- and tricobalt clusters, 110

ZINDO

Electronic structure and spectra of ruthenium diimine complexes by density functional theory and INDO/S. Comparison of the two methods, 187

Zirconocene

Structure and bonding of metallacyclocumulenes, radialenes, butadiyne complexes and their possible interconversion: a theoretical study, 204