

Subject Index of Volume 640

Allenylation

The deviating behavior of thiols in nucleophilic trapping reactions of chromiumcarbonyl phenyl complex substituted propargyl cation, 41

Allyl

Komplekatalyse: LIX. Die Katalyse der 1,4-*trans*-Polymerisation des Butadiens mit Tris(allyl)neodym(III) $\text{Nd}(\eta^3\text{-C}_3\text{H}_5)_3$ als Einkomponentenkatalysator—Kinetik und Reaktionsmechanismus, 1

Allylic alkylation

Synthesis of chiral 2-alkyl-8-quinolinyloxazoline ligands: reversal of enantioselectivity in the asymmetric palladium-catalyzed allylic alkylation, 65

Anchimeric assistance

Anchimeric assistance by γ -substituents Z, Z = MeO, PhO, MeS or PhS, in reactions of the bromides $(\text{Me}_3\text{Si})_2(\text{ZMe}_2\text{Si})\text{-CSiMe}_2\text{Br}$ with AgBF_4 , 29

Arene complexes

The deviating behavior of thiols in nucleophilic trapping reactions of chromiumcarbonyl phenyl complex substituted propargyl cation, 41

Arylamine ligands

A new homobimetallic arenediyl diplatinum(II) unit as building block for macromolecular synthesis. X-ray crystal structure of $[\text{C}_6\text{H}_2\{\text{CH}_2\text{NMe}_2\}_2\text{-1,5-}\{\text{PtCl}(\text{PPh}_3)\}_2\text{-2,4}]$, 166

Arylboronic esters

Bis(diphosphaferrocene) palladium(II) dimer complexes as efficient catalysts in the synthesis of arylboronic esters, 197

Asymmetric catalysis

Synthesis of chiral 2-alkyl-8-quinolinyloxazoline ligands: reversal of enantioselectivity in the asymmetric palladium-catalyzed allylic alkylation, 65

Bimetallic complexes

Synthesis of metallocenes of zirconium, hafnium, manganese, iron, tin, lead and half-sandwich complexes of rhodium and iridium containing the ligands $(\eta\text{-C}_5\text{R}_4\text{CR}'_2\text{PMe}_2)$, where R and R' may be H or Me, 93

Building blocks

A new homobimetallic arenediyl diplatinum(II) unit as building block for macromolecular synthesis. X-ray crystal structure of $[\text{C}_6\text{H}_2\{\text{CH}_2\text{NMe}_2\}_2\text{-1,5-}\{\text{PtCl}(\text{PPh}_3)\}_2\text{-2,4}]$, 166

Butadiene

Komplekatalyse: LIX. Die Katalyse der 1,4-*trans*-Polymerisation des Butadiens mit Tris(allyl)neodym(III) $\text{Nd}(\eta^3\text{-C}_3\text{H}_5)_3$ als Einkomponentenkatalysator—Kinetik und Reaktionsmechanismus, 1

Butadiene ligand

Interaction of half-sandwich alkylmolybdenum(III) complexes with $\text{B}(\text{C}_6\text{F}_5)_3$. The X-ray structure of $[\text{CpMo}(\eta^4\text{-C}_4\text{H}_6)(\mu\text{-Cl})(\mu\text{-CH}_2)(\text{O})\text{MoCp}][\text{CH}_3\text{B}(\text{C}_6\text{F}_5)_3]$, 113

Carbon monoxide

Selectivity for the methoxycarbonylation of ethylene versus

CO–ethylene copolymerization with catalysts based on C_4 -bridged bidentate phosphines and phospholes, 182

Catalysis

Bis(diphosphaferrocene) palladium(II) dimer complexes as efficient catalysts in the synthesis of arylboronic esters, 197

Komplekatalyse: LIX. Die Katalyse der 1,4-*trans*-Polymerisation des Butadiens mit Tris(allyl)neodym(III) $\text{Nd}(\eta^3\text{-C}_3\text{H}_5)_3$ als Einkomponentenkatalysator—Kinetik und Reaktionsmechanismus, 1

CD spectra

Synthesis and circular dichroism spectra of silver(I) complexes with *R,R*-DIOP (4*R*,5*R*-*trans*-4,5-bis[(diphenylphosphino)methyl]-2,2-dimethyl-1,3-dioxalane): crystal structures of $[\text{AgCl}(\text{R,R-DIOP})_2\cdot 2\text{CHCl}_3]$, $\{[\text{AgBr}(\text{R,R-DIOP})_2\cdot 2\text{CH}_2\text{Cl}_2\cdot 2\text{H}_2\text{O}]\}$, $[\text{AgI}(\text{R,R-DIOP})_2]$ and $[\text{AgSCN}(\text{R,R-DIOP})_2]$, 57

C–F bond activation

Density functional studies of the reactivity in the C–F bond activation of fluoromethane by bare lanthanum monocation, 121

C–H activation

A new homobimetallic arenediyl diplatinum(II) unit as building block for macromolecular synthesis. X-ray crystal structure of $[\text{C}_6\text{H}_2\{\text{CH}_2\text{NMe}_2\}_2\text{-1,5-}\{\text{PtCl}(\text{PPh}_3)\}_2\text{-2,4}]$, 166

Chelating cyclopentadienyl-phosphine complexes

Synthesis of metallocenes of zirconium, hafnium, manganese, iron, tin, lead and half-sandwich complexes of rhodium and iridium containing the ligands $(\eta\text{-C}_5\text{R}_4\text{CR}'_2\text{PMe}_2)$, where R and R' may be H or Me, 93

Chelating cyclopentadienyl-phosphine ligands

Synthesis of metallocenes of zirconium, hafnium, manganese, iron, tin, lead and half-sandwich complexes of rhodium and iridium containing the ligands $(\eta\text{-C}_5\text{R}_4\text{CR}'_2\text{PMe}_2)$, where R and R' may be H or Me, 93

Chirality

Syntheses and pharmacological characterization of achiral and chiral enantiopure C/Si/Ge-analogous derivatives of the muscarinic antagonist cycrimine: a study on C/Si/Ge bioisosterism, 140

Chiral ligands

Synthesis of chiral 2-alkyl-8-quinolinyloxazoline ligands: reversal of enantioselectivity in the asymmetric palladium-catalyzed allylic alkylation, 65

Chromium

Alkylaminophosphanyl substituted half-sandwich complexes of vanadium(III) and chromium(III): preparation and reactivity in ethylene polymerisation, 21

The deviating behavior of thiols in nucleophilic trapping reactions of chromiumcarbonyl phenyl complex substituted propargyl cation, 41

CO-factored force field

Force constant calculations for octahedral complexes of the type $\text{M}(\text{CO})_5\text{L}$, based on the CO-factored force field, 37

Conjugated diimine

Iron carbonyl complexes from 2-[2,3-diaza-4-(2-thienyl)buta-1,3-dienyl]thiophene: N–N bond cleavage and cyclometalation, 85

Copolymerization

Selectivity for the methoxycarbonylation of ethylene versus CO–ethylene copolymerization with catalysts based on C₄-bridged bidentate phosphines and phospholes, 182

Cotton effect

Synthesis and circular dichroism spectra of silver(I) complexes with *R,R*-DIOP (4*R*,5*R*-*trans*-4,5-bis[(diphenylphosphino)methyl]-2,2-dimethyl-1,3-dioxalane): crystal structures of [AgCl(*R,R*-DIOP)]₂·2CHCl₃, {[AgBr(*R,R*-DIOP)]₂·CH₂Cl₂·2H₂O}, [AgI(*R,R*-DIOP)]₂ and [AgSCN(*R,R*-DIOP)]₂, 57

Crystal structures

Cation–anion interactions in triphenyl telluronium salts. The crystal structures of (Ph₃Te)₂[MCl₆] (M = Pt, Ir), (Ph₃Te)-[AuCl₄], and (Ph₃Te)(NO₃)·HNO₃, 50

New ligand substitution modes of M₂(μ-SR)₂(CO)₈ (M = Mo, W; R = alkyl, aryl): synthesis, structure and properties of new di-nuclear metal(I) carbonyl thiolates, [M₂(μ-SPh)₂(CO)_xL_y]ⁿ⁻ (M = Mo, W; x = 6, 8; y = 0, 1; L = PhS, (Ph)₂PCH₂P(Ph)₂ and S₂CNEt₂; n = 1, 0) and a polymer [W₂(μ-SPh)₂(CO)₆(Ph)₂PCH₂P(Ph)₂]_m, 127

Synthesis and structure of *ansa*-cyclopentadienyl pyrrolyl titanium complexes: [(η⁵-C₅H₄)CH₂(2-C₄H₃N)]Ti(NMe₂)₂ and [1,3-{CH₂(2-C₄H₃N)}₂(η⁵-C₅H₃)]Ti(NMe₂)₂, 79

The initiation of ring-opening metathesis polymerisation of norbornadiene by seven-coordinate molybdenum(II) compounds. X-ray crystal structure of [Mo(μ-Cl)(SnCl₃)(CO)₃(η⁴-NBD)], 72

C/Si/Ge bioisosterism

Syntheses and pharmacological characterization of achiral and chiral enantiopure C/Si/Ge-analogous derivatives of the muscarinic antagonist cycrimine: a study on C/Si/Ge bioisosterism, 140

Cycloaddition

The deviating behavior of thiols in nucleophilic trapping reactions of chromiumcarbonyl phenyl complex substituted propargyl cation, 41

Cyclometalation

Iron carbonyl complexes from 2-[2,3-diaza-4-(2-thienyl)buta-1,3-dienyl]thiophene: N–N bond cleavage and cyclometalation, 85

Cyclopentadienyl

Synthesis and structure of *ansa*-cyclopentadienyl pyrrolyl titanium complexes: [(η⁵-C₅H₄)CH₂(2-C₄H₃N)]Ti(NMe₂)₂ and [1,3-{CH₂(2-C₄H₃N)}₂(η⁵-C₅H₃)]Ti(NMe₂)₂, 79

Derivatised ferrocenes

Synthesis of metallocenes of zirconium, hafnium, manganese, iron, tin, lead and half-sandwich complexes of rhodium and iridium containing the ligands (η-C₅R₄CR₂PMe₂), where R and R' may be H or Me, 93

Derivatised manganocenes

Synthesis of metallocenes of zirconium, hafnium, manganese, iron, tin, lead and half-sandwich complexes of rhodium and iridium containing the ligands (η-C₅R₄CR₂PMe₂), where R and R' may be H or Me, 93

DFT

Density functional studies of the reactivity in the C–F bond activation of fluoromethane by bare lanthanum monocation, 121

Diphosphaferrocene

Bis(diphosphaferrocene) palladium(II) dimer complexes as efficient catalysts in the synthesis of arylboronic esters, 197

Diphosphines

Selectivity for the methoxycarbonylation of ethylene versus CO–ethylene copolymerization with catalysts based on C₄-bridged bidentate phosphines and phospholes, 182

1,3-Disilacyclobutane

UV laser-induced photolysis of 1,3-disilacyclobutane in oxygen for chemical vapour deposition of nano-sized polyoxocarbosilane films, 170

Electrochemical reaction

Preparation and characterization of ruthenium(II), rhodium(III) and iridium(III) complexes of isocyanide bearing the azo group, 10

Electron transfer

Density functional studies of the reactivity in the C–F bond activation of fluoromethane by bare lanthanum monocation, 121

EPR properties

Interaction of half-sandwich alkylmolybdenum(III) complexes with B(C₆F₅)₃. The X-ray structure of [CpMo(η⁴-C₄H₆)(μ-Cl)(μ-CH₂)(O)MoCp][CH₃B(C₆F₅)₃], 113

Ethylene

Selectivity for the methoxycarbonylation of ethylene versus CO–ethylene copolymerization with catalysts based on C₄-bridged bidentate phosphines and phospholes, 182

Fluoromethane

Density functional studies of the reactivity in the C–F bond activation of fluoromethane by bare lanthanum monocation, 121

Force constants

Force constant calculations for octahedral complexes of the type M(CO)₅L, based on the CO-factored force field, 37

Functionalised metallocene compounds

Synthesis of metallocenes of zirconium, hafnium, manganese, iron, tin, lead and half-sandwich complexes of rhodium and iridium containing the ligands (η-C₅R₄CR₂PMe₂), where R and R' may be H or Me, 93

Germanium

Syntheses and pharmacological characterization of achiral and chiral enantiopure C/Si/Ge-analogous derivatives of the muscarinic antagonist cycrimine: a study on C/Si/Ge bioisosterism, 140

Half-sandwich

Interaction of half-sandwich alkylmolybdenum(III) complexes with B(C₆F₅)₃. The X-ray structure of [CpMo(η⁴-C₄H₆)(μ-Cl)(μ-CH₂)(O)MoCp][CH₃B(C₆F₅)₃], 113

Hexachloroiridate

Cation–anion interactions in triphenyl telluronium salts. The crystal structures of (Ph₃Te)₂[MCl₆] (M = Pt, Ir), (Ph₃Te)-[AuCl₄], and (Ph₃Te)(NO₃)·HNO₃, 50

Hexachloroplatinate

Cation–anion interactions in triphenyl telluronium salts. The crystal structures of (Ph₃Te)₂[MCl₆] (M = Pt, Ir), (Ph₃Te)-[AuCl₄], and (Ph₃Te)(NO₃)·HNO₃, 50

Homogeneous Ziegler–Natta catalysts

Alkylaminophosphanyl substituted half-sandwich complexes of vanadium(III) and chromium(III): preparation and reactivity in ethylene polymerisation, 21

Infrared spectra

Force constant calculations for octahedral complexes of the type M(CO)₅L, based on the CO-factored force field, 37

- Iridium**
Preparation and characterization of ruthenium(II), rhodium(III) and iridium(III) complexes of isocyanide bearing the azo group, 10
- Iron carbonyl complexes**
Iron carbonyl complexes from 2-[2,3-diaza-4-(2-thienyl)buta-1,3-dienyl]thiophene: N–N bond cleavage and cyclometalation, 85
- Isocyanazobenzene**
Preparation and characterization of ruthenium(II), rhodium(III) and iridium(III) complexes of isocyanide bearing the azo group, 10
- Isomerization**
The deviating behavior of thiols in nucleophilic trapping reactions of chromiumcarbonyl phenyl complex substituted propargyl cation, 41
- Lanthanum monocation**
Density functional studies of the reactivity in the C–F bond activation of fluoromethane by bare lanthanum monocation, 121
- Laser photolysis**
UV laser-induced photolysis of 1,3-disilacyclobutane in oxygen for chemical vapour deposition of nano-sized polyoxocarbosilane films, 170
- Matrix isolation**
Photolysis of $(\text{Me}_2\text{Si})_6$ in argon matrices doped with high concentrations (ca. 20%) of N_2O or $\text{C}_2\text{H}_4\text{O}$: formation of $(\text{Me}_2\text{SiO})_6$, 177
- Mechanism**
Anchimeric assistance by γ -substituents Z, Z = MeO, PhO, MeS or PhS, in reactions of the bromides $(\text{Me}_3\text{Si})_2(\text{ZMe}_2\text{Si})\text{CSiMe}_2\text{Br}$ with AgBF_4 , 29
- $(\text{Me}_2\text{Si})_6$
Photolysis of $(\text{Me}_2\text{Si})_6$ in argon matrices doped with high concentrations (ca. 20%) of N_2O or $\text{C}_2\text{H}_4\text{O}$: formation of $(\text{Me}_2\text{SiO})_6$, 177
- Metathesis**
The initiation of ring-opening metathesis polymerisation of norbornadiene by seven-coordinate molybdenum(II) compounds. X-ray crystal structure of $[\text{Mo}(\mu\text{-Cl})(\text{SnCl}_3)(\text{CO})_3(\eta^4\text{-NBD})]$, 72
- Methoxycarbonylation**
Selectivity for the methoxycarbonylation of ethylene versus CO–ethylene copolymerization with catalysts based on C_4 -bridged bidentate phosphines and phospholes, 182
- Molybdenum**
Interaction of half-sandwich alkylmolybdenum(III) complexes with $\text{B}(\text{C}_6\text{F}_5)_3$. The X-ray structure of $[\text{CpMo}(\eta^4\text{-C}_4\text{H}_6)(\mu\text{-Cl})(\mu\text{-CH}_2)(\text{O})\text{MoCp}][\text{CH}_3\text{B}(\text{C}_6\text{F}_5)_3]$, 113
The initiation of ring-opening metathesis polymerisation of norbornadiene by seven-coordinate molybdenum(II) compounds. X-ray crystal structure of $[\text{Mo}(\mu\text{-Cl})(\text{SnCl}_3)(\text{CO})_3(\eta^4\text{-NBD})]$, 72
- Mo_2S_2 -unit
New ligand substitution modes of $\text{M}_2(\mu\text{-SR})_2(\text{CO})_8$ (M = Mo, W; R = alkyl, aryl): synthesis, structure and properties of new di-nuclear metal(I) carbonyl thiolates, $[\text{M}_2(\mu\text{-SPh})_2(\text{CO})_x\text{L}_y]^n$ (M = Mo, W; x = 6, 8; y = 0, 1; L = PhS, $(\text{Ph})_2\text{PCH}_2\text{P}(\text{Ph})_2$ and S_2CNET_2 ; n = 1, 0) and a polymer $[\text{W}_2(\mu\text{-SPh})_2(\text{CO})_6(\text{Ph})_2\text{PCH}_2\text{P}(\text{Ph})_2]_m$, 127
- Muscarinic antagonists**
Syntheses and pharmacological characterization of achiral and chiral enantiopure C/Si/Ge-analogous derivatives of the muscarinic antagonist cycrimine: a study on C/Si/Ge bioisosterism, 140
- Nanostructured polyoxocarbosilane**
UV laser-induced photolysis of 1,3-disilacyclobutane in oxygen for chemical vapour deposition of nano-sized polyoxocarbosilane films, 170
- Neodymium**
Komplekzkatalyse: LIX. Die Katalyse der 1,4-*trans*-Polymerisation des Butadiens mit Tris(allyl)neodym(III) $\text{Nd}(\eta^3\text{-C}_3\text{H}_5)_3$ als Einkomponentenkatalysator—Kinetik und Reaktionsmechanismus, 1
- N–N Bond activation**
Iron carbonyl complexes from 2-[2,3-diaza-4-(2-thienyl)buta-1,3-dienyl]thiophene: N–N bond cleavage and cyclometalation, 85
- Norbornadiene**
The initiation of ring-opening metathesis polymerisation of norbornadiene by seven-coordinate molybdenum(II) compounds. X-ray crystal structure of $[\text{Mo}(\mu\text{-Cl})(\text{SnCl}_3)(\text{CO})_3(\eta^4\text{-NBD})]$, 72
- Olefin polymerisation**
Alkylaminophosphanyl substituted half-sandwich complexes of vanadium(III) and chromium(III): preparation and reactivity in ethylene polymerisation, 21
- ORD**
Syntheses and pharmacological characterization of achiral and chiral enantiopure C/Si/Ge-analogous derivatives of the muscarinic antagonist cycrimine: a study on C/Si/Ge bioisosterism, 140
- Organometallic chemistry**
Iron carbonyl complexes from 2-[2,3-diaza-4-(2-thienyl)buta-1,3-dienyl]thiophene: N–N bond cleavage and cyclometalation, 85
- Oxidation**
Photolysis of $(\text{Me}_2\text{Si})_6$ in argon matrices doped with high concentrations (ca. 20%) of N_2O or $\text{C}_2\text{H}_4\text{O}$: formation of $(\text{Me}_2\text{SiO})_6$, 177
UV laser-induced photolysis of 1,3-disilacyclobutane in oxygen for chemical vapour deposition of nano-sized polyoxocarbosilane films, 170
- Palladium**
Bis(diphosphaferrocene) palladium(II) dimer complexes as efficient catalysts in the synthesis of arylboronic esters, 197
- Palladium complexes**
Synthesis of chiral 2-alkyl-8-quinolinyloxazoline ligands: reversal of enantioselectivity in the asymmetric palladium-catalyzed allylic alkylation, 65
- Pentacarbonyls**
Force constant calculations for octahedral complexes of the type $\text{M}(\text{CO})_5\text{L}$, based on the CO-factored force field, 37
- Photochemical reaction**
Preparation and characterization of ruthenium(II), rhodium(III) and iridium(III) complexes of isocyanide bearing the azo group, 10
- Photolysis**
Photolysis of $(\text{Me}_2\text{Si})_6$ in argon matrices doped with high concentrations (ca. 20%) of N_2O or $\text{C}_2\text{H}_4\text{O}$: formation of $(\text{Me}_2\text{SiO})_6$, 177
- Platinum**
A new homobimetallic arenediyl diplatinum(II) unit as building block for macromolecular synthesis. X-ray crystal structure of $[\text{C}_6\text{H}_2\{\text{CH}_2\text{NMe}_2\}_2\text{-1,5-}\{\text{PtCl}(\text{PPh}_3)\}_2\text{-2,4}]$, 166
- Polymerisation**
The initiation of ring-opening metathesis polymerisation of norbornadiene by seven-coordinate molybdenum(II) compounds. X-ray crystal structure of $[\text{Mo}(\mu\text{-Cl})(\text{SnCl}_3)(\text{CO})_3(\eta^4\text{-NBD})]$, 72

Polymerization

Komplekatalyse: LIX. Die Katalyse der 1,4-*trans*-Polymerisation des Butadiens mit Tris(allyl)neodym(III) $\text{Nd}(\eta^3\text{-C}_3\text{H}_5)_3$ als Einkomponentenkatalysator—Kinetik und Reaktionsmechanismus, 1

Pyrrolyl

Synthesis and structure of *ansa*-cyclopentadienyl pyrrolyl titanium complexes: $[(\eta^5\text{-C}_5\text{H}_4\text{CH}_2(2\text{-C}_4\text{H}_3\text{N}))\text{Ti}(\text{NMe}_2)_2$ and $[1,3\text{-}\{\text{CH}_2(2\text{-C}_4\text{H}_3\text{N})\}_2(\eta^5\text{-C}_5\text{H}_3)]\text{Ti}(\text{NMe}_2)_2$, 79

Rhodium

Preparation and characterization of ruthenium(II), rhodium(III) and iridium(III) complexes of isocyanide bearing the azo group, 10

R,R-DIOP

Synthesis and circular dichroism spectra of silver(I) complexes with *R,R*-DIOP (4*R*,5*R*-*trans*-4,5-bis[(diphenylphosphino)methyl]-2,2-dimethyl-1,3-dioxalane): crystal structures of $[\text{AgCl}(\text{R,R-DIOP})_2]_2 \cdot 2\text{CHCl}_3$, $\{[\text{AgBr}(\text{R,R-DIOP})_2]_2 \cdot \text{CH}_2\text{Cl}_2 \cdot 2\text{H}_2\text{O}$, $[\text{AgI}(\text{R,R-DIOP})_2]$ and $[\text{AgSCN}(\text{R,R-DIOP})_2]$, 57

Ruthenium

Preparation and characterization of ruthenium(II), rhodium(III) and iridium(III) complexes of isocyanide bearing the azo group, 10

Selectivity

Selectivity for the methoxycarbonylation of ethylene versus CO-ethylene copolymerization with catalysts based on C_4 -bridged bidentate phosphines and phospholes, 182

Silene

UV laser-induced photolysis of 1,3-disilacyclobutane in oxygen for chemical vapour deposition of nano-sized polyoxocarbosilane films, 170

Silicon

Anchimeric assistance by γ -substituents Z, Z = MeO, PhO, MeS or PhS, in reactions of the bromides $(\text{Me}_3\text{Si})_2(\text{ZMe}_2\text{Si})\text{-CSiMe}_2\text{Br}$ with AgBF_4 , 29

Syntheses and pharmacological characterization of achiral and chiral enantiopure C/Si/Ge-analogous derivatives of the muscarinic antagonist cycrimine: a study on C/Si/Ge bioisosterism, 140

Silver complexes

Synthesis and circular dichroism spectra of silver(I) complexes with *R,R*-DIOP (4*R*,5*R*-*trans*-4,5-bis[(diphenylphosphino)methyl]-2,2-dimethyl-1,3-dioxalane): crystal structures of $[\text{AgCl}(\text{R,R-DIOP})_2]_2 \cdot 2\text{CHCl}_3$, $\{[\text{AgBr}(\text{R,R-DIOP})_2]_2 \cdot \text{CH}_2\text{Cl}_2 \cdot 2\text{H}_2\text{O}$, $[\text{AgI}(\text{R,R-DIOP})_2]$ and $[\text{AgSCN}(\text{R,R-DIOP})_2]$, 57

Spectroscopy and electrochemistry

New ligand substitution modes of $\text{M}_2(\mu\text{-SR})_2(\text{CO})_8$ (M = Mo, W; R = alkyl, aryl): synthesis, structure and properties of new di-nuclear metal(I) carbonyl thiolates, $[\text{M}_2(\mu\text{-SPh})_2(\text{CO})_x\text{L}_y]^{n-}$ (M = Mo, W; x = 6, 8; y = 0, 1; L = PhS, (Ph)₂PCH₂P(Ph)₂ and S₂CNET₂; n = 1, 0) and a polymer $[\text{W}_2(\mu\text{-SPh})_2(\text{CO})_6(\text{Ph})_2\text{PCH}_2\text{P}(\text{Ph})_2]_{nm}$, 127

Structures

Synthesis and circular dichroism spectra of silver(I) complexes with *R,R*-DIOP (4*R*,5*R*-*trans*-4,5-bis[(diphenylphosphino)methyl]-2,2-dimethyl-1,3-dioxalane): crystal structures of $[\text{AgCl}(\text{R,R-DIOP})_2]_2 \cdot 2\text{CHCl}_3$, $\{[\text{AgBr}(\text{R,R-DIOP})_2]_2 \cdot \text{CH}_2\text{Cl}_2 \cdot 2\text{H}_2\text{O}$, $[\text{AgI}(\text{R,R-DIOP})_2]$ and $[\text{AgSCN}(\text{R,R-DIOP})_2]$, 57

Substitution mode

New ligand substitution modes of $\text{M}_2(\mu\text{-SR})_2(\text{CO})_8$ (M = Mo, W; R = alkyl, aryl): synthesis, structure and properties of new di-nuclear metal(I) carbonyl thiolates, $[\text{M}_2(\mu\text{-SPh})_2(\text{CO})_x\text{L}_y]^{n-}$ (M = Mo, W; x = 6, 8; y = 0, 1; L = PhS, (Ph)₂PCH₂P(Ph)₂ and S₂CNET₂; n = 1, 0) and a polymer $[\text{W}_2(\mu\text{-SPh})_2(\text{CO})_6(\text{Ph})_2\text{PCH}_2\text{P}(\text{Ph})_2]_{nm}$, 127

Tetrachloroaurate

Cation-anion interactions in triphenyl telluronium salts. The crystal structures of $(\text{Ph}_3\text{Te})_2[\text{MCl}_6]$ (M = Pt, Ir), $(\text{Ph}_3\text{Te})\text{-}[\text{AuCl}_4]$, and $(\text{Ph}_3\text{Te})(\text{NO}_3)\text{-HNO}_3$, 50

Tetrachloroplatinate

Cation-anion interactions in triphenyl telluronium salts. The crystal structures of $(\text{Ph}_3\text{Te})_2[\text{MCl}_6]$ (M = Pt, Ir), $(\text{Ph}_3\text{Te})\text{-}[\text{AuCl}_4]$, and $(\text{Ph}_3\text{Te})(\text{NO}_3)\text{-HNO}_3$, 50

Tin

The initiation of ring-opening metathesis polymerisation of norbornadiene by seven-coordinate molybdenum(II) compounds. X-ray crystal structure of $[\text{Mo}(\mu\text{-Cl})(\text{SnCl}_3)(\text{CO})_3(\eta^4\text{-NBD})]$, 72

Titanium

Synthesis and structure of *ansa*-cyclopentadienyl pyrrolyl titanium complexes: $[(\eta^5\text{-C}_5\text{H}_4\text{CH}_2(2\text{-C}_4\text{H}_3\text{N}))\text{Ti}(\text{NMe}_2)_2$ and $[1,3\text{-}\{\text{CH}_2(2\text{-C}_4\text{H}_3\text{N})\}_2(\eta^5\text{-C}_5\text{H}_3)]\text{Ti}(\text{NMe}_2)_2$, 79

Triphenyltelluronium

Cation-anion interactions in triphenyl telluronium salts. The crystal structures of $(\text{Ph}_3\text{Te})_2[\text{MCl}_6]$ (M = Pt, Ir), $(\text{Ph}_3\text{Te})\text{-}[\text{AuCl}_4]$, and $(\text{Ph}_3\text{Te})(\text{NO}_3)\text{-HNO}_3$, 50

Trisyl

Anchimeric assistance by γ -substituents Z, Z = MeO, PhO, MeS or PhS, in reactions of the bromides $(\text{Me}_3\text{Si})_2(\text{ZMe}_2\text{Si})\text{-CSiMe}_2\text{Br}$ with AgBF_4 , 29

Vanadium

Alkylaminophosphanyl substituted half-sandwich complexes of vanadium(III) and chromium(III): preparation and reactivity in ethylene polymerisation, 21

 W_2S_2 unit

New ligand substitution modes of $\text{M}_2(\mu\text{-SR})_2(\text{CO})_8$ (M = Mo, W; R = alkyl, aryl): synthesis, structure and properties of new di-nuclear metal(I) carbonyl thiolates, $[\text{M}_2(\mu\text{-SPh})_2(\text{CO})_x\text{L}_y]^{n-}$ (M = Mo, W; x = 6, 8; y = 0, 1; L = PhS, (Ph)₂PCH₂P(Ph)₂ and S₂CNET₂; n = 1, 0) and a polymer $[\text{W}_2(\mu\text{-SPh})_2(\text{CO})_6(\text{Ph})_2\text{PCH}_2\text{P}(\text{Ph})_2]_{nm}$, 127

X-ray structure

Interaction of half-sandwich alkylmolybdenum(III) complexes with $\text{B}(\text{C}_6\text{F}_5)_3$. The X-ray structure of $[\text{CpMo}(\eta^4\text{-C}_4\text{H}_6)(\mu\text{-Cl})(\mu\text{-CH}_2(\text{O})\text{MoCp})][\text{CH}_3\text{B}(\text{C}_6\text{F}_5)_3]$, 113