

## Contents of Volume 640

### Regular Papers

|   |     |
|---|-----|
| Komplekxkatalyse: LIX. Die Katalyse der 1,4- <i>trans</i> -Polymerisation des Butadiens mit Tris(allyl)neodym(III) Nd( $\eta^3$ -C <sub>3</sub> H <sub>5</sub> ) <sub>3</sub> als Einkomponentenkatalysator—Kinetik und Reaktionsmechanismus<br>S. Maiwald, H. Weißenborn, C. Sommer, G. Müller, R. Taube . . . . .   | 1   |
| Preparation and characterization of ruthenium(II), rhodium(III) and iridium(III) complexes of isocyanide bearing the azo group<br>Y. Yamamoto, H. Nakamura, J.-F. Ma . . . . .  | 10  |
| Alkylaminophosphanyl substituted half-sandwich complexes of vanadium(III) and chromium(III): preparation and reactivity in ethylene polymerisation<br>V.V. Kotov, E.V. Avtomonov, J. Sundermeyer, E. Aitola, T. Repo, D.A. Lemenovskii . . . . .  | 21  |
| Anchimeric assistance by $\gamma$ -substituents Z, Z = MeO, PhO, MeS or PhS, in reactions of the bromides (Me <sub>3</sub> Si) <sub>2</sub> (ZMe <sub>2</sub> Si)CSiMe <sub>2</sub> Br with AgBF <sub>4</sub><br>C. Eaborn, A. Kowalewska, J. David Smith, W.A. Stańczyk . . . . .  | 29  |
| Force constant calculations for octahedral complexes of the type M(CO) <sub>5</sub> L, based on the CO-factored force field<br>D. Karakaş, C. Kaya . . . . .  | 37  |
| The deviating behavior of thiols in nucleophilic trapping reactions of chromiumcarbonyl phenyl complex substituted propargyl cation<br>A. Netz, K. Polborn, T.J.J. Müller . . . . .   | 41  |
| Cation–anion interactions in triphenyl telluronium salts. The crystal structures of (Ph <sub>3</sub> Te) <sub>2</sub> [MCl <sub>6</sub> ] (M = Pt, Ir), (Ph <sub>3</sub> Te)[AuCl <sub>4</sub> ], and (Ph <sub>3</sub> Te)(NO <sub>3</sub> )·HNO <sub>3</sub><br>R. Oilunkaniemi, J. Pietikäinen, R.S. Laitinen, M. Ahlgren . . . . .   | 50  |
| Synthesis and circular dichroism spectra of silver(I) complexes with <i>R,R</i> -DIOP (4 <i>R</i> ,5 <i>R</i> - <i>trans</i> -4,5-bis[(diphenylphosphino)methyl]-2,2-dimethyl-1,3-dioxalane): crystal structures of [AgCl( <i>R,R</i> -DIOP)] <sub>2</sub> ·2CHCl <sub>3</sub> , {[AgBr( <i>R,R</i> -DIOP)] <sub>2</sub> } <sub>2</sub> ·CH <sub>2</sub> Cl <sub>2</sub> ·2H <sub>2</sub> O, [AgI( <i>R,R</i> -DIOP)] <sub>2</sub> and [AgSCN( <i>R,R</i> -DIOP)] <sub>2</sub><br>B. Wu, X.-T. Wu, X. Tian, W.-H. Sun . . . . . | 57  |
| Synthesis of chiral 2-alkyl-8-quinolinyl-oxazoline ligands: reversal of enantioselectivity in the asymmetric palladium-catalyzed allylic alkylation<br>X.-G. Li, X. Cheng, J.-A. Ma, Q.-L. Zhou . . . . .   | 65  |
| The initiation of ring-opening metathesis polymerisation of norbornadiene by seven-coordinate molybdenum(II) compounds. X-ray crystal structure of [Mo( $\mu$ -Cl)(SnCl <sub>3</sub> )(CO) <sub>3</sub> ( $\eta^4$ -NBD)]<br>T. Szymańska-Buzar, T. Głowiak, I. Czełuśniak . . . . .  | 72  |
| Synthesis and structure of <i>ansa</i> -cyclopentadienyl pyrrolyl titanium complexes: [( $\eta^5$ -C <sub>5</sub> H <sub>4</sub> )CH <sub>2</sub> (2-C <sub>4</sub> H <sub>3</sub> N)]Ti(NMe <sub>2</sub> ) <sub>2</sub> and [1,3-{CH <sub>2</sub> (2-C <sub>4</sub> H <sub>3</sub> N)} <sub>2</sub> ( $\eta^5$ -C <sub>5</sub> H <sub>3</sub> )]Ti(NMe <sub>2</sub> )<br>W.S. Seo, Y.J. Cho, S.C. Yoon, J.T. Park, Y. Park . . . . .   | 79  |
| Iron carbonyl complexes from 2-[2,3-diaza-4-(2-thienyl)buta-1,3-dienyl]thiophene: N–N bond cleavage and cyclometalation<br>C.-J. Lin, W.-S. Hwang, M.Y. Chiang . . . . .  | 85  |
| Synthesis of metallocenes of zirconium, hafnium, manganese, iron, tin, lead and half-sandwich complexes of rhodium and iridium containing the ligands ( $\eta$ -C <sub>5</sub> R <sub>4</sub> CR <sub>2</sub> PMe <sub>2</sub> ), where R and R' may be H or Me<br>R.M. Bellabarba, G.P. Clancy, P.T. Gomes, A.M. Martins, L.H. Rees, M.L.H. Green . . . . .  | 93  |
| Interaction of half-sandwich alkylmolybdenum(III) complexes with B(C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> . The X-ray structure of [CpMo( $\eta^4$ -C <sub>4</sub> H <sub>6</sub> )( $\mu$ -Cl)( $\mu$ -CH <sub>2</sub> )(O)MoCp][CH <sub>3</sub> B(C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> ]<br>E. Le Grogneç, R. Poli, P. Richard, R. Llusar, S. Uriel . . . . .   | 113 |

|   |     |
|---|-----|
| Density functional studies of the reactivity in the C–F bond activation of fluoromethane by bare lanthanum monocation<br>D. Zhang, C. Zhang, C. Liu . . . . .   | 121 |
| New ligand substitution modes of $M_2(\mu-SR)_2(CO)_8$ ( $M = Mo, W$ ; $R = \text{alkyl, aryl}$ ): synthesis, structure and properties of new di-nuclear metal(I) carbonyl thiolates, $[M_2(\mu-SPh)_2(CO)_xL_y]^n-$ ( $M = Mo, W$ ; $x = 6, 8$ ; $y = 0, 1$ ; $L = PhS, (Ph)_2PCH_2P(Ph)_2$ and $S_2CNET_2$ ; $n = 1, 0$ ) and a polymer $[W_2(\mu-SPh)_2(CO)_6(Ph)_2PCH_2P(Ph)_2]_m$<br>B. Zhuang, H. Sun, G. Pan, L. He, Q. Wei, Z. Zhou, S. Peng, K. Wu . . . . . | 127 |
| Syntheses and pharmacological characterization of achiral and chiral enantiopure C/Si/Ge-analogous derivatives of the muscarinic antagonist cycrimine: a study on C/Si/Ge bioisosterism<br>R. Tacke, T. Kornek, T. Heinrich, C. Burschka, M. Penka, M. Pülm, C. Keim, E. Mutschler, G. Lambrecht . . . . .  | 140 |
| A new homobimetallic arenediyl diplatinum(II) unit as building block for macromolecular synthesis. X-ray crystal structure of $[C_6H_2\{CH_2NMe_2\}_2-1,5-\{PtCl(PPh_3)\}_2-2,4]$<br>M.D. Meijer, A.W. Kleij, M. Lutz, A.L. Spek, G. van Koten . . . . .  | 166 |
| UV laser-induced photolysis of 1,3-disilacyclobutane in oxygen for chemical vapour deposition of nano-sized polyoxocarbosilane films<br>J. Pola, J. Vitek, Z. Bastl, J. Šubrt . . . . .   | 170 |
| Photolysis of $(Me_2Si)_6$ in argon matrices doped with high concentrations (ca. 20%) of $N_2O$ or $C_2H_4O$ : formation of $(Me_2SiO)_6$<br>M.J. Almond, J.P. Cannady, T.A. Darling, J.S. Ogden, R. Walsh . . . . .  | 177 |
| Selectivity for the methoxycarbonylation of ethylene versus CO–ethylene copolymerization with catalysts based on $C_4$ -bridged bidentate phosphines and phospholes<br>S. Doherty, E.G. Robins, J.G. Knight, C.R. Newman, B. Rhodes, P.A. Champkin, W. Clegg . . . . .  | 182 |
| Bis(diphosphaferrocene) palladium(II) dimer complexes as efficient catalysts in the synthesis of arylboronic esters<br>M. Melaimi, F. Mathey, P. Le Floch . . . . .   | 197 |
| Author Index of Volume 640 . . . . .  | 201 |
| Subject Index of Volume 640 . . . . .   | 203 |
| Contents of volume of Volume 640 . . . . .  | 207 |