



ELSEVIER

Journal of Organometallic Chemistry 649 (2002) 303–305

Journal  
of Organo  
metallic  
Chemistry

www.elsevier.com/locate/jorganchem

## Contents of Volume 649

### Regular Papers

Synthesis of functional derivatives of the [3,3'-Co(1,2-C <sub>2</sub> B <sub>9</sub> H <sub>11</sub> ) <sub>2</sub> ] <sup>-</sup> anion I.B. Sivaev, Z.A. Starikova, S. Sjöberg, V.I. Bregadze . . . . .	1
Synthesis and reactivity of transition metal complexes containing halogenated boryl ligands S. Aldridge, R.J. Calder, R.E. Baghurst, M.E. Light, M.B. Hursthouse . . . . .	9
Synthesis and crystal structure of highly soluble ansa-titano- and zirconocene dichloride complexes [Me <sub>2</sub> Si( $\eta^5$ -C <sub>5</sub> H <sub>2</sub> (SiMe <sub>3</sub> ) <sub>2</sub> ) <sub>2</sub> ]MCl <sub>2</sub> (M = Ti, Zr) B. Douziech, R. Choukroun, C. Lorber, B. Donnadieu . . . . .	15
Alkynethiolate ligands in the syntheses of iron carbonyl derivatives. Crystal structure of [ $\eta^5$ -C <sub>5</sub> H <sub>5</sub> ]Fe(CO) <sub>2</sub> (SC≡CSiMe <sub>3</sub> ) E. Delgado, B. Donnadieu, S. García, F. Zamora . . . . .	21
Synthesis of di- and trisilanes with potentially chelating substituents J. Belzner, U. Dehnert, D. Schär, B. Rohde, P. Müller, I. Usón . . . . .	25
The reaction of [H <sub>4</sub> Ru <sub>4</sub> (CO) <sub>12</sub> ] with 1-penten-3-yne: dimerization and trimerization through the triple bonds F.J. Zuno-Cruz, G. Sánchez-Cabrera, M.J. Rosales-Hoz, H. Nöth . . . . .	43
Cyclopentadienyl, indenyl, and fluorenyl complexes of sodium with 15-crown-5 T. Kähler, U. Behrens, S. Neander, F. Olbrich . . . . .	50
Synthesis, characterization, and reactivity of half-sandwich Ru(II) complexes containing phosphine, arsine, stibine, and bismuthine ligands E. Becker, C. Slugovc, E. Rüba, C. Standfest-Hauser, K. Mereiter, R. Schmid, K. Kirchner . . . . .	55
Acetonitrile activation in di-iron $\mu$ -carbyne complexes: synthesis and structure of the cyanomethyl complex [Fe <sub>2</sub> ( $\mu$ -CNMe <sub>2</sub> )( $\mu$ -CO)(CO)(CH <sub>2</sub> CN)(Cp) <sub>2</sub> ] V.G. Albano, L. Busetto, F. Marchetti, M. Monari, V. Zanotti . . . . .	64
C(sp <sup>3</sup> )–H···O and C(sp <sup>2</sup> )–H···O hydrogen bonds in acyclic- and cyclic-organotellurium carboxylates P.C. Srivastava, S. Bajpai, R. Lath, C. Ram, S. Bajpai, R.J. Butcher, M. Zimmer, M. Veith . . . . .	70
Reactivity of Me <sub>3</sub> M (M = Al, Ga, In) with benzylamine to form the dimers, [Me <sub>2</sub> M(H)CH <sub>2</sub> Ph] <sub>2</sub> , and the hexamers, [MeAlNCH <sub>2</sub> Ph] <sub>6</sub> and [MeGaNCH <sub>2</sub> Ph] <sub>6</sub> : molecular structures of <i>trans</i> -[Me <sub>2</sub> M(H)CH <sub>2</sub> Ph] <sub>2</sub> and [MeAlNCH <sub>2</sub> Ph] <sub>6</sub> E.K. Styron, C.H. Lake, D.H. Powell, L.K. Krannich, C.L. Watkins . . . . .	78
Synthesis of ‘three-petal’ carbocyclic systems based on <i>s-cis</i> -diferrocenyltrienes E.I. Klimova, M. Martínez García, T. Klimova, C. Alvarez Toledano, R. Alfredo Toscano, L. Ruiz Ramírez . . . . .	86
Synthesis and characterisation of new platinum ethynyl dimers and polymers with pendant ferrocenyl groups N.J. Long, A.J.P. White, D.J. Williams, M. Younus . . . . .	94
Rydberg spectra of bis( $\eta^6$ -benzene)chromium derivatives: Gas-phase photoabsorption study of bis( $\eta^6$ -1,2,4-trimethylbenzene)chromium and bis( $\eta^6$ -1,2,4,5-tetramethylbenzene)chromium S.Y. Ketkov, G.K. Fukin, L.N. Zakharov, J.C. Green, G. Clancy . . . . .	100
New insights into the reaction of <i>t</i> -butylhydroperoxide with dichloro- and dimethyl(dioxo)molybdenum(VI) M. Groarke, I.S. Gonçalves, W.A. Herrmann, F.E. Kühn . . . . .	108
X-ray crystal structures of novel platinum(II) and palladium(II) complexes of dialkyl phosphonated phosphines G. Guerrero, P.H. Mutin, F. Dahan, A. Vioux . . . . .	113

Attachment to manganese or cobalt of a bulky tri(organosilyl)methyl ligand containing an NMe <sub>2</sub> or an OMe donor group S.S. Al-Juaid, C. Eaborn, S.M. El-Hamruni, P.B. Hitchcock, J.D. Smith, S.E. Sözerli Can . . . . .	121
EXAFS investigation of metal organic synthesis tools W. Hörner, H. Bertagnoli . . . . .	128
Dicationic triple-decker complexes with a bridging boratabenzene ligand A.R. Kudinov, D.A. Loginov, Z.A. Starikova, P.V. Petrovskii . . . . .	136
Complexation of <i>N</i> -heterocyclic silylenes to Pd(0): formation of highly labile homoleptic complexes Pd[Si('BuNCH <sub>2</sub> CH <sub>2</sub> N'Bu)] <sub>4</sub> and Pd[Si('BuNCHCHN'Bu)] <sub>3</sub> W.A. Herrmann, P. Härter, C.W.K. Gstöttmayr, F. Bielert, N. Seebotth, P. Sirsch . . . . .	141
Coordination chemistry of tin: Part III. The crystal structure and thermal behavior of dipotassium dimethyl-tetrafluoro-stannate dihydrate, K <sub>2</sub> [(CH <sub>3</sub> ) <sub>2</sub> SnF <sub>4</sub> ]·2H <sub>2</sub> O I. Abdelhalim Ahmed, G. Kastner, H. Reuter, D. Schultze . . . . .	147
Synthesis and crystal structures of Group 6 metal carbonyl complexes containing S-rich bis(pyrazol-1-yl)methane ligands L.-F. Tang, W.-L. Jia, Z.-H. Wang, J.-T. Wang, H.-G. Wang . . . . .	152
Syntheses and structural characterization of some mono- and di- <i>p</i> -nitrophenyl[3]ferrocenophanes: X-ray structure of [3]ferrocenophane, 3- <i>p</i> -nitrophenyl[3]ferrocenophane, and 3,4'-bis( <i>p</i> -nitrophenyl)[3]ferrocenophane; DFT calculations on ferrocene derivatives O. Kadkin, C. Näther, W. Friedrichsen . . . . .	161
Alkyne oligomerization catalyzed by molybdenum(0) complexes G.A. Ardizzoia, S. Brenna, G. LaMonica, A. Maspero, N. Masciocchi . . . . .	173
Syntheses of the B <sub>(8)</sub> -hydroxy- and B <sub>(8,8')</sub> -dihydroxy-derivatives of the bis(1,2-dicarbollido)-3-cobalt(1)-ate ion by its reductive acetoxylation and hydroxylation: molecular structure of [8,8'-μ-CH <sub>3</sub> C(O) <sub>2</sub> <(1,2-C <sub>2</sub> B <sub>9</sub> H <sub>10</sub> ) <sub>2</sub> -3-Co] <sup>0</sup> zwitterion determined by X-ray diffraction analysis J. Plešek, B. Grüner, J. Báča, J. Fusek, I. Císařová . . . . .	181
Synthese von [(η <sup>5</sup> -C <sub>5</sub> H <sub>5</sub> ) <sub>2</sub> Ti(Cl)(C≡CSiMe <sub>3</sub> )]Ni(CO) und dessen Reaktionsverhalten gegenüber Phosphiten: die Festkörperstruktur von (CO) <sub>2</sub> Ni[P(OC <sub>6</sub> H <sub>4</sub> CH <sub>3</sub> -2) <sub>3</sub> ] <sub>2</sub> E. Meichel, T. Stein, J. Kralik, G. Rheinwald, H. Lang . . . . .	191
Synthesis, characterisation and molecular structure of [Rh(COE) <sub>2</sub> (acac)] (COE = cyclooctene, η <sup>2</sup> -C <sub>8</sub> H <sub>14</sub> ), an important starting material for the preparation of rhodium catalyst precursors J.M. Burke, R. Benjamin Coapes, A.E. Goeta, J.A.K. Howard, T.B. Marder, E.G. Robins, S.A. Westcott . . . . .	199
A tetrameric copper(I) alkoxide with a π-tethered ligand: 2-allyl-6-methylphenoxy copper(I) B. Gustafsson, M. Häkansson, G. Westman, S. Jagner . . . . .	204
Conversion of alkynes to cyclic imides and anhydrides using reactive iron carbonyls prepared from Fe(CO) <sub>5</sub> and Fe <sub>3</sub> (CO) <sub>12</sub> M. Periasamy, C. Rameshkumar, A. Mukkanti . . . . .	209
Synthesis and complexation of a new facultative tridentate S <sub>2</sub> Te donor ligand MeS(CH <sub>2</sub> ) <sub>3</sub> Te(CH <sub>2</sub> ) <sub>3</sub> SMe: crystal structures of [Rh(Cp*)(S <sub>2</sub> Te)][PF <sub>6</sub> ] <sub>2</sub> and [PtCl(S <sub>2</sub> Te)][PF <sub>6</sub> ] <sub>2</sub> M. Hesford, W. Levason, S.D. Orchard, G. Reid . . . . .	214
The first metal complexes of bis(diisopropylamino)carbene: synthesis, structure and ligand properties K. Denk, P. Sirsch, W.A. Herrmann . . . . .	219
Novel titana-2,4-cyclopentadienes by 1,1-alkylboration B. Wrackmeyer, A. Pedall, J. Weidinger . . . . .	225
From bis(silylamino)tin dichlorides via di(1-alkynyl)-bis(silylamino)tin to new heterocycles by 1,1-organoboration B. Wrackmeyer, A. Pedall, W. Milius, O.L. Tok, Y.N. Bubnov . . . . .	232
Silatetrazoline: Eigenschaften und Strukturen cyclischer thermostabiler Tetrazenderivate H.-W. Lerner, M. Bolte, N. Wiberg . . . . .	246
The bent metallocene geometries of potassium polyalkyl cyclopentadienyl THF solvates: monosolvated [(THF)K(μ-C <sub>5</sub> Me <sub>5</sub> )] <sub>n</sub> , disolvated [(THF) <sub>2</sub> K(μ-C <sub>5</sub> Me <sub>5</sub> )] <sub>n</sub> and the tethered olefin complex [(THF)K(μ-C <sub>5</sub> Me <sub>4</sub> SiMe <sub>2</sub> CH <sub>2</sub> CH=CH <sub>2</sub> )] <sub>n</sub> W.J. Evans, J.C. Brady, C.H. Fujimoto, D.G. Giarikos, J.W. Ziller . . . . .	252

Asymmetric addition of diethylzinc to aromatic aldehydes by chiral ferrocene-based catalysts T.-J. Kim, H.-Y. Lee, E.-S. Ryu, D.-K. Park, C.S. Cho, S.C. Shim, J.H. Jeong . . . . .	258
Si–C bond cleavage in the reaction of gallium chloride with lithium bis(trimethylsilyl)amide and thermolysis of base adducts of dichloro(trimethylsilyl)amido gallium compounds B. Luo, V.G. Young Jr., W.L. Gladfelter . . . . .	268
Synthesis of 1,4-disilacyclohexa-2,5-dienes J. Yang, I. Guzei, J.G. Verkade . . . . .	276
<b>Note</b>	
Oxidation of primary and secondary alcohols catalyzed by a pentamethylcyclopentadienyliridium complex K.-i. Fujita, S. Furukawa, R. Yamaguchi . . . . .	289
Book review . . . . .	293
Author Index of Volume 649 . . . . .	295
Subject Index of Volume 649 . . . . .	297
Contents of Volume 649 . . . . .	303