

Subject Index of Volume 649

Acetonitrile

Acetonitrile activation in di-iron μ -carbyne complexes: synthesis and structure of the cyanomethyl complex $[\text{Fe}_2(\mu\text{-CNMe}_2)(\mu\text{-CO})(\text{CO})(\text{CH}_2\text{CN})(\text{Cp})_2]$, 64

Acetylacetonate

Synthesis, characterisation and molecular structure of $[\text{Rh}(\text{COE})_2(\text{acac})]$ (COE = cyclooctene, $\eta^2\text{-C}_8\text{H}_{14}$), an important starting material for the preparation of rhodium catalyst precursors, 199

Acetylene reactions

Synthesis of 1,4-disilacyclohexa-2,5-dienes, 276

Aldehydes

Asymmetric addition of diethylzinc to aromatic aldehydes by chiral ferrocene-based catalysts, 258

Alkene

A tetrameric copper(I) alkoxide with a π -tethered ligand: 2-allyl-6-methylphenoxocopper(I), 204

Alkoxide

A tetrameric copper(I) alkoxide with a π -tethered ligand: 2-allyl-6-methylphenoxocopper(I), 204

Alkylboration

From bis(silylamino)tin dichlorides via di(1-alkynyl)-bis(silylamino)tin to new heterocycles by 1,1-organoboration, 232

Alkylperoxides

New insights into the reaction of *t*-butylhydroperoxide with dichloro- and dimethyl(dioxo)molybdenum(VI), 108

Alkyne

Synthesis and characterisation of new platinum ethynyl dimers and polymers with pendant ferrocenyl groups, 94

Alkynes

Conversion of alkynes to cyclic imides and anhydrides using reactive iron carbonyls prepared from $\text{Fe}(\text{CO})_5$ and $\text{Fe}_3(\text{CO})_{12}$, 209

From bis(silylamino)tin dichlorides via di(1-alkynyl)-bis(silylamino)tin to new heterocycles by 1,1-organoboration, 232

Novel titana-2,4-cyclopentadienes by 1,1-alkylboration, 225

Alkynethiolates derivatives

Alkynethiolate ligands in the syntheses of iron carbonyl derivatives. Crystal structure of $[(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})_2(\text{SC}\equiv\text{CSiMe}_3)]$, 21

Allylboration

From bis(silylamino)tin dichlorides via di(1-alkynyl)-bis(silylamino)tin to new heterocycles by 1,1-organoboration, 232

Allyl complexes

Synthesis, characterization, and reactivity of half-sandwich Ru(II) complexes containing phosphine, arsine, stibine, and bismutine ligands, 55

2-Allyl-6-methylphenol

A tetrameric copper(I) alkoxide with a π -tethered ligand: 2-allyl-6-methylphenoxocopper(I), 204

Amides

From bis(silylamino)tin dichlorides via di(1-alkynyl)-bis(silylamino)tin to new heterocycles by 1,1-organoboration, 232

Novel titana-2,4-cyclopentadienes by 1,1-alkylboration, 225

Amidogallium compounds

Si–C bond cleavage in the reaction of gallium chloride with lithium bis(trimethylsilyl)amide and thermolysis of base adducts of dichloro(trimethylsilyl)amido gallium compounds, 268

ansa-Metallocene

Synthesis and crystal structure of highly soluble ansa-titano- and zirconocene dichloride complexes $[\text{Me}_2\text{Si}(\eta^5\text{-C}_5\text{H}_2(\text{SiMe}_3)_2)_2\text{MCl}_2$ (M = Ti, Zr), 15

Arene coordination

Synthesis, characterization, and reactivity of half-sandwich Ru(II) complexes containing phosphine, arsine, stibine, and bismutine ligands, 55

Arsines

Synthesis, characterization, and reactivity of half-sandwich Ru(II) complexes containing phosphine, arsine, stibine, and bismutine ligands, 55

Asymmetric addition

Asymmetric addition of diethylzinc to aromatic aldehydes by chiral ferrocene-based catalysts, 258

Bent metallocenes

The bent metallocene geometries of potassium polyalkyl cyclopentadienyl THF solvates: monosolvated $[(\text{THF})\text{K}(\mu\text{-C}_5\text{-Me}_5)]_n$, disolvated $[(\text{THF})_2\text{K}(\mu\text{-C}_5\text{Me}_5)]_n$ and the tethered olefin complex $[(\text{THF})\text{K}(\mu\text{-C}_5\text{Me}_4\text{SiMe}_2\text{CH}_2\text{CH}=\text{CH}_2)]_n$, 252

Benzylamine

Reactivity of Me_3M (M = Al, Ga, In) with benzylamine to form the dimers, $[\text{Me}_2\text{MN}(\text{H})\text{CH}_2\text{Ph}]_2$, and the hexamers, $[\text{MeAlNCH}_2\text{Ph}]_6$ and $[\text{MeGaNCH}_2\text{Ph}]_6$: molecular structures of *trans*- $[\text{Me}_2\text{MN}(\text{H})\text{CH}_2\text{Ph}]_2$ and $[\text{MeAlNCH}_2\text{Ph}]_6$, 78

Bisarene complexes

Rydberg spectra of bis(η^6 -benzene)chromium derivatives: Gas-phase photoabsorption study of bis(η^6 -1,2,4-trimethylbenzene)chromium and bis(η^6 -1,2,4,5-tetramethylbenzene)chromium, 100

Bis(1,2-dicarbollido)-3-cobalt(1-)ate ion

Syntheses of the $\text{B}_{(8)}$ -hydroxy- and $\text{B}_{(8,8)}$ -dihydroxy-derivatives of the bis(1,2-dicarbollido)-3-cobalt(1-)ate ion by its reductive acetoxylation and hydroxylation: molecular structure of $[8,8'\text{-}\mu\text{-CH}_3\text{C}(\text{O})_2\text{C} < (1,2\text{-C}_2\text{B}_9\text{H}_{10})_2\text{-3-Co}]^0$ zwitterion determined by X-ray diffraction analysis, 181

Bismutines

Synthesis, characterization, and reactivity of half-sandwich Ru(II) complexes containing phosphine, arsine, stibine, and bismutine ligands, 55

Bis(pyrazol-1-yl)methane

Synthesis and crystal structures of Group 6 metal carbonyl complexes containing S-rich bis(pyrazol-1-yl)methane ligands, 152

Boratabenzene

Dicationic triple-decker complexes with a bridging boratabenzene ligand, 136

- Boron**
 Dicationic triple-decker complexes with a bridging boratabenzene ligand, 136
 Synthesis and reactivity of transition metal complexes containing halogenated boryl ligands, 9
- Boronic acid**
 Synthesis, characterisation and molecular structure of [Rh(COE)₂(acac)] (COE = cyclooctene, η²-C₈H₁₄), an important starting material for the preparation of rhodium catalyst precursors, 199
- Boron neutron capture therapy**
 Synthesis of functional derivatives of the [3,3'-Co(1,2-C₂B₉H₁₁)₂]⁻ anion, 1
- Boryl**
 Synthesis and reactivity of transition metal complexes containing halogenated boryl ligands, 9
- Bridged ferrocenes**
 Syntheses and structural characterization of some mono- and di-*p*-nitrophenyl[3]ferrocenophanes: X-ray structure of [3]ferrocenophane, 3-*p*-nitrophenyl[3]ferrocenophane, and 3,4'-bis-(*p*-nitrophenyl)[3]ferrocenophane; DFT calculations on ferrocene derivatives, 161
- Bulky ligands**
 Attachment to manganese or cobalt of a bulky tri(organosilyl)methyl ligand containing an NMe₂ or an OMe donor group, 121
- Carboraboranes**
 Syntheses of the B₍₈₎-hydroxy- and B_(8,8')-dihydroxy-derivatives of the bis(1,2-dicarbollido)-3-cobalt(1-)-ate ion by its reductive acetoxylation and hydroxylation: molecular structure of [8,8'-μ-CH₃C(O)₂<(1,2-C₂B₉H₁₀)₂-3-Co]⁰ zwitterion determined by X-ray diffraction analysis, 181
- Carbene complexes**
 The first metal complexes of bis(diisopropylamino)carbene: synthesis, structure and ligand properties, 219
- Carbenoids**
 EXAFS investigation of metal organic synthesis tools, 128
- Carbyne complexes**
 Acetonitrile activation in di-iron μ-carbyne complexes: synthesis and structure of the cyanomethyl complex [Fe₂(μ-CNMe₂)(μ-CO)(CO)(CH₂CN)(Cp)₂], 64
- Catalysis**
 Alkyne oligomerization catalyzed by molybdenum(0) complexes, 173
- Catalyst**
 Synthesis, characterisation and molecular structure of [Rh(COE)₂(acac)] (COE = cyclooctene, η²-C₈H₁₄), an important starting material for the preparation of rhodium catalyst precursors, 199
- Chiral ferrocenes**
 Asymmetric addition of diethylzinc to aromatic aldehydes by chiral ferrocene-based catalysts, 258
- Chlorogallium compounds**
 Si-C bond cleavage in the reaction of gallium chloride with lithium bis(trimethylsilyl)amide and thermolysis of base adducts of dichloro(trimethylsilyl)amido gallium compounds, 268
- Chromium**
 Rydberg spectra of bis(η⁶-benzene)chromium derivatives: Gas-phase photoabsorption study of bis(η⁶-1,2,4-trimethylbenzene)chromium and bis(η⁶-1,2,4,5-tetramethylbenzene)chromium, 100
- Cobalt**
 Dicationic triple-decker complexes with a bridging boratabenzene ligand, 136
- Cobalt bis(dicarbollide)**
 Synthesis of functional derivatives of the [3,3'-Co(1,2-C₂B₉H₁₁)₂]⁻ anion, 1
- Copper**
 Synthesis and complexation of a new facultative tridentate S₂Te donor ligand MeS(CH₂)₃Te(CH₂)₃SMe: crystal structures of [Rh(Cp*)(S₂Te)][PF₆]₂ and [PtCl(S₂Te)]PF₆, 214
- Copper(I)**
 A tetrameric copper(I) alkoxide with a π-tethered ligand: 2-allyl-6-methylphenoxocopper(I), 204
- 15-Crown-5**
 Cyclopentadienyl, indenyl, and fluorenyl complexes of sodium with 15-crown-5, 50
- Crystal structure**
 Alkyne oligomerization catalyzed by molybdenum(0) complexes, 173
 A tetrameric copper(I) alkoxide with a π-tethered ligand: 2-allyl-6-methylphenoxocopper(I), 204
- Crystal structures**
 Acetonitrile activation in di-iron μ-carbyne complexes: synthesis and structure of the cyanomethyl complex [Fe₂(μ-CNMe₂)(μ-CO)(CO)(CH₂CN)(Cp)₂], 64
 Coordination chemistry of tin: Part III. The crystal structure and thermal behavior of dipotassium dimethyl-tetrafluoro-stannate dihydrate, K₂[(CH₃)₂SnF₄]₂·2H₂O, 147
 Synthesis of functional derivatives of the [3,3'-Co(1,2-C₂B₉H₁₁)₂]⁻ anion, 1
- Cyclic imides**
 Conversion of alkynes to cyclic imides and anhydrides using reactive iron carbonyls prepared from Fe(CO)₅ and Fe₃(CO)₁₂, 209
- Cyclopentadienide**
 The bent metallocene geometries of potassium polyalkyl cyclopentadienyl THF solvates: monosolvated [(THF)K(μ-C₅-Me₃)_{*n*}], disolvated [(THF)₂K(μ-C₅Me₃)_{*n*}] and the tethered olefin complex [(THF)K(μ-C₅Me₄SiMe₂CH₂CH=CH₂)_{*n*}], 252
- Cyclopentadienyl**
 Cyclopentadienyl, indenyl, and fluorenyl complexes of sodium with 15-crown-5, 50
- Cyclopentadienyl iron complexes**
 Alkynethiolate ligands in the syntheses of iron carbonyl derivatives. Crystal structure of [(η⁵-C₅H₅)Fe(CO)₂(SC=CSiMe₃)], 21
- Cyclotrimerization**
 Alkyne oligomerization catalyzed by molybdenum(0) complexes, 173
- DFT calculations**
 Syntheses and structural characterization of some mono- and di-*p*-nitrophenyl[3]ferrocenophanes: X-ray structure of [3]ferrocenophane, 3-*p*-nitrophenyl[3]ferrocenophane, and 3,4'-bis-(*p*-nitrophenyl)[3]ferrocenophane; DFT calculations on ferrocene derivatives, 161
- Dialkyl phosphonated phosphines**
 X-ray crystal structures of novel platinum(II) and palladium(II) complexes of dialkyl phosphonated phosphines, 113
- Dibenzylamine**
 Reactivity of Me₃M (M = Al, Ga, In) with benzylamine to form the dimers, [Me₂MN(H)CH₂Ph]₂, and the hexamers, [MeAlNCH₂Ph]₆ and [MeGaNCH₂Ph]₆: molecular structures of *trans*-[Me₂MN(H)CH₂Ph]₂ and [MeAlNCH₂Ph]₆, 78
- Diethylzinc**
 Asymmetric addition of diethylzinc to aromatic aldehydes by chiral ferrocene-based catalysts, 258

- Dinuclear
Acetonitrile activation in di-iron μ -carbyne complexes: synthesis and structure of the cyanomethyl complex $[\text{Fe}_2(\mu\text{-CNMe}_2)(\mu\text{-CO})(\text{CO})(\text{CH}_2\text{CN})(\text{Cp})_2]$, 64
- 1,4-Disilacyclohexadiene synthesis
Synthesis of 1,4-disilacyclohexa-2,5-dienes, 276
- Disilane reactions
Synthesis of 1,4-disilacyclohexa-2,5-dienes, 276
- Double carbonylation
Conversion of alkynes to cyclic imides and anhydrides using reactive iron carbonyls prepared from $\text{Fe}(\text{CO})_5$ and $\text{Fe}_3(\text{CO})_{12}$, 209
- DSC
Coordination chemistry of tin: Part III. The crystal structure and thermal behavior of dipotassium dimethyl-tetrafluoro-stannate dihydrate, $\text{K}_2[(\text{CH}_3)_2\text{SnF}_4]\cdot 2\text{H}_2\text{O}$, 147
- Electrochemistry
Synthesis and crystal structure of highly soluble ansa-titano- and zirconocene dichloride complexes $[\text{Me}_2\text{Si}(\eta^5\text{-C}_5\text{H}_2\text{-}(\text{SiMe}_3)_2)_2\text{MCl}_2$ (M = Ti, Zr), 15
- Electronic absorption spectra
Rydberg spectra of bis(η^6 -benzene)chromium derivatives: Gas-phase photoabsorption study of bis(η^6 -1,2,4-trimethylbenzene)chromium and bis(η^6 -1,2,4,5-tetramethylbenzene)chromium, 100
- Enynes
The reaction of $[\text{H}_4\text{Ru}_4(\text{CO})_{12}]$ with 1-penten-3-yne: dimerization and trimerization through the triple bonds, 43
- Epoxidation
New insights into the reaction of *t*-butylhydroperoxide with dichloro- and dimethyl(dioxo)molybdenum(VI), 108
- EXAFS
EXAFS investigation of metal organic synthesis tools, 128
- Ferrocene
Synthesis and characterisation of new platinum ethynyl dimers and polymers with pendant ferrocenyl groups, 94
Synthesis of 'three-petal' carbocyclic systems based on *s-cis*-diferrocenyltrienes, 86
- [3]Ferrocenophane
Syntheses and structural characterization of some mono- and di-*p*-nitrophenyl[3]ferrocenophanes: X-ray structure of [3]ferrocenophane, 3-*p*-nitrophenyl[3]ferrocenophane, and 3,4'-bis(*p*-nitrophenyl)[3]ferrocenophane; DFT calculations on ferrocene derivatives, 161
- Fluorenyl
Cyclopentadienyl, indenyl, and fluorenyl complexes of sodium with 15-crown-5, 50
- Fragmentation
Synthesis of 'three-petal' carbocyclic systems based on *s-cis*-diferrocenyltrienes, 86
- Functional derivatives
Synthesis of functional derivatives of the $[\text{3,3'-Co}(1,2\text{-C}_2\text{B}_9\text{H}_{11})_2]^-$ anion, 1
- Gomberg arylation reaction
Syntheses and structural characterization of some mono- and di-*p*-nitrophenyl[3]ferrocenophanes: X-ray structure of [3]ferrocenophane, 3-*p*-nitrophenyl[3]ferrocenophane, and 3,4'-bis(*p*-nitrophenyl)[3]ferrocenophane; DFT calculations on ferrocene derivatives, 161
- Grignard analogues
Attachment to manganese or cobalt of a bulky tri(organosilyl)methyl ligand containing an NMe_2 or an OMe donor group, 121
- Grignard compounds
EXAFS investigation of metal organic synthesis tools, 128
- Group 6 metal carbonyl complexes
Synthesis and crystal structures of Group 6 metal carbonyl complexes containing S-rich bis(pyrazol-1-yl)methane ligands, 152
- Halosilanes
Synthesis of di- and trisilanes with potentially chelating substituents, 25
- Homoannular alkylation
Synthesis of 'three-petal' carbocyclic systems based on *s-cis*-diferrocenyltrienes, 86
- Hydroboration
Synthesis, characterisation and molecular structure of $[\text{Rh}(\text{COE})_2(\text{acac})]$ (COE = cyclooctene, $\eta^2\text{-C}_8\text{H}_{14}$), an important starting material for the preparation of rhodium catalyst precursors, 199
- Hydrogen bond
Synthesis and reactivity of transition metal complexes containing halogenated boryl ligands, 9
- Hydrogen bonding
X-ray crystal structures of novel platinum(II) and palladium(II) complexes of dialkyl phosphonated phosphines, 113
- Hydrogen transfer
Oxidation of primary and secondary alcohols catalyzed by a pentamethylcyclopentadienyliridium complex, 289
- Hydroxy and dihydroxy derivatives
Syntheses of the $\text{B}_{(8)}$ -hydroxy- and $\text{B}_{(8,8)}$ -dihydroxy-derivatives of the bis(1,2-dicarbollido)-3-cobalt(1-)-ate ion by its reductive acetoxylation and hydroxylation: molecular structure of $[\text{8,8}'\text{-}\mu\text{-CH}_3\text{C}(\text{O})_2\text{-(1,2-C}_2\text{B}_9\text{H}_{10})_2\text{-3-Co}]^0$ zwitterion determined by X-ray diffraction analysis, 181
- Indenyl
Cyclopentadienyl, indenyl, and fluorenyl complexes of sodium with 15-crown-5, 50
- Infrared spectroscopy
A tetrameric copper(I) alkoxide with a π -tethered ligand: 2-allyl-6-methylphenoxocopper(I), 204
- Intermolecular C–H \cdots O hydrogen bonds
C(sp³)–H \cdots O and C(sp²)–H \cdots O hydrogen bonds in acyclic- and cyclic-organotellurium carboxylates, 70
- Iridium
Dicationic triple-decker complexes with a bridging boratabenzene ligand, 136
- Iridium catalyst
Oxidation of primary and secondary alcohols catalyzed by a pentamethylcyclopentadienyliridium complex, 289
- Iridium complexes
The first metal complexes of bis(diisopropylamino)carbene: synthesis, structure and ligand properties, 219
- Iron
Dicationic triple-decker complexes with a bridging boratabenzene ligand, 136
Synthesis and reactivity of transition metal complexes containing halogenated boryl ligands, 9
- Iron carbonyl complexes
Conversion of alkynes to cyclic imides and anhydrides using reactive iron carbonyls prepared from $\text{Fe}(\text{CO})_5$ and $\text{Fe}_3(\text{CO})_{12}$, 209
- Manganese
Synthesis and reactivity of transition metal complexes containing halogenated boryl ligands, 9

Metallacarboranes

Syntheses of the $B_{(8)}$ -hydroxy- and $B_{(8,8)}$ -dihydroxy-derivatives of the bis(1,2-dicarbollido)-3-cobalt(1-)ate ion by its reductive acetoxylation and hydroxylation: molecular structure of $[8,8'\text{-}\mu\text{-CH}_3\text{C}(\text{O})_2\text{<(1,2-C}_2\text{B}_9\text{H}_{10}\text{)}_2\text{-3-Co}]^0$ zwitterion determined by X-ray diffraction analysis, 181

Methyl substituents

Rydberg spectra of bis(η^6 -benzene)chromium derivatives: Gas-phase photoabsorption study of bis(η^6 -1,2,4-trimethylbenzene)chromium and bis(η^6 -1,2,4,5-tetramethylbenzene)chromium, 100

Molybdenum

Alkyne oligomerization catalyzed by molybdenum(0) complexes, 173

New insights into the reaction of *t*-butylhydroperoxide with dichloro- and dimethyl(dioxo)molybdenum(VI), 108

MS

Coordination chemistry of tin: Part III. The crystal structure and thermal behavior of dipotassium dimethyl-tetrafluoro-stannate dihydrate, $K_2[(\text{CH}_3)_2\text{SnF}_4]\cdot 2\text{H}_2\text{O}$, 147

N-Heterocyclic carbenes

The first metal complexes of bis(diisopropylamino)carbene: synthesis, structure and ligand properties, 219

N-heterocyclic silylenes

Complexation of *N*-heterocyclic silylenes to Pd(0): formation of highly labile homoleptic complexes $\text{Pd}[\text{Si}(\text{t-BuNCH}_2\text{-CH}_2\text{N}^t\text{Bu})_4]$ and $\text{Pd}[\text{Si}(\text{t-BuNCHCHN}^t\text{Bu})_3]$, 141

NMR

Coordination chemistry of tin: Part III. The crystal structure and thermal behavior of dipotassium dimethyl-tetrafluoro-stannate dihydrate, $K_2[(\text{CH}_3)_2\text{SnF}_4]\cdot 2\text{H}_2\text{O}$, 147

From bis(silylamino)tin dichlorides via di(1-alkynyl)-bis(silylamino)tin to new heterocycles by 1,1-organoboration, 232

NMR spectroscopy

A tetrameric copper(I) alkoxide with a π -tethered ligand: 2-allyl-6-methylphenoxocopper(I), 204

Nuclear medicine

Synthesis of functional derivatives of the $[3,3'\text{-Co}(1,2\text{-C}_2\text{B}_9\text{H}_{11})_2]^-$ anion, 1

Olefin

Synthesis, characterisation and molecular structure of $[\text{Rh}(\text{COE})_2(\text{acac})]$ (COE = cyclooctene, $\eta^2\text{-C}_8\text{H}_{14}$), an important starting material for the preparation of rhodium catalyst precursors, 199

Oligomerization

The reaction of $[\text{H}_4\text{Ru}_4(\text{CO})_{12}]$ with 1-penten-3-yne: dimerization and trimerization through the triple bonds, 43

Oligosilanes

Synthesis of di- and trisilanes with potentially chelating substituents, 25

1,1-Organoboration

Novel titana-2,4-cyclopentadienes by 1,1-alkylboration, 225

Organocobalt

Attachment to manganese or cobalt of a bulky tri(organosilyl)methyl ligand containing an NMe_2 or an OMe donor group, 121

Organomanganese

Attachment to manganese or cobalt of a bulky tri(organosilyl)methyl ligand containing an NMe_2 or an OMe donor group, 121

Organotellurium carboxylates

$\text{C}(\text{sp}^3)\text{-H}\cdots\text{O}$ and $\text{C}(\text{sp}^2)\text{-H}\cdots\text{O}$ hydrogen bonds in acyclic- and cyclic-organotellurium carboxylates, 70

Oxidation

Oxidation of primary and secondary alcohols catalyzed by a pentamethylcyclopentadienyliridium complex, 289

Palladium(II) complexes

X-ray crystal structures of novel platinum(II) and palladium(II) complexes of dialkyl phosphonated phosphines, 113

Pd complexes

Complexation of *N*-heterocyclic silylenes to Pd(0): formation of highly labile homoleptic complexes $\text{Pd}[\text{Si}(\text{t-BuNCH}_2\text{-CH}_2\text{N}^t\text{Bu})_4]$ and $\text{Pd}[\text{Si}(\text{t-BuNCHCHN}^t\text{Bu})_3]$, 141

Pentamethylcyclopentadienyl complex

Oxidation of primary and secondary alcohols catalyzed by a pentamethylcyclopentadienyliridium complex, 289

Phenoxide

A tetrameric copper(I) alkoxide with a π -tethered ligand: 2-allyl-6-methylphenoxocopper(I), 204

Platinum

Synthesis and characterisation of new platinum ethynyl dimers and polymers with pendant ferrocenyl groups, 94

Synthesis and complexation of a new facultative tridentate S_2Te donor ligand $\text{MeS}(\text{CH}_2)_3\text{Te}(\text{CH}_2)_3\text{SMe}$: crystal structures of $[\text{Rh}(\text{Cp}^*)(\text{S}_2\text{Te})][\text{PF}_6]_2$ and $[\text{PtCl}(\text{S}_2\text{Te})][\text{PF}_6]$, 214

Platinum(II) complexes

X-ray crystal structures of novel platinum(II) and palladium(II) complexes of dialkyl phosphonated phosphines, 113

Polymer

Synthesis and characterisation of new platinum ethynyl dimers and polymers with pendant ferrocenyl groups, 94

Potassium

The bent metallocene geometries of potassium polyalkyl cyclopentadienyl THF solvates: monosolvated $[(\text{THF})\text{K}(\mu\text{-C}_5\text{Me}_5)]_n$, disolvated $[(\text{THF})_2\text{K}(\mu\text{-C}_5\text{Me}_5)]_n$ and the tethered olefin complex $[(\text{THF})\text{K}(\mu\text{-C}_5\text{Me}_4\text{SiMe}_2\text{CH}_2\text{-CH}=\text{CH}_2)]_n$, 252

Powder diffraction

Alkyne oligomerization catalyzed by molybdenum(0) complexes, 173

Reductive acetoxylation

Syntheses of the $B_{(8)}$ -hydroxy- and $B_{(8,8)}$ -dihydroxy-derivatives of the bis(1,2-dicarbollido)-3-cobalt(1-)ate ion by its reductive acetoxylation and hydroxylation: molecular structure of $[8,8'\text{-}\mu\text{-CH}_3\text{C}(\text{O})_2\text{<(1,2-C}_2\text{B}_9\text{H}_{10}\text{)}_2\text{-3-Co}]^0$ zwitterion determined by X-ray diffraction analysis, 181

Reformatsky compounds

EXAFS investigation of metal organic synthesis tools, 128

Rhodium

Dicationic triple-decker complexes with a bridging boratabenzene ligand, 136

Synthesis and complexation of a new facultative tridentate S_2Te donor ligand $\text{MeS}(\text{CH}_2)_3\text{Te}(\text{CH}_2)_3\text{SMe}$: crystal structures of $[\text{Rh}(\text{Cp}^*)(\text{S}_2\text{Te})][\text{PF}_6]_2$ and $[\text{PtCl}(\text{S}_2\text{Te})][\text{PF}_6]$, 214

Synthesis, characterisation and molecular structure of $[\text{Rh}(\text{COE})_2(\text{acac})]$ (COE = cyclooctene, $\eta^2\text{-C}_8\text{H}_{14}$), an important starting material for the preparation of rhodium catalyst precursors, 199

Rhodium complexes

The first metal complexes of bis(diisopropylamino)carbene: synthesis, structure and ligand properties, 219

Ruthenium

Dicationic triple-decker complexes with a bridging boratabenzene ligand, 136

Synthesis, characterization, and reactivity of half-sandwich Ru(II) complexes containing phosphine, arsine, stibine, and bis-mutine ligands, 55

Rydberg spectroscopy

Rydberg spectra of bis(η^6 -benzene)chromium derivatives: Gas-phase photoabsorption study of bis(η^6 -1,2,4-trimethylbenzene)chromium and bis(η^6 -1,2,4,5-tetramethylbenzene)chromium, 100

Sandwich compounds

Dicationic triple-decker complexes with a bridging boratabenzene ligand, 136

s-cis-Diferrocenyltrienes

Synthesis of 'three-petal' carbocyclic systems based on *s-cis*-diferrocenyltrienes, 86

Si–C bond cleavage

Si–C bond cleavage in the reaction of gallium chloride with lithium bis(trimethylsilyl)amide and thermolysis of base adducts of dichloro(trimethylsilyl)amido gallium compounds, 268

Silicon–silicon bond formation

Synthesis of di- and trisilanes with potentially chelating substituents, 25

 μ -Silylene complexes

Complexation of *N*-heterocyclic silylenes to Pd(0): formation of highly labile homoleptic complexes Pd[Si('BuNCH₂-CH₂N'Bu)]₄ and Pd[Si('BuNCHCHN'Bu)]₃, 141

Silylenes

Synthesis of di- and trisilanes with potentially chelating substituents, 25

Sodium

Cyclopentadienyl, indenyl, and fluorenyl complexes of sodium with 15-crown-5, 50

Spirocyclodimers

Synthesis of 'three-petal' carbocyclic systems based on *s-cis*-diferrocenyltrienes, 86

Stibines

Synthesis, characterization, and reactivity of half-sandwich Ru(II) complexes containing phosphine, arsine, stibine, and bisimidine ligands, 55

Structure determination

Synthesis and crystal structure of highly soluble ansa-titano- and zirconocene dichloride complexes [Me₂Si(η^5 -C₅H₂(SiMe₃)₂)MCl₂ (M = Ti, Zr), 15

Substitution

Synthesis and reactivity of transition metal complexes containing halogenated boryl ligands, 9

Supramolecular assemblies

C(sp³)-H...O and C(sp²)-H...O hydrogen bonds in acyclic- and cyclic-organotellurium carboxylates, 70

Telluroether

Synthesis and complexation of a new facultative tridentate S₂Te donor ligand MeS(CH₂)₃Te(CH₂)₃SMe: crystal structures of [Rh(Cp*)(S₂Te)][PF₆]₂ and [PtCl(S₂Te)]PF₆, 214

Tethered olefins

The bent metallocene geometries of potassium polyalkyl cyclopentadienyl THF solvates: monosolvated [(THF)K(μ -C₅-Me₅)]_n, disolvated [(THF)₂K(μ -C₅Me₅)]_n and the tethered olefin complex [(THF)K(μ -C₅Me₄SiMe₂CH₂CH=CH₂)]_n, 252

Tetraruthenium clusters

The reaction of [H₄Ru₄(CO)₁₂] with 1-penten-3-yne: dimerization and trimerization through the triple bonds, 43

Thermolysis

Reactivity of Me₃M (M = Al, Ga, In) with benzylamine to form the dimers, [Me₂MN(H)CH₂Ph]₂, and the hexamers, [MeAlNCH₂Ph]₆ and [MeGaNCH₂Ph]₆: molecular structures of *trans*-[Me₂MN(H)CH₂Ph]₂ and [MeAlNCH₂Ph]₆, 78

Thiocarbene complexes

Acetonitrile activation in di-iron μ -carbyne complexes: synthesis and structure of the cyanomethyl complex [Fe₂(μ -CNMe₂)(μ -CO)(CO)(CH₂CN)(Cp)₂], 64

Thioether

Synthesis and complexation of a new facultative tridentate S₂Te donor ligand MeS(CH₂)₃Te(CH₂)₃SMe: crystal structures of [Rh(Cp*)(S₂Te)][PF₆]₂ and [PtCl(S₂Te)]PF₆, 214

Tin

Coordination chemistry of tin: Part III. The crystal structure and thermal behavior of dipotassium dimethyl-tetrafluoro-stannate dihydrate, K₂[(CH₃)₂SnF₄] \cdot 2H₂O, 147

From bis(silylamino)tin dichlorides via di(1-alkynyl)-bis(silylamino)tin to new heterocycles by 1,1-organoboration, 232

Titanacyclopentadienes

Novel titana-2,4-cyclopentadienes by 1,1-alkylboration, 225

Titanium

Novel titana-2,4-cyclopentadienes by 1,1-alkylboration, 225

Synthesis and crystal structure of highly soluble ansa-titano- and zirconocene dichloride complexes [Me₂Si(η^5 -C₅H₂(SiMe₃)₂)MCl₂ (M = Ti, Zr), 15

Trimethylaluminum

Reactivity of Me₃M (M = Al, Ga, In) with benzylamine to form the dimers, [Me₂MN(H)CH₂Ph]₂, and the hexamers, [MeAlNCH₂Ph]₆ and [MeGaNCH₂Ph]₆: molecular structures of *trans*-[Me₂MN(H)CH₂Ph]₂ and [MeAlNCH₂Ph]₆, 78

Trimethylgallium

Reactivity of Me₃M (M = Al, Ga, In) with benzylamine to form the dimers, [Me₂MN(H)CH₂Ph]₂, and the hexamers, [MeAlNCH₂Ph]₆ and [MeGaNCH₂Ph]₆: molecular structures of *trans*-[Me₂MN(H)CH₂Ph]₂ and [MeAlNCH₂Ph]₆, 78

Trimethylindium

Reactivity of Me₃M (M = Al, Ga, In) with benzylamine to form the dimers, [Me₂MN(H)CH₂Ph]₂, and the hexamers, [MeAlNCH₂Ph]₆ and [MeGaNCH₂Ph]₆: molecular structures of *trans*-[Me₂MN(H)CH₂Ph]₂ and [MeAlNCH₂Ph]₆, 78

Triple bonds

The reaction of [H₄Ru₄(CO)₁₂] with 1-penten-3-yne: dimerization and trimerization through the triple bonds, 43

Triple-decker complexes

Dicationic triple-decker complexes with a bridging boratabenzene ligand, 136

X-ray

From bis(silylamino)tin dichlorides via di(1-alkynyl)-bis(silylamino)tin to new heterocycles by 1,1-organoboration, 232

X-ray crystallography

Syntheses of the B₍₈₎-hydroxy- and B_(8,8')-dihydroxy-derivatives of the bis(1,2-dicarbollido)-3-cobalt(1-ate) ion by its reductive acetoxylation and hydroxylation: molecular structure of [8,8'- μ -CH₃C(O)₂<(1,2-C₂B₉H₁₀)₂-3-Co]⁰ zwitterion determined by X-ray diffraction analysis, 181

X-ray crystal structures

Synthesis and crystal structures of Group 6 metal carbonyl complexes containing S-rich bis(pyrazol-1-yl)methane ligands, 152

X-ray diffraction

Alkynethiolate ligands in the syntheses of iron carbonyl derivatives. Crystal structure of [(η^5 -C₅H₅)Fe(CO)₂-(SC≡CSiMe₃)], 21

X-ray diffraction analysis

Synthesis of 'three-petal' carbocyclic systems based on *s-cis*-diferrocenyltrienes, 86

X-ray structure

Dicationic triple-decker complexes with a bridging boratabenzene ligand, 136

X-ray structures

Complexation of *N*-heterocyclic silylenes to Pd(0): formation of highly labile homoleptic complexes Pd[Si('Bu-NCH₂CH₂N'Bu)]₄ and Pd[Si('BuNCHCHN'Bu)]₃, 141

Si–C bond cleavage in the reaction of gallium chloride with lithium bis(trimethylsilyl)amide and thermolysis of base adducts of dichloro(trimethylsilyl)amido gallium compounds, 268

Zirconium

Synthesis and crystal structure of highly soluble ansa-titano- and zirconocene dichloride complexes [Me₂Si(η⁵-C₅H₂(SiMe₃)₂)₂MC₂] (M = Ti, Zr), 15