



ELSEVIER

Journal of Organometallic Chemistry 658 (2002) 281

Journal
of Organo
metallic
Chemistry

www.elsevier.com/locate/jorganchem

Erratum

Erratum to “Synthesis and reactivity of $\text{Al}(\text{Et})(\text{q}')_2$ ($\text{q}' = 2\text{-methyl-8-quinolinolato}$) and crystal structures of $[\text{Al}(\text{Et})_2\text{q}]_2$ and $\text{Al}(\text{q}')_2\text{q}$ ($\text{q} = 8\text{-quinolinolato}$)”

[Journal of Organometallic Chemistry 654 (2002) 229–232]☆

Isao Yamaguchi, Takayuki Iijima, Takakazu Yamamoto*

Chemical Resources Laboratory, Tokyo Institute of Technology, 4259 Nagatsuta, Midori-ku, 226-8503 Yokohama, Japan

There are errors in the published version of the above-mentioned paper. Synthesis of **2** in experimental part of the published article should read:

Synthesis of 2.

To a toluene solution (50 ml) of AlEt_3 , (4.60 ml 30.0 mmol) was added dropwise a toluene solution (80 ml) of 2-methyl-8-quinolinol (9.55 g, 60.0 mmol) for 1 h. After stirring for 12 h, the resulting precipitate was removed by filtration. The filtrate was dried by evaporation under vacuum to obtain **2** as a yellow solid (10.9 g, 97%). Element Anal. Found: C, 70.81; H, 5.39; N, 7.49%. Calc. for $\text{C}_{22}\text{H}_{21}\text{AlN}_2\text{O}_2$: C, 70.96; H, 5.68; N, 7.52%. $^1\text{H-NMR}$ (400 MHz, CD_2Cl_2): δ 8.23 (d, 2H), 7.45 (d, 2H), 7.38 (t, 2H), 7.13 (dd, 2H), 6.91 (dd, 2H), 3.07 (s, 6H), 0.60 (t, 3H), 0.13 (m, 1H), -0.06 (m, 1H).

☆ PII of original article: S0022-328X(02)01405-5

* Corresponding author. Tel.: +81-45-924-5220; fax: +81-45-924-5276

E-mail address: tyamamot@res.titech.ac.jp (T. Yamamoto).