

## Contents of Volume 657

Editorial	
M. Vaultier . . . . .	1
Recent developments in the chemistry of the nine-vertex monocarbaboranes	
B. Štibr, B. Wrackmeyer . . . . .	3
Imide- and amide-supported Group 5 and 6 metallacarboranes	
A.K. Hughes . . . . .	9
The use of Kitaigorodskii's <i>Aufbau</i> principle in the solid-state study of crystalline borane compounds. A preliminary account	
C. O'Dowd, J.D. Kennedy, M. Thornton-Pett . . . . .	20
Chemistry on a metallathiaborane cluster Part 4: reactions of 11-vertex rhodathiaboranes with bidentate phosphines and their subsequent rearrangements	
O. Volkov, R. Macias, N.P. Rath, L. Barton . . . . .	40
Molecular construction based on icosahedral carboranes and aromatic <i>N,N'</i> -dimethylurea groups. Aromatic layered molecules and a transition metal complex	
Y. Endo, C. Songkram, R. Yamasaki, A. Tanatani, H. Kagechika, K. Takaishi, K. Yamaguchi . . . . .	48
Functionalized cobalt bis(dicarbollide) ions as selective extraction reagents for removal of $M^{2+}$ and $M^{3+}$ cations from nuclear waste, crystal and molecular structures of the $[8,8'-\mu\text{-CIP(O)(O)}_2\text{-(1,2-C}_2\text{B}_9\text{H}_{10})_2\text{-}3,3'\text{-Co}]HN(C_2H_5)_3$ and $[8,8'\text{-}\mu\text{-Et}_2\text{NP(O)(O)}_2\text{-(1,2-C}_2\text{B}_9\text{H}_{10})_2\text{-}3,3'\text{-Co}]HN(CH_3)_3$	
J. Plešek, B. Grüner, I. Císařová, J. Báča, P. Selucky, J. Rais . . . . .	59
Synthesis and structure of 2,4,6-tris[tris(dimethylamino)silyl]borazine: $\{[(CH_3)_2N]_3SiNH\}_3B_3N_3H_3$	
F. Cheng, B. Toury, S.J. Archibald, J.S. Bradley . . . . .	71
High resolution solid state NMR investigation of various boron nitride preceramic polymers	
C. Gervais, F. Babonneau . . . . .	75
Synthesis of $[B_{12}H_{12}]^{2-}$ based extractants and their application for the treatment of nuclear wastes	
R. Bernard, D. Cornu, B. Grüner, J.-F. Dozol, P. Miele, B. Bonnetot . . . . .	83
Pyrolysis of poly[2,4,6-tri(methylamino)borazine] and its conversion into BN fibers	
S. Bernard, D. Cornu, P. Miele, H. Vincent, J. Bouix . . . . .	91
Porous boron nitride supports obtained from molecular precursors. Influence of the precursor formulation and of the thermal treatment on the properties of the BN ceramic	
J.A. Perdigon-Melon, A. Auroux, D. Cornu, P. Miele, B. Toury, B. Bonnetot . . . . .	98
Boron nitride thin fibres obtained from a new copolymer borazine-tri(methylamino)borazine precursor	
C. Duriez, E. Framery, B. Toury, P. Toutois, P. Miele, M. Vaultier, B. Bonnetot . . . . .	107
Synthesis and structure of rhodium complexes with monoanionic carborane ligand $[9\text{-SMe}_2\text{-7,8-C}_2\text{B}_9\text{H}_{10}]^-$	
A.R. Kudinov, D.S. Perekalin, P.V. Petrovskii, K.A. Lyssenko, G.V. Grintselev-Knyazev, Z.A. Starikova . . . . .	115
Allylboration of functionalized isoquinolines	
F.V. Pastukhov, I.V. Yampolsky, Y.N. Bubnov . . . . .	123
Syntheses with organoboranes. XIII. Synthesis of $\omega$ -(4-bromophenyl)alkanoic acids and their borylation	
M. Zaidlewicz, A. Wolan . . . . .	129
Chiral boronates—versatile reagents in asymmetric synthesis	
S. Thormeier, B. Carboni, D.E. Kaufmann . . . . .	136

Reactivity of some poly-1-alkynylsilicon and -tin compounds towards triallylborane—routes to novel heterocycles B. Wrackmeyer, M.H. Bhatti, S. Ali, O.L. Tok, Y.N. Bubnov . . . . .	146
Dihydrogen bonding of decahydro- <i>closو</i> -decaborate(2-) and dodecahydro- <i>closو</i> -dodecaborate(2-) anions with proton donors: experimental and theoretical investigation E.S. Shubina, E.V. Bakhmutova, A.M. Filin, I.B. Sivaev, L.N. Teplitskaya, A.L. Chistyakov, I.V. Stankevich, V.I. Bakhmutov, V.I. Bregadze, L.M. Epstein . . . . .	155
Synthesis and derivatization of the 2-amino- <i>closو</i> -decaborate anion [2-B <sub>10</sub> H <sub>9</sub> NH <sub>3</sub> ] <sup>-</sup> I.B. Sivaev, N.A. Votinova, V.I. Bragin, Z.A. Starikova, L.V. Goeva, V.I. Bregadze, S. Sjöberg . . . . .	163
Polysubstituted derivatives of 3-( $\eta^5$ -Cp)-4-SMe <sub>2</sub> -3,1,2-FeC <sub>2</sub> B <sub>9</sub> H <sub>10</sub> . X-ray structure of 3-( $\eta^5$ -Cp)-4-SMe <sub>2</sub> -7,8-(CF <sub>3</sub> COO) <sub>2</sub> -12-HgCl-3,1,2-FeC <sub>2</sub> B <sub>9</sub> H <sub>7</sub> V.I. Bregadze, I.A. Lobanova, S.V. Timofeev, A.R. Kudinov, V.I. Meshcheryakov, O.L. Tok, P.V. Petrovskii, Z.A. Starikova . . . . .	171
Polyhedral monocarbaborane chemistry: The C,B- <i>para</i> -diphenyl monocarbadodecaborane anion [1,12-Ph <sub>2</sub> - <i>closو</i> -1-CB <sub>11</sub> H <sub>10</sub> ] <sup>-</sup> A. Franken, C.A. Kilner, M. Thornton-Pett, J.D. Kennedy . . . . .	176
Polyhedral monocarbaborane chemistry: The <i>closو</i> -[PhCB <sub>7</sub> H <sub>7</sub> ] <sup>-</sup> and <i>closو</i> -[PhCB <sub>10</sub> H <sub>10</sub> ] <sup>-</sup> anions: the two missing species in the <i>closو</i> -[PhCB <sub>n</sub> H <sub>n</sub> ] <sup>-</sup> sequence A. Franken, C.A. Kilner, M. Thornton-Pett, J.D. Kennedy . . . . .	180
Pd(II) bromide complexes of 1,2-bis(diphenylphosphino)-1,2-dicarba- <i>closو</i> -dodecaborane. Crystal structures of [PdBr <sub>2</sub> (1,2-(PPh <sub>2</sub> ) <sub>2</sub> -1,2-C <sub>2</sub> B <sub>10</sub> H <sub>10</sub> )]·CH <sub>2</sub> Cl <sub>2</sub> , [PdBr <sub>1.133</sub> Cl <sub>0.867</sub> (1,2-(PPh <sub>2</sub> ) <sub>2</sub> -1,2-C <sub>2</sub> B <sub>10</sub> H <sub>10</sub> )]·CH <sub>2</sub> Cl <sub>2</sub> and [PdBrCl <sub>0.541</sub> Me <sub>0.459</sub> (1,2-(PPh <sub>2</sub> ) <sub>2</sub> -1,2-C <sub>2</sub> B <sub>10</sub> H <sub>10</sub> )]·CHCl <sub>3</sub> S. Paavola, F. Teixidor, C. Viñas, R. Kivekäs . . . . .	187
X-ray analysis and structural characterization of 2-phenyl-6-aza-1,3-dioxa-2-borabenzocyclononenones H.I. Beltrán, L.S. Zamudio-Rivera, T. Mancilla, R. Santillan, N. Farfán . . . . .	194
Azanonaborane-pyridine derivatives [(R'C <sub>5</sub> H <sub>4</sub> N)B <sub>8</sub> H <sub>11</sub> NHR']: synthesis, structure and some molecular-orbital calculations C. Bauer, D. Gabel, T. Borrmann, J.D. Kennedy, C.A. Kilner, M. Thornton-Pett, U. Dörfler . . . . .	205
Retention of the B(3)-X (X = Br, I) bond in <i>closو</i> -o-carborane derivatives after nucleophilic attack. The first synthesis of [3-X-7-R-7,8- <i>nido</i> -C <sub>2</sub> B <sub>9</sub> H <sub>10</sub> ] <sup>-</sup> (X = Br, I). Crystal structure of [HNMe <sub>3</sub> ][3-I-7,8- <i>nido</i> -C <sub>2</sub> B <sub>9</sub> H <sub>11</sub> ] G. Barberà, C. Viñas, F. Teixidor, A.J. Welch, G.M. Rosair . . . . .	217
Phosphine–boranes incorporating the carborane cluster R. Núñez, C. Viñas, F. Teixidor, R. Sillanpää, R. Kivekäs . . . . .	224
Modulation of the C–C distance in disubstituted 1,2-R <sub>2</sub> -o-carboranes. Crystal structure of <i>closو</i> 1,2-(SPh) <sub>2</sub> -1,2-C <sub>2</sub> B <sub>10</sub> H <sub>10</sub> J. Llop, C. Viñas, J.M. Oliva, F. Teixidor, M.A. Flores, R. Kivekas, R. Sillanpää . . . . .	232
Cobaltabisdicarbollide anion [Co(C <sub>2</sub> B <sub>9</sub> H <sub>11</sub> ) <sub>2</sub> ] <sup>-</sup> as doping agent on intelligent membranes for ion capture C. Masalles, F. Teixidor, S. Borrós, C. Viñas . . . . .	239
Recent studies on RR'S·C <sub>2</sub> B <sub>9</sub> H <sub>11</sub> charge-compensated ligands: Crystal structures of 10-(S(CH <sub>3</sub> ) <sub>2</sub> )-7,8-C <sub>2</sub> B <sub>9</sub> H <sub>11</sub> and 10-(S(CH <sub>2</sub> ) <sub>4</sub> )-7,8-C <sub>2</sub> B <sub>9</sub> H <sub>11</sub> O. Tutusaus, F. Teixidor, R. Núñez, C. Viñas, R. Sillanpää, R. Kivekäs . . . . .	247
Macropolyhedral boron-containing cluster chemistry: Models for intermediates en route to globular and discoidal megaloborane assemblies. Structures of [ <i>nido</i> -B <sub>10</sub> H <sub>12</sub> ( <i>nido</i> -B <sub>5</sub> H <sub>8</sub> ) <sub>2</sub> ] and [(CH <sub>2</sub> CH <sub>2</sub> C <sub>5</sub> H <sub>4</sub> N)- <i>arachno</i> -B <sub>10</sub> H <sub>10</sub> (NC <sub>5</sub> H <sub>4</sub> - <i>closو</i> -C <sub>2</sub> B <sub>10</sub> H <sub>10</sub> )] as determined by synchrotron X-ray diffraction analysis J. Bould, M.G.S. Londesborough, D.L. Ormsby, J.A. Hugh MacBride, K. Wade, C.A. Kilner, W. Clegg, S.J. Teat, M. Thornton-Pett, R. Greatrex, J.D. Kennedy . . . . .	256
Macropolyhedral boron-containing cluster chemistry: A novel triple-cluster structural motif. Isolation and characterisation of contiguous twenty-vertex [(PPh <sub>3</sub> ) <sub>4</sub> ClPd <sub>4</sub> B <sub>16</sub> H <sub>17</sub> (PPh <sub>3</sub> ) <sub>2</sub> ] M.G.S. Londesborough, C.A. Kilner, M. Thornton-Pett, J.D. Kennedy . . . . .	262
Palladium-catalyzed cross-coupling reactions of arylboronic acids and 2-I- <i>p</i> -carborane L. Eriksson, I.P. Beletskaya, V.I. Bregadze, I.B. Sivaev, S. Sjöberg . . . . .	267
Macropolyhedral thiaboranes with unusual <i>arachno</i> subclusters—a computational investigation M. Hofmann, S.K. Goll . . . . .	273
Erratum . . . . .	279
Author Index of Volume 657 . . . . .	281
Subject Index of Volume 657 . . . . .	283
Contents of Volume 657 . . . . .	289