



ELSEVIER

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Journal of Organometallic Chemistry 663 (2002) 299–303

Journal
of Organo
metallic
Chemistrywww.elsevier.com/locate/jorgchem

Subject Index of Volume 663

- Acetate
Mono-indenyl and cyclopentadienyl derivatives of molybdenum(IV) with a 16 valence-electron configuration, 78
- Acetylacetonate
Mono-indenyl and cyclopentadienyl derivatives of molybdenum(IV) with a 16 valence-electron configuration, 78
- Adaptable coordination modes
Adaptable coordination modes of conjugated 1,3-diene: uniqueness of *s-trans* coordination, 5
- Alkali metals
The activation of C–H bonds in internal alkenes and dienes by {NiCp} species formed in the reaction of nickelocene with alkali metals, 249
- Alkene activation
The activation of C–H bonds in internal alkenes and dienes by {NiCp} species formed in the reaction of nickelocene with alkali metals, 249
- Alkene polymerization
Why propene is not polymerized by (Cp*₂YH)₂: reactions of yttrium alkyl complexes with alkenes produce allyl and vinyl yttrium complexes, 91
- Alkyl complexes
Dialkyl titanium complexes that contain a sulfur-linked bis(phenolato) ligand: The structure of an olefin polymerization catalyst precursor, 158
- Alkyls
Synthesis and structures of crystalline bis(trimethylsilyl)methanido complexes of potassium, calcium and ytterbium, 263
- Allenylidene
Allenylidene-ruthenium-arene precatalyst for ring opening metathesis polymerisation (ROMP), 235
- Allyl complex
Why propene is not polymerized by (Cp*₂YH)₂: reactions of yttrium alkyl complexes with alkenes produce allyl and vinyl yttrium complexes, 91
- Amidine
Mono-indenyl and cyclopentadienyl derivatives of molybdenum(IV) with a 16 valence-electron configuration, 78
- Amino acid
Cyclopalladation of Schiff bases from phenylalanine and 2-phenylglycine, 277
- Anionic
The first anionic arenediethynylgold(I) complexes, 40
- ansa*-Zirconocene
Mechanisms of Sn-to-Zr cyclopentadienyl transfer in the formation of Me₂Si-bridged zirconocenes from sila-stanna-tetrahydro-s-indacenes, 58
- Arduengo carbenes
Structural characterization of Group 4 transition metal halide bis-Arduengo carbene complexes MCl₄L₂: X-ray crystal structure analyses and DFT calculations, 192
- Arenediethynyl
The first anionic arenediethynylgold(I) complexes, 40
- (η⁶-Arene)-ruthenium(II) complexes
Neutral and cationic (η⁶-arene)-ruthenium(II) complexes containing the iminophosphorane–phosphine ligand Ph₂PCH₂P(=N-*p*-C₅F₄N)Ph₂: influence of the arene ring in catalytic transfer hydrogenation of cyclohexanone, 32
- Arsine
Ruthenium hydrides bearing SbPh₃ and AsPh₃ ligands: characterization of the bis(dihydrogen) complexes [Cp*⁺Ru(H₂)₂(EPh₃)]⁺ (Cp* = C₅Me₅; E = Sb, As), 151
- Aryl transfer
Four- five- and six-co-ordinated pentafluorophenylthallium(III) complexes. Crystal structures of [Tl(C₆F₅)₃(OPPh₃)], [Tl(C₆F₅)₂Cl(OPPh₃)₂] and [Tl(C₆F₅)₂(OdppmO)₂][Tl(C₆F₅)₂Cl₂], 289
- Axial chirality
Synthesis and X-ray structures of (biphenoxide)Ti species: four and five coordinate complexes that crystallize as mixtures of diastereomers, 70
- Bent titanocenes
Synthesis and crystal structures of thermally stable titanocenes, 134
- B-H-M agostic bond
Synthesis and characterisation of the *exo-nido* molybdacborane complex Mo(η-C₃H₅)(CO)₂(7,8-μ-SCH₂CH₂S-7,8-*nido*-C₂B₉H₁₀). Strong B–H–Mo 3-centre bonding, 221
- Biphenol
Synthesis and X-ray structures of (biphenoxide)Ti species: four and five coordinate complexes that crystallize as mixtures of diastereomers, 70
- Borneol
Synthesis and X-ray structures of (biphenoxide)Ti species: four and five coordinate complexes that crystallize as mixtures of diastereomers, 70
- Boronic acids
Cross-coupling reactions with boronic acids in water catalysed by oxime-derived palladacycles, 46
- Calcium
Synthesis and structures of crystalline bis(trimethylsilyl)methanido complexes of potassium, calcium and ytterbium, 263
- Carbon–carbon bond formation
Proton induced coupling reactions in dinuclear σ-alkynyl-μ-methylene-rhodium complexes, 145
- Carbonyl substitution reactions
Carbonyl substitution chemistry of neutral and cationic triruthenium cluster complexes derived from 1,1-dimethylhydrazine. Incorporation of two and three phosphane ligands, 227
- Catalytic transfer hydrogenation
Neutral and cationic (η⁶-arene)-ruthenium(II) complexes containing the iminophosphorane–phosphine ligand Ph₂PCH₂P(=N-*p*-C₅F₄N)Ph₂: influence of the arene ring in catalytic transfer hydrogenation of cyclohexanone, 32
- C–H bond activation
The activation of C–H bonds in internal alkenes and dienes by {NiCp} species formed in the reaction of nickelocene with alkali metals, 249

Chelate complexes

Synthesis, structure and catalytic activity of new iminophenolato complexes of scandium and yttrium, 63

Chiral ligand

New ruthenium(II) complexes containing the chiral ligand (4*S*)-2-[(*S*_p)-2-(diphenylphosphino)ferrocenyl]-4-(methylethyl)oxazoline (FcPN). X-ray structures of *mer-trans*-[RuCl₂(dppm)-(FcPN)] (dppm = bis(diphenylphosphino)methane) and *cis*-[RuCl₂(CO)(py)(FcPN)], 204

Clusters

The activation of C–H bonds in internal alkenes and dienes by {NiCp} species formed in the reaction of nickelocene with alkali metals, 249

Conjugated 1,3-diene

Adaptable coordination modes of conjugated 1,3-diene: uniqueness of *s-trans* coordination, 5

Constrained-geometry

New optically active 'constrained-geometry' cyclopentadienyl-phosphine ligands and their metal complexes, 183

Controlled radical polymerization

Half-sandwich molybdenum(III) compounds containing diazadiene ligands and their use in the controlled radical polymerization of styrene, 269

Cp**Ru*Cl(PPh₃)₂

Reaction of Cp**Ru*Cl(PPh₃)₂ with dioxygen and formation of a neutral complex Cp**Ru*Cl(O₂)(PPh₃), 127

Cross-coupling reactions

Cross-coupling reactions with boronic acids in water catalysed by oxime-derived palladacycles, 46

Crystal and molecular structures

C–Br versus C–H bond activation in palladium(II) cyclopalladated compounds. Crystal and molecular structure of [Pd{C₆H₄C(H)=NCy}(MeCOCHCOMe)], 239

Crystal structures

Carbonyl substitution chemistry of neutral and cationic triruthenium cluster complexes derived from 1,1-dimethylhydrazine. Incorporation of two and three phosphane ligands, 227

Four- five- and six-co-ordinated pentafluorophenylthallium(III) complexes. Crystal structures of [Tl(C₆F₅)₃(OPPh₃)], [Tl(C₆F₅)₂Cl(OPPh₃)₂] and [Tl(C₆F₅)₂(OdppmO)₂][Tl(C₆F₅)₂Cl₂], 289

Synthesis of Group 10 polyfluorothiolate mono- and bi-nuclear complexes. Crystal structures of [Ni(SC₆HF₄)₂(dpppe)], [(dpppe)Ni(μ-SC₆HF₄)₂Pd(C₆F₅)₂] and [(dpppe)Ni(μ-SC₆F₅)₂Pd(C₆F₅)₂], 98

Cyclic olefins

Allenylidene-ruthenium-arene precatalyst for ring opening metathesis polymerisation (ROMP), 235

Cyclometallation

Cyclopalladation of Schiff bases from phenylalanine and 2-phenylglycine, 277

Cyclopentadienyl

Half-sandwich molybdenum(III) compounds containing diazadiene ligands and their use in the controlled radical polymerization of styrene, 269

Mono-indenyl and cyclopentadienyl derivatives of molybdenum(IV) with a 16 valence-electron configuration, 78

The activation of C–H bonds in internal alkenes and dienes by {NiCp} species formed in the reaction of nickelocene with alkali metals, 249

Cyclopentadienyl-amido

Hydrazonide titanium derivatives: Synthesis, characterization and catalytic activity in olefin polymerization. Molecular structure of [Ti(η⁵-C₅H₄SiMe₃)Cl(μ-N₂CPh₂)₂], 173

Cyclopentadienyl-phosphine ligands

New optically active 'constrained-geometry' cyclopentadienyl-phosphine ligands and their metal complexes, 183

Deprotonation

Synthesis and characterisation of the *exo-nido* molybdacarborane complex Mo(η-C₃H₅)(CO)₂(7,8-μ-SCH₂CH₂S-7,8-*nido*-C₂B₉H₁₀). Strong B–H–Mo 3-centre bonding, 221

DFT calculation

Structural characterization of Group 4 transition metal halide bis-Arduengo carbene complexes MCl₄L₂: X-ray crystal structure analyses and DFT calculations, 192

DFT calculations

Theoretical analysis of bis(ethylene) complexes of molybdenum and tungsten, 83

Diastereomeric structures

Synthesis and X-ray structures of (biphenoxide)Ti species: four and five coordinate complexes that crystallize as mixtures of diastereomers, 70

Diazadiene ligands

Half-sandwich molybdenum(III) compounds containing diazadiene ligands and their use in the controlled radical polymerization of styrene, 269

Dihydrogen complexes

Ruthenium hydrides bearing SbPh₃ and AsPh₃ ligands: characterization of the bis(dihydrogen) complexes [Cp**Ru*(H₂)₂(EPh₃)]⁺ (Cp* = C₅Me₅; E = Sb, As), 151

Dimethylcyclopentenes

Zirconocene mediated cyclization and isomerization of 1,3,6-heptatriene, 13

1,1-Dimethylhydrazine

Carbonyl substitution chemistry of neutral and cationic triruthenium cluster complexes derived from 1,1-dimethylhydrazine. Incorporation of two and three phosphane ligands, 227

Dioxygen

Reaction of Cp**Ru*Cl(PPh₃)₂ with dioxygen and formation of a neutral complex Cp**Ru*Cl(O₂)(PPh₃), 127

Dithiocarbamate

Mono-indenyl and cyclopentadienyl derivatives of molybdenum(IV) with a 16 valence-electron configuration, 78

Dithiophosphate

Mono-indenyl and cyclopentadienyl derivatives of molybdenum(IV) with a 16 valence-electron configuration, 78

d⁰-Metal complexes

Structural characterization of Group 4 transition metal halide bis-Arduengo carbene complexes MCl₄L₂: X-ray crystal structure analyses and DFT calculations, 192

Ethene polymerization

Structural characterization of Group 4 transition metal halide bis-Arduengo carbene complexes MCl₄L₂: X-ray crystal structure analyses and DFT calculations, 192

Ethylene complexes

Theoretical analysis of bis(ethylene) complexes of molybdenum and tungsten, 83

Ferrocenyloxazoline

New ruthenium(II) complexes containing the chiral ligand (4*S*)-2-[(*S*_p)-2-(diphenylphosphino)ferrocenyl]-4-(methylethyl)oxazoline (FcPN). X-ray structures of *mer-trans*-[RuCl₂(dppm)-(FcPN)] (dppm = bis(diphenylphosphino)methane) and *cis*-[RuCl₂(CO)(py)(FcPN)], 204

Ferrocenylphosphine

New ruthenium(II) complexes containing the chiral ligand (4*S*)-2-[(*S*_p)-2-(diphenylphosphino)ferrocenyl]-4-(methylethyl)oxazoline (FcPN). X-ray structures of *mer-trans*-[RuCl₂(dppm)-(FcPN)] (dppm = bis(diphenylphosphino)methane) and *cis*-[RuCl₂(CO)(py)(FcPN)], 204

Fluxionality

Structural and dynamic studies in solution of anionic dinuclear azolato-bridged palladium(II) complexes, 108

Gold(I)

The first anionic arenediethynylgold(I) complexes, 40

Group 10

Synthesis of Group 10 polyfluorothiolate mono- and bi-nuclear complexes. Crystal structures of $[\text{Ni}(\text{SC}_6\text{HF}_4)_2(\text{dppe})]$, $[(\text{dppe})\text{Ni}(\mu\text{-SC}_6\text{HF}_4)_2\text{Pd}(\text{C}_6\text{F}_5)_2]$ and $[(\text{dppe})\text{Ni}(\mu\text{-SC}_6\text{F}_5)_2\text{Pd}(\text{C}_6\text{F}_5)_2]$, 98

Hemilabile ligands

Neutral and cationic (η^6 -arene)-ruthenium(II) complexes containing the iminophosphorane–phosphine ligand $\text{Ph}_2\text{PCH}_2\text{P}(=\text{N}-p\text{-C}_5\text{F}_4\text{N})\text{Ph}_2$: influence of the arene ring in catalytic transfer hydrogenation of cyclohexanone, 32

1,3,6-Heptatriene

Zirconocene mediated cyclization and isomerization of 1,3,6-heptatriene, 13

Homo- and heterobinuclear

Synthesis of Group 10 polyfluorothiolate mono- and bi-nuclear complexes. Crystal structures of $[\text{Ni}(\text{SC}_6\text{HF}_4)_2(\text{dppe})]$, $[(\text{dppe})\text{Ni}(\mu\text{-SC}_6\text{HF}_4)_2\text{Pd}(\text{C}_6\text{F}_5)_2]$ and $[(\text{dppe})\text{Ni}(\mu\text{-SC}_6\text{F}_5)_2\text{Pd}(\text{C}_6\text{F}_5)_2]$, 98

Hydrazonide

Hydrazonide titanium derivatives: Synthesis, characterization and catalytic activity in olefin polymerization. Molecular structure of $[\text{Ti}(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)\text{Cl}(\mu\text{-N}_2\text{CPh}_2)]_2$, 173

Hydride complexes

Ruthenium hydrides bearing SbPh_3 and AsPh_3 ligands: characterization of the bis(dihydrogen) complexes $[\text{Cp}^*\text{Ru}(\text{H}_2)_2(\text{EPh}_3)]^+$ ($\text{Cp}^* = \text{C}_5\text{Me}_5$; $\text{E} = \text{Sb}, \text{As}$), 151

Hydrides

A study of the coordination ability of 2,5-di(2-pyridyl)phospholes on Ru centres, 118

Imines

Cyclopalladation of Schiff bases from phenylalanine and 2-phenylglycine, 277

Iminophosphoranes

Neutral and cationic (η^6 -arene)-ruthenium(II) complexes containing the iminophosphorane–phosphine ligand $\text{Ph}_2\text{PCH}_2\text{P}(=\text{N}-p\text{-C}_5\text{F}_4\text{N})\text{Ph}_2$: influence of the arene ring in catalytic transfer hydrogenation of cyclohexanone, 32

Indazol

Structural and dynamic studies in solution of anionic dinuclear azolato-bridged palladium(II) complexes, 108

Indenyl

Mono-indenyl and cyclopentadienyl derivatives of molybdenum(IV) with a 16 valence-electron configuration, 78

 π – π Interactions

Synthesis of Group 10 polyfluorothiolate mono- and bi-nuclear complexes. Crystal structures of $[\text{Ni}(\text{SC}_6\text{HF}_4)_2(\text{dppe})]$, $[(\text{dppe})\text{Ni}(\mu\text{-SC}_6\text{HF}_4)_2\text{Pd}(\text{C}_6\text{F}_5)_2]$ and $[(\text{dppe})\text{Ni}(\mu\text{-SC}_6\text{F}_5)_2\text{Pd}(\text{C}_6\text{F}_5)_2]$, 98

Lithiation

Lithiophenylalkyllithiums: new dilithium reagents having both sp^2 - and sp^3 -hybridised remote carbanionic centres, 21

Lithium compounds

Lithiophenylalkyllithiums: new dilithium reagents having both sp^2 - and sp^3 -hybridised remote carbanionic centres, 21

Luminescent properties

A luminescent Pt_3Pb cluster with three $\text{Pt}(\text{II}) \rightarrow \text{Pb}(\text{II})$ bonds, 284

MALDI-TOF-MS

Mass spectrometry of polystyrene and polypropene ruthenium complexes. A new tool for polymer characterization, 213

Menthol

Synthesis and X-ray structures of (biphenoxide)Ti species: four and five coordinate complexes that crystallize as mixtures of diastereomers, 70

Metallate(II) anions

Synthesis and structures of crystalline bis(trimethylsilyl)methanido complexes of potassium, calcium and ytterbium, 263

 μ -Methylene ligand

Proton induced coupling reactions in dinuclear σ -alkynyl- μ -methylene-rhodium complexes, 145

Molybdacarborane

Synthesis and characterisation of the *exo-nido* molybdacarborane complex $\text{Mo}(\eta\text{-C}_3\text{H}_5)(\text{CO})_2(7,8\text{-}\mu\text{-SCH}_2\text{CH}_2\text{S-7,8-nido-C}_2\text{B}_9\text{H}_{10})$. Strong B–H–Mo 3-centre bonding, 221

Molybdenum

Half-sandwich molybdenum(III) compounds containing diazadiene ligands and their use in the controlled radical polymerization of styrene, 269

Mono-indenyl and cyclopentadienyl derivatives of molybdenum(IV) with a 16 valence-electron configuration, 78

Theoretical analysis of bis(ethylene) complexes of molybdenum and tungsten, 83

Nickel

The activation of C–H bonds in internal alkenes and dienes by $\{\text{NiCp}\}$ species formed in the reaction of nickelocene with alkali metals, 249

nido-carboranylthioethers

Synthesis and characterisation of the *exo-nido* molybdacarborane complex $\text{Mo}(\eta\text{-C}_3\text{H}_5)(\text{CO})_2(7,8\text{-}\mu\text{-SCH}_2\text{CH}_2\text{S-7,8-nido-C}_2\text{B}_9\text{H}_{10})$. Strong B–H–Mo 3-centre bonding, 221

Olefin polymerization

Hydrazonide titanium derivatives: Synthesis, characterization and catalytic activity in olefin polymerization. Molecular structure of $[\text{Ti}(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)\text{Cl}(\mu\text{-N}_2\text{CPh}_2)]_2$, 173

Organolithium

Lithiophenylalkyllithiums: new dilithium reagents having both sp^2 - and sp^3 -hybridised remote carbanionic centres, 21

Organometallic compounds

Synthesis of Group 10 polyfluorothiolate mono- and bi-nuclear complexes. Crystal structures of $[\text{Ni}(\text{SC}_6\text{HF}_4)_2(\text{dppe})]$, $[(\text{dppe})\text{Ni}(\mu\text{-SC}_6\text{HF}_4)_2\text{Pd}(\text{C}_6\text{F}_5)_2]$ and $[(\text{dppe})\text{Ni}(\mu\text{-SC}_6\text{F}_5)_2\text{Pd}(\text{C}_6\text{F}_5)_2]$, 98

Oxime derived palladacycles

Cross-coupling reactions with boronic acids in water catalysed by oxime-derived palladacycles, 46

Palladium

Cyclopalladation of Schiff bases from phenylalanine and 2-phenylglycine, 277

Structural and dynamic studies in solution of anionic dinuclear azolato-bridged palladium(II) complexes, 108

Palladium complexes

Mono- and di-nuclear complexes of *ortho*-palladated and -platinated 4,4'-dimethylazobenzene with bis(diphenylphosphino)methane. More data on *transphobia*, 164

Palladium(II) cyclopalladated compounds

C–Br versus C–H bond activation in palladium(II) cyclopalladated compounds. Crystal and molecular structure of $[\text{Pd}\{\text{C}_6\text{H}_4\text{C}(\text{H})=\text{NCy}\}(\text{MeCOCHCOMe})]$, 239

Paramagnetic organometallics

Half-sandwich molybdenum(III) compounds containing diazadiene ligands and their use in the controlled radical polymerization of styrene, 269

Pentafluorophenyl complexes

Four- five- and six-co-ordinated pentafluorophenylthallium(III)

- complexes. Crystal structures of $[\text{Ti}(\text{C}_6\text{F}_5)_3(\text{OPPh}_3)]$, $[\text{Ti}(\text{C}_6\text{F}_5)_2\text{Cl}(\text{OPPh}_3)_2]$ and $[\text{Ti}(\text{C}_6\text{F}_5)_2(\text{OdppmO})_2][\text{Ti}(\text{C}_6\text{F}_5)_2\text{Cl}_2]$, 289
- Pentafluorophenyl derivatives**
A luminescent Pt_3Pb cluster with three $\text{Pt}(\text{II}) \rightarrow \text{Pb}(\text{II})$ bonds, 284
- Phenolato ligands**
Dialkyl titanium complexes that contain a sulfur-linked bis(phenolato) ligand: The structure of an olefin polymerization catalyst precursor, 158
- Phosphine oxide**
Four- five- and six-co-ordinated pentafluorophenylthallium(III) complexes. Crystal structures of $[\text{Ti}(\text{C}_6\text{F}_5)_3(\text{OPPh}_3)]$, $[\text{Ti}(\text{C}_6\text{F}_5)_2\text{Cl}(\text{OPPh}_3)_2]$ and $[\text{Ti}(\text{C}_6\text{F}_5)_2(\text{OdppmO})_2][\text{Ti}(\text{C}_6\text{F}_5)_2\text{Cl}_2]$, 289
- Phospholes**
A study of the coordination ability of 2,5-di(2-pyridyl)phospholes on Ru centres, 118
- Platinum complexes**
A luminescent Pt_3Pb cluster with three $\text{Pt}(\text{II}) \rightarrow \text{Pb}(\text{II})$ bonds, 284
Mono- and di-nuclear complexes of *ortho*-palladated and -platinated 4,4'-dimethylazobenzene with bis(diphenylphosphino)methane. More data on *transphobia*, 164
- Polymerisation**
Allenylidene-ruthenium-arene precatalyst for ring opening metathesis polymerisation (ROMP), 235
- Polymerization catalysts**
Dialkyl titanium complexes that contain a sulfur-linked bis(phenolato) ligand: The structure of an olefin polymerization catalyst precursor, 158
- Polymer mass distribution**
Mass spectrometry of polystyrene and polypropene ruthenium complexes. A new tool for polymer characterization, 213
- Polymer modification**
Mass spectrometry of polystyrene and polypropene ruthenium complexes. A new tool for polymer characterization, 213
- Potassium**
Synthesis and structures of crystalline bis(trimethylsilyl)methanido complexes of potassium, calcium and ytterbium, 263
- Pt \rightarrow Pb bonds**
A luminescent Pt_3Pb cluster with three $\text{Pt}(\text{II}) \rightarrow \text{Pb}(\text{II})$ bonds, 284
- Pyrazol**
Structural and dynamic studies in solution of anionic dinuclear azolato-bridged palladium(II) complexes, 108
- Rhodium**
New optically active 'constrained-geometry' cyclopentadienylphosphine ligands and their metal complexes, 183
- Rhodium complexes**
Proton induced coupling reactions in dinuclear σ -alkynyl- μ -methylene-rhodium complexes, 145
- Ring-opening polymerisation**
Synthesis, structure and catalytic activity of new iminophenolato complexes of scandium and yttrium, 63
- ROMP**
Allenylidene-ruthenium-arene precatalyst for ring opening metathesis polymerisation (ROMP), 235
- Ru organometallics**
Mass spectrometry of polystyrene and polypropene ruthenium complexes. A new tool for polymer characterization, 213
- Ruthenium**
Carbonyl substitution chemistry of neutral and cationic triruthenium cluster complexes derived from 1,1-dimethylhydrazine. Incorporation of two and three phosphane ligands, 227
Ruthenium hydrides bearing SbPh_3 and AsPh_3 ligands: characterization of the bis(dihydrogen) complexes $[\text{Cp}^*\text{Ru}(\text{H}_2)_2(\text{EPh}_3)]^+$ ($\text{Cp}^* = \text{C}_5\text{Me}_5$; $\text{E} = \text{Sb}, \text{As}$), 151
- Ruthenium catalyst**
Allenylidene-ruthenium-arene precatalyst for ring opening metathesis polymerisation (ROMP), 235
- Ruthenium clusters**
Carbonyl substitution chemistry of neutral and cationic triruthenium cluster complexes derived from 1,1-dimethylhydrazine. Incorporation of two and three phosphane ligands, 227
- Ruthenium complexes**
A study of the coordination ability of 2,5-di(2-pyridyl)phospholes on Ru centres, 118
- Ruthenium(II)**
New ruthenium(II) complexes containing the chiral ligand (4*S*)-2-[(*S*_p)-2-(diphenylphosphino)ferrocenyl]-4-(methylethyl)oxazoline (FcPN). X-ray structures of *mer-trans*- $[\text{RuCl}_2(\text{dppm})(\text{FcPN})]$ (dppm = bis(diphenylphosphino)methane) and *cis*- $[\text{RuCl}_2(\text{CO})(\text{py})(\text{FcPN})]$, 204
- Ruthenium neutral complexes**
Reaction of $\text{Cp}^*\text{RuCl}(\text{PPh}_3)_2$ with dioxygen and formation of a neutral complex $\text{Cp}^*\text{RuCl}(\text{O}_2)(\text{PPh}_3)$, 127
- Scandium**
Synthesis, structure and catalytic activity of new iminophenolato complexes of scandium and yttrium, 63
- Siloxanes**
Preparation of Ti(IV), Zr(IV) and Sn(IV) metal alkyls containing the (Pr^iO)₃SiCH₂ fragment, 256
- Silver**
Four- five- and six-co-ordinated pentafluorophenylthallium(III) complexes. Crystal structures of $[\text{Ti}(\text{C}_6\text{F}_5)_3(\text{OPPh}_3)]$, $[\text{Ti}(\text{C}_6\text{F}_5)_2\text{Cl}(\text{OPPh}_3)_2]$ and $[\text{Ti}(\text{C}_6\text{F}_5)_2(\text{OdppmO})_2][\text{Ti}(\text{C}_6\text{F}_5)_2\text{Cl}_2]$, 289
- Silyl substituents**
Synthesis and crystal structures of thermally stable titanocenes, 134
- Stibine**
Ruthenium hydrides bearing SbPh_3 and AsPh_3 ligands: characterization of the bis(dihydrogen) complexes $[\text{Cp}^*\text{Ru}(\text{H}_2)_2(\text{EPh}_3)]^+$ ($\text{Cp}^* = \text{C}_5\text{Me}_5$; $\text{E} = \text{Sb}, \text{As}$), 151
- s-trans* Coordination
Adaptable coordination modes of conjugated 1,3-diene: uniqueness of *s-trans* coordination, 5
- Termination**
Why propene is not polymerized by $(\text{Cp}^*_2\text{YH})_2$: reactions of yttrium alkyl complexes with alkenes produce allyl and vinyl yttrium complexes, 91
- Thallium**
Four- five- and six-co-ordinated pentafluorophenylthallium(III) complexes. Crystal structures of $[\text{Ti}(\text{C}_6\text{F}_5)_3(\text{OPPh}_3)]$, $[\text{Ti}(\text{C}_6\text{F}_5)_2\text{Cl}(\text{OPPh}_3)_2]$ and $[\text{Ti}(\text{C}_6\text{F}_5)_2(\text{OdppmO})_2][\text{Ti}(\text{C}_6\text{F}_5)_2\text{Cl}_2]$, 289
- Thiolates**
Synthesis of Group 10 polyfluorothiolate mono- and bi-nuclear complexes. Crystal structures of $[\text{Ni}(\text{SC}_6\text{HF}_4)_2(\text{dppe})]$, $[(\text{dppe})\text{Ni}(\mu\text{-SC}_6\text{HF}_4)_2\text{Pd}(\text{C}_6\text{F}_5)_2]$ and $[(\text{dppe})\text{Ni}(\mu\text{-SC}_6\text{F}_5)_2\text{Pd}(\text{C}_6\text{F}_5)_2]$, 98
- Through-space coupling**
Structural and dynamic studies in solution of anionic dinuclear azolato-bridged palladium(II) complexes, 108
- Tin alkyls**
Preparation of Ti(IV), Zr(IV) and Sn(IV) metal alkyls containing the (Pr^iO)₃SiCH₂ fragment, 256
- Tin organyls**
Mechanisms of Sn-to-Zr cyclopentadienyl transfer in the formation of Me_2Si -bridged zirconocenes from sila-stanna-tetrahydro-s-indacenes, 58
- Titanium**
Dialkyl titanium complexes that contain a sulfur-linked bis(phenolato) ligand: The structure of an olefin polymerization catalyst precursor, 158

- nolato) ligand: The structure of an olefin polymerization catalyst precursor, 158
- Hydrazonide titanium derivatives: Synthesis, characterization and catalytic activity in olefin polymerization. Molecular structure of $[\text{Ti}(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)\text{Cl}(\mu\text{-N}_2\text{CPh}_2)]_2$, 173
- Synthesis and crystal structures of thermally stable titanocenes, 134
- Titanium alkyls
- Preparation of Ti(IV), Zr(IV) and Sn(IV) metal alkyls containing the $(\text{Pr}^f\text{O})_3\text{SiCH}_2$ fragment, 256
- Titanocenes
- Synthesis and crystal structures of thermally stable titanocenes, 134
- Transmetalation
- Mechanisms of Sn-to-Zr cyclopentadienyl transfer in the formation of Me_2Si -bridged zirconocenes from sila-stanna-tetrahydro-s-indacenes, 58
- Transphobia
- Mono- and di-nuclear complexes of *ortho*-palladated and -platinated 4,4'-dimethylazobenzene with bis(diphenylphosphino)methane. More data on *transphobia*, 164
- Tungsten
- Theoretical analysis of bis(ethylene) complexes of molybdenum and tungsten, 83
- Vinyl complex
- Why propene is not polymerized by $(\text{Cp}^*_2\text{YH})_2$: reactions of yttrium alkyl complexes with alkenes produce allyl and vinyl yttrium complexes, 91
- X-ray crystallography
- C–Br versus C–H bond activation in palladium(II) cyclopalladated compounds. Crystal and molecular structure of $[\text{Pd}\{\text{C}_6\text{H}_4\text{C}(\text{H})=\text{NCy}\}(\text{MeCOCHCOMe})]$, 239
- Synthesis and crystal structures of thermally stable titanocenes, 134
- X-ray crystal structure analyses
- Structural characterization of Group 4 transition metal halide bis-Arduengo carbene complexes MCl_4L_2 : X-ray crystal structure analyses and DFT calculations, 192
- X-ray structures
- Synthesis and structures of crystalline bis(trimethylsilyl)methanido complexes of potassium, calcium and ytterbium, 263
- Ytterbium(II)
- Synthesis and structures of crystalline bis(trimethylsilyl)methanido complexes of potassium, calcium and ytterbium, 263
- Yttrium
- Synthesis, structure and catalytic activity of new iminophenolato complexes of scandium and yttrium, 63
- Zirconacyclopentanes
- Zirconocene mediated cyclization and isomerization of 1,3,6-heptatriene, 13
- Zirconium alkyls
- Preparation of Ti(IV), Zr(IV) and Sn(IV) metal alkyls containing the $(\text{Pr}^f\text{O})_3\text{SiCH}_2$ fragment, 256