



ELSEVIER

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SCIENCE @ DIRECT®

Journal of Organometallic Chemistry 664 (2002) 313–318

Journal  
of Organo  
metallic  
Chemistry[www.elsevier.com/locate/jorgchem](http://www.elsevier.com/locate/jorgchem)

## Subject Index of Volume 664

 $\pi$ -Acidity

Synthesis and electrochemical behaviour of  $\sigma, \eta^2$ -acetylide-bridged early-late complexes; The solid-state structure of  $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2]\text{Pd}(\text{PPh}_3)$ , 123

## Alkyl

Monomere Kupfer(I)-Alkyle mit  $\beta$ -Wasserstoffatomen und Kupfer(I)-Aryle mit kondensierten Aromaten; die Festkörperstruktur von  $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2]\text{Cu}^n\text{C}_4\text{H}_9$ , 142

## Alkyne

Stabilisierung niedervalenter Ni(CO)-Bausteine durch  $[\text{Ti}](\text{C}\equiv\text{CR})_2$ ; Reaktionsverhalten von  $\{[\text{Ti}](\text{C}\equiv\text{CR})_2\}\text{Ni}(\text{CO})$  gegenüber Triphenylphosphan und Phosphiten, 150

## Alkyne and diyne oligomerisation

The crystal structure of  $[\text{Mo}(\text{NCS})(\text{CO})_2(\eta^3\text{-C}_3\text{H}_5)(\text{NCMe})_2]\cdot\text{MeCN}$  and the reactions of  $\{\text{Mo}(\text{CO})_2(\eta^3\text{-C}_3\text{H}_5)^+\}$  containing species with symmetric alkynes, 176

## Alkyne/dialkyltungsten complexes

Steric isomerization of alkyne-dialkyltungsten complexes with a chelating diaryloxo ligand: crystal structures of  $\text{C}_s$ - and  $\text{C}_1$ - $\text{W}(\eta^2\text{-RC}\equiv\text{CR})[2,2'\text{-S}(4\text{-Me-6-R}'\text{-C}_6\text{H}_2\text{O})_2](\text{CH}_2\text{SiMe}_3)_2$ , 234

## Alkynyl

Monomere Kupfer(I)-Alkyle mit  $\beta$ -Wasserstoffatomen und Kupfer(I)-Aryle mit kondensierten Aromaten; die Festkörperstruktur von  $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2]\text{Cu}^n\text{C}_4\text{H}_9$ , 142

## Alkynyl-transfer

Stabilisierung niedervalenter Ni(CO)-Bausteine durch  $[\text{Ti}](\text{C}\equiv\text{CR})_2$ ; Reaktionsverhalten von  $\{[\text{Ti}](\text{C}\equiv\text{CR})_2\}\text{Ni}(\text{CO})$  gegenüber Triphenylphosphan und Phosphiten, 150

## Aluminium

Syntheses of polyaluminium compounds by hydroalumination reactions, chelating Lewis acids possessing two and four coordinatively unsaturated aluminium atoms, 110

## Amines

Facile syntheses of new multidentate (phosphino)amines: X-ray structure of  $1,4\text{-}\{(\text{OC})_4\text{Mo}(\text{Ph}_2\text{P})_2\text{NCH}_2\}_2\text{C}_6\text{H}_4$ , 294

## Amorphous polypropylene

Indenyl-amido titanium and zirconium dimethyl complexes: improved synthesis and use in propylene polymerization, 5

## Antimony selenium

Syntheses and structures of  $(\text{R}_2\text{Bi})_2\text{E}$  (E = S, Te) and *cyclo*- $(\text{RSbSe})_2[\text{W}(\text{CO})_5]_2$  [R = CH(SiMe<sub>3</sub>)<sub>2</sub>], 130

## Arene complexes

Osmacycles containing the  $\{(\text{Ph}_2\text{PE})_2\text{N}\}^-$  ligand (E = S or Se): synthesis and crystal structures, 85

## Arsenic

Reactivity of *cyclo*- $(\text{PhX})_6$  (X = As, P) towards  $[\text{M}_3\text{L}_2(\text{CO})_{10}]$  (M = Ru, L = CO or NCMe; M = Fe, L = CO), 27

## Aryl

Monomere Kupfer(I)-Alkyle mit  $\beta$ -Wasserstoffatomen und Kupfer(I)-Aryle mit kondensierten Aromaten; die Festkörperstruktur von  $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2]\text{Cu}^n\text{C}_4\text{H}_9$ , 142

## Aryl dithiocarboxylate

Reactions of cyclomanganated complexes with carbon disulfide: routes to  $\eta^2$ -aryldithiocarboxylate-Mn(CO)<sub>4</sub> complexes and to the trithiocarbonate complex  $(\mu_3\text{-CS}_3)_2\text{Mn}_4(\text{CO})_{16}$ , 288

## Benzene

Highly selective synthesis of biphenyl by the  $\text{Pd}(\text{OAc})_2/\text{HPA}/\text{O}_2/\text{AcOH}$  catalyst system, 59

## Biphenyl

Highly selective synthesis of biphenyl by the  $\text{Pd}(\text{OAc})_2/\text{HPA}/\text{O}_2/\text{AcOH}$  catalyst system, 59

## Bismuth

Syntheses and structures of  $(\text{R}_2\text{Bi})_2\text{E}$  (E = S, Te) and *cyclo*- $(\text{RSbSe})_2[\text{W}(\text{CO})_5]_2$  [R = CH(SiMe<sub>3</sub>)<sub>2</sub>], 130

## Boron

Alkyl derivatives of Stanna-closo-dodecaborate, 116

## Boron-nitrogen backbone

Stepwise assembly and structural characterization of oligonuclear ferrocene aggregates with boron-nitrogen backbone, 94

## Bulky ligands

Sodium and potassium derivatives of diphenyl(trimethylsilyl)methane, 182

## Butadienyl

Synthesis and reactivity of organochalogen ester substituted  $\eta^3$ -butadienyl complexes of Mo(II): crystal structure of  $[\text{MoCl}(\text{CO})_2(\eta^3\text{-CH}_2\text{C}(\text{COSePh})\text{C}=\text{CH}_2)(1,10\text{-phenanthroline})]\cdot 0.5\text{CH}_2\text{Cl}_2$ , 277

## 2-Butyne

Nitrile exchange reactions of  $[\text{W}(\text{CO})(\text{NCMe})\{\text{P}(\text{O}^i\text{Pr})_3\}_2(\eta^2\text{-MeC}_2\text{Me})][\text{BPh}_4]$ ; X-ray crystal structures of  $[\text{W}(\text{CO})(\text{L})\{\text{P}(\text{O}^i\text{Pr})_3\}_2(\eta^2\text{-MeC}_2\text{Me})][\text{BPh}_4]$  {L = NC<sup>i</sup>Pr, NC<sup>i</sup>Bu, NCCH<sub>2</sub>Ph, 1,2-C<sub>6</sub>H<sub>4</sub>(NCCH<sub>2</sub>)<sub>2</sub>}, 37

## Carbazole

Oxidative insertion into the N-H bond of carbazole, indole and pyrrole with zerovalent metals, 170

## Carbon disulfide

Reactions of cyclomanganated complexes with carbon disulfide: routes to  $\eta^2$ -aryldithiocarboxylate-Mn(CO)<sub>4</sub> complexes and to the trithiocarbonate complex  $(\mu_3\text{-CS}_3)_2\text{Mn}_4(\text{CO})_{16}$ , 288

## Carbonyl

Stabilisierung niedervalenter Ni(CO)-Bausteine durch  $[\text{Ti}](\text{C}\equiv\text{CR})_2$ ; Reaktionsverhalten von  $\{[\text{Ti}](\text{C}\equiv\text{CR})_2\}\text{Ni}(\text{CO})$  gegenüber Triphenylphosphan und Phosphiten, 150

Synthesis and reactions of the cationic seven-coordinate complexes,  $[\text{MoI}(\text{CO})_3(\text{NCR})\{\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2\}][\text{BF}_4]$  (R = Me, n = 1, 2 and 3; R = Et, n = 1). Crystal structures of  $[\text{MoI}_2(\text{CO})_2(\text{PPh}_3)\{\text{Ph}_2\text{P}(\text{CH}_2)\text{PPh}_2\}]$ ,  $[\text{MoI}(\text{CO})_2\{\text{Ph}_2\text{P}(\text{CH}_2)\text{PPh}_2\}_2][\text{BF}_4]\cdot\text{CH}_2\text{Cl}_2$ ,  $[\text{MoCl}_3\text{O}\{\text{Ph}_2\text{PO}(\text{CH}_2)\text{OPPh}_2\}]$  and  $[\text{Mo}_3(\mu_3\text{-I})_2\{\mu_2\text{-Ph}_2\text{P}(\text{CH}_2)\text{PPh}_2\}_3]\text{I}$ , 45

## Carbonyl complexes

Facile syntheses of new multidentate (phosphino)amines: X-ray structure of  $1,4\text{-}\{(\text{OC})_4\text{Mo}(\text{Ph}_2\text{P})_2\text{NCH}_2\}_2\text{C}_6\text{H}_4$ , 294

## Catalysis

New complexes of palladium(II) with chelating heterocyclic nitrogen ligands. Characterization and catalytic activity in CO-styrene copolymerization. Crystal structure of  $[\text{Pd}(\text{bipy})_2][\text{BAR}'_4]_2$ , Ar' = [3,5-(CF<sub>3</sub>)<sub>2</sub>C<sub>6</sub>H<sub>3</sub>], 77

## Catalyst separation

Polyether phosphine oxide induced phase separable homogeneous catalysis for hydroformylation of higher olefins, 1

## Cationic

Nitrile exchange reactions of  $[\text{WI}(\text{CO})(\text{NCMe})\{\text{P}(\text{O}^i\text{Pr})_3\}_2(\eta^2\text{-MeC}_2\text{Me})][\text{BPh}_4]$ ; X-ray crystal structures of  $[\text{WI}(\text{CO})(\text{L})\{\text{P}(\text{O}^i\text{Pr})_3\}_2(\eta^2\text{-MeC}_2\text{Me})][\text{BPh}_4]$   $\{\text{L} = \text{NC}^i\text{Pr}, \text{NC}^i\text{Bu}, \text{NCCH}_2\text{Ph}, 1,2\text{-C}_6\text{H}_4(\text{NCCH}_2)_2\}$ , 37

Synthesis and reactions of the cationic seven-coordinate complexes,  $[\text{MoI}(\text{CO})_3(\text{NCR})\{\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2\}][\text{BF}_4]$  ( $\text{R} = \text{Me}, n = 1, 2$  and  $3; \text{R} = \text{Et}, n = 1$ ). Crystal structures of  $[\text{MoI}_2(\text{CO})_2(\text{PPh}_3)\{\text{Ph}_2\text{P}(\text{CH}_2)\text{PPh}_2\}]$ ,  $[\text{MoI}(\text{CO})_2\{\text{Ph}_2\text{P}(\text{CH}_2)\text{PPh}_2\}_2][\text{BF}_4] \cdot \text{CH}_2\text{Cl}_2$ ,  $[\text{MoCl}_3\text{O}\{\text{Ph}_2\text{PO}(\text{CH}_2)\text{OPPh}_2\}]$  and  $[\text{Mo}_3(\mu_3\text{-I})_2\{\mu_2\text{-Ph}_2\text{P}(\text{CH}_2)\text{PPh}_2\}_3]\text{I}$ , 45

C–H... $\pi$  intermolecular contact

Preparation and structural characterization of six new diorganotin(IV) complexes of the  $\text{R}_2\text{Sn}(\text{SaleanH}_2)$  and  $\text{R}_2\text{Sn}(\text{SalceanH}_2)$  type ( $\text{R} = \text{Me}, ^i\text{Bu}, \text{Ph}$ ), 188

## Chiral

Synthesis, reaction and enantiomeric resolution of chiral clusters containing  $\text{SRuNiM}$  ( $\text{M} = \text{Mo}, \text{W}$ ) core, 201

*cis* and *trans* isomers

Preparation and structural characterization of six new diorganotin(IV) complexes of the  $\text{R}_2\text{Sn}(\text{SaleanH}_2)$  and  $\text{R}_2\text{Sn}(\text{SalceanH}_2)$  type ( $\text{R} = \text{Me}, ^i\text{Bu}, \text{Ph}$ ), 188

## Cluster

Alkylderivatives of Stanna-*closo*-dodecaborate, 116

Reactivity of *cyclo*-(PhX)<sub>6</sub> ( $\text{X} = \text{As}, \text{P}$ ) towards  $[\text{M}_3\text{L}_2(\text{CO})_{10}]$  ( $\text{M} = \text{Ru}, \text{L} = \text{CO}$  or  $\text{NCMe}$ ;  $\text{M} = \text{Fe}, \text{L} = \text{CO}$ ), 27

Synthesis, reaction and enantiomeric resolution of chiral clusters containing  $\text{SRuNiM}$  ( $\text{M} = \text{Mo}, \text{W}$ ) core, 201

## Copolymerization

New complexes of palladium(II) with chelating heterocyclic nitrogen ligands. Characterization and catalytic activity in CO–styrene copolymerization. Crystal structure of  $[\text{Pd}(\text{bipy})_2][\text{BAR}'_4]_2$ ,  $\text{Ar}' = [3,5\text{-}(\text{CF}_3)_2\text{C}_6\text{H}_3]$ , 77

## Copper(I)

Monomere Kupfer(I)-Alkyle mit  $\beta$ -Wasserstoffatomen und Kupfer(I)-Aryle mit kondensierten Aromaten; die Festkörperstruktur von  $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2]\text{Cu}^i\text{C}_4\text{H}_9$ , 142

## CO-substitution

Stabilisierung niedervalenter  $\text{Ni}(\text{CO})$ -Bausteine durch  $[\text{Ti}](\text{C}\equiv\text{CR})_2$ ; Reaktionsverhalten von  $\{[\text{Ti}](\text{C}\equiv\text{CR})_2\}\text{Ni}(\text{CO})$  gegenüber Triphenylphosphan und Phosphiten, 150

## Crystal structure

Facile syntheses of new multidentate (phosphino)amines: X-ray structure of  $1,4\text{-}\{(\text{OC})_4\text{Mo}(\text{Ph}_2\text{P})_2\text{NCH}_2\}_2\text{C}_6\text{H}_4$ , 294

Osmacycles containing the  $\{(\text{Ph}_2\text{PE})_2\text{N}\}^-$  ligand ( $\text{E} = \text{S}$  or  $\text{Se}$ ): synthesis and crystal structures, 85

Synthesis and reactions of the cationic seven-coordinate complexes,  $[\text{MoI}(\text{CO})_3(\text{NCR})\{\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2\}][\text{BF}_4]$  ( $\text{R} = \text{Me}, n = 1, 2$  and  $3; \text{R} = \text{Et}, n = 1$ ). Crystal structures of  $[\text{MoI}_2(\text{CO})_2(\text{PPh}_3)\{\text{Ph}_2\text{P}(\text{CH}_2)\text{PPh}_2\}]$ ,  $[\text{MoI}(\text{CO})_2\{\text{Ph}_2\text{P}(\text{CH}_2)\text{PPh}_2\}_2][\text{BF}_4] \cdot \text{CH}_2\text{Cl}_2$ ,  $[\text{MoCl}_3\text{O}\{\text{Ph}_2\text{PO}(\text{CH}_2)\text{OPPh}_2\}]$  and  $[\text{Mo}_3(\mu_3\text{-I})_2\{\mu_2\text{-Ph}_2\text{P}(\text{CH}_2)\text{PPh}_2\}_3]\text{I}$ , 45

Synthesis, reaction and enantiomeric resolution of chiral clusters containing  $\text{SRuNiM}$  ( $\text{M} = \text{Mo}, \text{W}$ ) core, 201

## Crystal structures

Nitrile exchange reactions of  $[\text{WI}(\text{CO})(\text{NCMe})\{\text{P}(\text{O}^i\text{Pr})_3\}_2(\eta^2\text{-MeC}_2\text{Me})][\text{BPh}_4]$ ; X-ray crystal structures of  $[\text{WI}(\text{CO})(\text{L})\{\text{P}(\text{O}^i\text{Pr})_3\}_2(\eta^2\text{-MeC}_2\text{Me})][\text{BPh}_4]$   $\{\text{L} = \text{NC}^i\text{Pr}, \text{NC}^i\text{Bu}, \text{NCCH}_2\text{Ph}, 1,2\text{-C}_6\text{H}_4(\text{NCCH}_2)_2\}$ , 37

Steric isomerization of alkyne–dialkyltungsten complexes with a chelating diaryloxo ligand: crystal structures of  $\text{C}_s$ - and  $\text{C}_1$ - $\text{W}(\eta^2\text{-RC}\equiv\text{CR})[2,2'\text{-S}(4\text{-Me-6-R}'\text{C}_6\text{H}_2\text{O})_2](\text{CH}_2\text{SiMe}_3)_2$ , 234

Synthesis, spectroscopic and structural aspects of some tetraorganodistannoxanes with internally functionalized oxime. 2.

Crystal and molecular structure of  $\{[\text{Me}_2\text{Sn}(\text{ON}=\text{C}(\text{Me})\text{-C}_5\text{H}_4\text{N})_2\text{O}\}_2 \cdot 2[2\text{-NC}_5\text{H}_4(\text{Me})\text{C}=\text{NOH}]$ , 66

## Cyclic voltammetry

Lithiation of ferrocenylamines and vanadium dinitrogen chemistry, 245

Synthesis and electrochemical behaviour of  $\sigma, \eta^2$ -acetylide-bridged early–late complexes; The solid-state structure of  $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2]\text{Pd}(\text{PPh}_3)$ , 123

## Cyclomanganates

Reactions of cyclomanganated complexes with carbon disulfide: routes to  $\eta^2$ -aryldithiocarboxylate- $\text{Mn}(\text{CO})_4$  complexes and to the trithiocarbonate complex  $(\mu_3\text{-CS}_3)_2\text{Mn}_4(\text{CO})_{16}$ , 288

## Cyclopentadienyl ligands

Arylimido niobium(V) complexes: mononuclear and dendritic derivatives, 258

## Dendrimers

Arylimido niobium(V) complexes: mononuclear and dendritic derivatives, 258

## Density functional theory

Specific solvent effect on  $\text{R}_2\text{ZrCl}_2$  ( $\text{R} = \text{butyl}, \text{ethyl}$ ) reactivity, a density functional study, 268

## Dinitrogen uptake

Lithiation of ferrocenylamines and vanadium dinitrogen chemistry, 245

## Diphenylmethyl

Sodium and potassium derivatives of diphenyl(trimethylsilyl)methane, 182

## Early–late complex

Synthesis and electrochemical behaviour of  $\sigma, \eta^2$ -acetylide-bridged early–late complexes; The solid-state structure of  $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2]\text{Pd}(\text{PPh}_3)$ , 123

## Electronic absorption spectra

Gas-phase electronic absorption spectra of metallocene dihydrides  $\text{M}(\eta^5\text{-C}_5\text{H}_5)_2\text{H}_2$  ( $\text{M} = \text{Mo}, \text{W}$ ), 106

## Enantiomeric resolution

Synthesis, reaction and enantiomeric resolution of chiral clusters containing  $\text{SRuNiM}$  ( $\text{M} = \text{Mo}, \text{W}$ ) core, 201

*fac*–*fac* and *fac*–*mer* configuration

Preparation and structural characterization of six new diorganotin(IV) complexes of the  $\text{R}_2\text{Sn}(\text{SaleanH}_2)$  and  $\text{R}_2\text{Sn}(\text{SalceanH}_2)$  type ( $\text{R} = \text{Me}, ^i\text{Bu}, \text{Ph}$ ), 188

## Ferrocenylamines

Lithiation of ferrocenylamines and vanadium dinitrogen chemistry, 245

## Formate

The reaction of samarium(II) with manganese carbonyl: unexpected conversion of CO to formate. X-ray crystal structures of  $[\text{Sm}(\text{Tp}^{\text{Me}2})_2]\text{Mn}(\text{CO})_5$  and  $\{[\text{Sm}(\text{Tp}^{\text{Me}2})_2](\mu\text{-HCO}_2)\}_2\text{Mn}(\text{CO})_5$  ( $\text{Tp}^{\text{Me}2} = \text{HB}(3,5\text{-dimethylpyrazolyl})_2$ ), 298

## Fullerenes

Oxorhenium(V) '3+2' mixed-ligand complexes carrying the SNO/SN donor and  $\text{C}_{60}$ , 208

## Hemilability

Hemilability of the primary amine–metal bond in polyamine (Group 6) metal carbonyl complexes, 214

 $\text{H}_3\text{PMO}_{12}\text{O}_{40}$ 

Highly selective synthesis of biphenyl by the  $\text{Pd}(\text{OAc})_2/\text{HPA}/\text{O}_2/\text{AcOH}$  catalyst system, 59

 $\beta$ -Hydride transfer

Monomere Kupfer(I)-Alkyle mit  $\beta$ -Wasserstoffatomen und Kupfer(I)-Aryle mit kondensierten Aromaten; die Festkörperstruktur von  $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2]\text{Cu}^i\text{C}_4\text{H}_9$ , 142

## Hydroalumination

Syntheses of polyaluminium compounds by hydroalumination reactions, chelating Lewis acids possessing two and four coordinatively unsaturated aluminium atoms, 110

## Hydroformylation

Polyether phosphine oxide induced phase separable homogeneous catalysis for hydroformylation of higher olefins, 1

## Hydrogen bridging

Alkylderivatives of Stanna-*closa*-dodecaborate, 116

 $\beta$ -Hydrogen migration

Specific solvent effect on  $R_2ZrCl_2$  (R = butyl, ethyl) reactivity, a density functional study, 268

## Hydrolysis

The reaction of samarium(II) with manganese carbonyl: unexpected conversion of CO to formate. X-ray crystal structures of  $[Sm(Tp^{Me_2})_2]Mn(CO)_5$  and  $\{[Sm(Tp^{Me_2})_2]_2(\mu-HCO_2)\}Mn(CO)_5$  ( $Tp^{Me_2}$  = HB(3,5-dimethylpyrazolyl), 298

## Imines

Hemilability of the primary amine–metal bond in polyamine- (Group 6) metal carbonyl complexes, 214

## Indenyl amido titanium dimethyl complexes

Indenyl-amido titanium and zirconium dimethyl complexes: improved synthesis and use in propylene polymerization, 5

## Indole

Oxidative insertion into the N–H bond of carbazole, indole and pyrrole with zerovalent metals, 170

## Inorganic heterocycle

Osmacycles containing the  $\{(Ph_2PE)_2N\}^-$  ligand (E = S or Se): synthesis and crystal structures, 85

## Insertion

Reactions of cyclomanganated complexes with carbon disulfide: routes to  $\eta^2$ -aryldithiocarboxylate- $Mn(CO)_4$  complexes and to the trithiocarbonate complex  $(\mu_3-CS_3)_2Mn_4(CO)_{16}$ , 288

## Intermolecular bonds

Crystal and molecular structure of ( $\alpha$ -naphthyl)TeI<sub>3</sub>, 306

## Internally functionalized oxime

Synthesis, spectroscopic and structural aspects of some tetraorganodistannoxanes with internally functionalized oxime. 2. Crystal and molecular structure of  $\{[Me_2Sn(ON=C(Me)-C_5H_4N)]_2O\}_2 \cdot 2[2-NC_5H_4(Me)C=NOH]$ , 66

## Iodide

Synthesis and reactions of the cationic seven-coordinate complexes,  $[MoI(CO)_3(NCR)\{Ph_2P(CH_2)_n PPh_2\}][BF_4]$  (R = Me,  $n = 1, 2$  and 3; R = Et,  $n = 1$ ). Crystal structures of  $[MoI_2(CO)_2(PPh_3)\{Ph_2P(CH_2)PPh_2\}]$ ,  $[MoI(CO)_2\{Ph_2P(CH_2)PPh_2\}_2][BF_4] \cdot CH_2Cl_2$ ,  $[MoCl_3O\{Ph_2PO(CH_2)OPPh_2\}]$  and  $[Mo_3(\mu_3-I)_2\{\mu_2-Ph_2P(CH_2)PPh_2\}_3]I$ , 45

## Iodo

Nitrile exchange reactions of  $[WI(CO)(NCMe)\{P(O^iPr)_3\}_2(\eta^2-MeC_2Me)][BPh_4]$ ; X-ray crystal structures of  $[WI(CO)(L)\{P(O^iPr)_3\}_2(\eta^2-MeC_2Me)][BPh_4]$   $\{L = NC^iPr, NC^tBu, NCCH_2Ph, 1,2-C_6H_4(NCCH_2)_2\}$ , 37

## Iron

Preparation and organic transformation reactions of  $CpFe(CO)_2-SCO-3-C_6H_4-SO_2Cl$ , 228

Reactivity of *cyclo*-(PhX)<sub>6</sub> (X = As, P) towards  $[M_3L_2(CO)_{10}]$  (M = Ru, L = CO or NCMe; M = Fe, L = CO), 27

## Lanthanide

The reaction of samarium(II) with manganese carbonyl: unexpected conversion of CO to formate. X-ray crystal structures of  $[Sm(Tp^{Me_2})_2]Mn(CO)_5$  and  $\{[Sm(Tp^{Me_2})_2]_2(\mu-HCO_2)\}Mn(CO)_5$  ( $Tp^{Me_2}$  = HB(3,5-dimethylpyrazolyl), 298

## Lithiation

Lithiation of ferrocenylamines and vanadium dinitrogen chemistry, 245

## Manganese

Reactions of cyclomanganated complexes with carbon disulfide: routes to  $\eta^2$ -aryldithiocarboxylate- $Mn(CO)_4$  complexes and to the trithiocarbonate complex  $(\mu_3-CS_3)_2Mn_4(CO)_{16}$ , 288

The reaction of samarium(II) with manganese carbonyl: unexpected conversion of CO to formate. X-ray crystal structures of  $[Sm(Tp^{Me_2})_2]Mn(CO)_5$  and  $\{[Sm(Tp^{Me_2})_2]_2(\mu-HCO_2)\}Mn(CO)_5$  ( $Tp^{Me_2}$  = HB(3,5-dimethylpyrazolyl), 298

## Metal carbonyl complexes

Hemilability of the primary amine–metal bond in polyamine- (Group 6) metal carbonyl complexes, 214

## Metallocene hydrides

Gas-phase electronic absorption spectra of metallocene dihydrides  $M(\eta^5-C_5H_5)_2H_2$  (M = Mo, W), 106

## Mixed ligand complexes

Oxorhenium(V) '3+2' mixed-ligand complexes carrying the SNO/SN donor and C<sub>60</sub>, 208

## Molybdenum

Facile syntheses of new multidentate (phosphino)amines: X-ray structure of 1,4- $\{(OC)_4Mo(Ph_2P)_2NCH_2\}_2C_6H_4$ , 294

Gas-phase electronic absorption spectra of metallocene dihydrides  $M(\eta^5-C_5H_5)_2H_2$  (M = Mo, W), 106

Synthesis and reactivity of organochalogen ester substituted  $\eta^3$ -butadienyl complexes of Mo(II): crystal structure of  $[MoCl(CO)_2(\eta^3-CH_2C(COSePh)C=CH_2)(1,10-phenanthroline)] \cdot 0.5 CH_2Cl_2$ , 277

## Molybdenum(II)

Synthesis and reactions of the cationic seven-coordinate complexes,  $[MoI(CO)_3(NCR)\{Ph_2P(CH_2)_n PPh_2\}][BF_4]$  (R = Me,  $n = 1, 2$  and 3; R = Et,  $n = 1$ ). Crystal structures of  $[MoI_2(CO)_2(PPh_3)\{Ph_2P(CH_2)PPh_2\}]$ ,  $[MoI(CO)_2\{Ph_2P(CH_2)PPh_2\}_2][BF_4] \cdot CH_2Cl_2$ ,  $[MoCl_3O\{Ph_2PO(CH_2)OPPh_2\}]$  and  $[Mo_3(\mu_3-I)_2\{\mu_2-Ph_2P(CH_2)PPh_2\}_3]I$ , 45

## Molybdenum(II)allyl

The crystal structure of  $[Mo(NCS)(CO)_2(\eta^3-C_3H_5)(NCMe)_2] \cdot MeCN$  and the reactions of  $\{Mo(CO)_2(\eta^3-C_3H_5)^+\}$  containing species with symmetric alkynes, 176

## Mössbauer spectra

Lithiation of ferrocenylamines and vanadium dinitrogen chemistry, 245

## N-Heterocyclic carbene ligand

Synthesis and structure of 1,2,4-triazol-2-ium-5-ylidene complexes of Hg(II), Pd(II), Ni(II), Ni(0), Rh(I) and Ir(I), 70

## Nickel

Oxidative insertion into the N–H bond of carbazole, indole and pyrrole with zerovalent metals, 170

Stabilisierung niedervalenter Ni(CO)-Bausteine durch  $[Ti](C \equiv CR)_2$ ; Reaktionsverhalten von  $\{[Ti](C \equiv CR)_2\}Ni(CO)$  gegenüber Triphenylphosphan und Phosphiten, 150

Synthesis and electrochemical behaviour of  $\sigma, \eta^2$ -acetylide-bridged early–late complexes; The solid-state structure of  $[(\eta^5-C_5H_4SiMe_3)_2Ti(C \equiv CSiMe_3)_2]Pd(PPh_3)$ , 123

## Niobium

Arylimido niobium(V) complexes: mononuclear and dendritic derivatives, 258

## Nitrile

Nitrile exchange reactions of  $[WI(CO)(NCMe)\{P(O^iPr)_3\}_2(\eta^2-MeC_2Me)][BPh_4]$ ; X-ray crystal structures of  $[WI(CO)(L)\{P(O^iPr)_3\}_2(\eta^2-MeC_2Me)][BPh_4]$   $\{L = NC^iPr, NC^tBu, NCCH_2Ph, 1,2-C_6H_4(NCCH_2)_2\}$ , 37

Synthesis and reactions of the cationic seven-coordinate complexes,  $[MoI(CO)_3(NCR)\{Ph_2P(CH_2)_n PPh_2\}][BF_4]$  (R = Me,  $n = 1, 2$  and 3; R = Et,  $n = 1$ ). Crystal structures of  $[MoI_2(CO)_2(PPh_3)\{Ph_2P(CH_2)PPh_2\}]$ ,  $[MoI(CO)_2\{Ph_2P(CH_2)PPh_2\}_2][BF_4] \cdot CH_2Cl_2$ ,  $[MoCl_3O\{Ph_2PO(CH_2)OPPh_2\}]$  and  $[Mo_3(\mu_3-I)_2\{\mu_2-Ph_2P(CH_2)PPh_2\}_3]I$ , 45

## Nitrogen ligands

New complexes of palladium(II) with chelating heterocyclic nitrogen ligands. Characterization and catalytic activity in CO–styrene copolymerization. Crystal structure of  $[\text{Pd}(\text{bipy})_2] - [\text{BAR}'_4]_2$ ,  $\text{Ar}' = [3,5\text{-}(\text{CF}_3)_2\text{C}_6\text{H}_3]$ , 77

## N Ligands

Arylimido niobium(V) complexes: mononuclear and dendritic derivatives, 258

## NMR spectroscopy

Facile syntheses of new multidentate (phosphino)amines: X-ray structure of 1,4- $\{(\text{OC})_4\text{Mo}(\text{Ph}_2\text{P})_2\text{NCH}_2\}_2\text{C}_6\text{H}_4$ , 294

Osmacycles containing the  $\{(\text{Ph}_2\text{PE})_2\text{N}\}^-$  ligand (E = S or Se): synthesis and crystal structures, 85

## Nucleophilic substitution

Alkylderivatives of Stanna-*closo*-dodecaborate, 116

Preparation and organic transformation reactions of  $\text{CpFe}(\text{CO})_2\text{-SCO-3-C}_6\text{H}_4\text{-SO}_2\text{Cl}$ , 228

## Oligonuclear ferrocene complexes

Stepwise assembly and structural characterization of oligonuclear ferrocene aggregates with boron–nitrogen backbone, 94

## Organochalcogen

Synthesis and reactivity of organochalogen ester substituted  $\eta^3$ -butadienyl complexes of Mo(II): crystal structure of  $[\text{MoCl}(\text{CO})_2(\eta^3\text{-CH}_2\text{C}(\text{COSePh})\text{C}=\text{CH}_2)(1,10\text{-phenanthroline})] \cdot 0.5\text{CH}_2\text{Cl}_2$ , 277

Organometallic  $\pi$ -tweezer

Monomere Kupfer(I)-Alkyle mit  $\beta$ -Wasserstoffatomen und Kupfer(I)-Aryle mit kondensierten Aromaten; die Festkörperstruktur von  $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2]\text{Cu}^u\text{C}_4\text{H}_9$ , 142

## Organotellurium triiodides

Crystal and molecular structure of ( $\alpha$ -naphthyl)TeI<sub>3</sub>, 306

## Osmacycle

Osmacycles containing the  $\{(\text{Ph}_2\text{PE})_2\text{N}\}^-$  ligand (E = S or Se): synthesis and crystal structures, 85

## Oxidative dimerization

Highly selective synthesis of biphenyl by the  $\text{Pd}(\text{OAc})_2/\text{HPA}/\text{O}_2/\text{AcOH}$  catalyst system, 59

## Palladium

Oxidative insertion into the N–H bond of carbazole, indole and pyrrole with zerovalent metals, 170

Synthesis and electrochemical behaviour of  $\sigma,\eta^2$ -acetylide-bridged early–late complexes; The solid-state structure of  $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2]\text{Pd}(\text{PPh}_3)$ , 123

## Palladium catalysis

One-pot palladium-catalyzed phosphination of aryl iodides with  $\text{Ph}_2\text{PSnR}_3$ , 223

## Palladium complexes

New complexes of palladium(II) with chelating heterocyclic nitrogen ligands. Characterization and catalytic activity in CO–styrene copolymerization. Crystal structure of  $[\text{Pd}(\text{bipy})_2][\text{BAR}'_4]_2$ ,  $\text{Ar}' = [3,5\text{-}(\text{CF}_3)_2\text{C}_6\text{H}_3]$ , 77

## P–C cross-coupling

One-pot palladium-catalyzed phosphination of aryl iodides with  $\text{Ph}_2\text{PSnR}_3$ , 223

Pd(OAc)<sub>2</sub>

Highly selective synthesis of biphenyl by the  $\text{Pd}(\text{OAc})_2/\text{HPA}/\text{O}_2/\text{AcOH}$  catalyst system, 59

## Pentamethylcyclopentadienyl

Cyclic thioether and acyclic thioether–thiolate complexes of pentamethylcyclopentadienyl ruthenium(II, III), 161

## Phosphane

Stabilisierung niedervalenter Ni(CO)-Bausteine durch  $[\text{Ti}](\text{C}\equiv\text{CR})_2$ ; Reaktionsverhalten von  $\{[\text{Ti}](\text{C}\equiv\text{CR})_2\}\text{Ni}(\text{CO})$  gegenüber Triphenylphosphan und Phosphiten, 150

## Phosphine

Synthesis and reactions of the cationic seven-coordinate complexes,  $[\text{MoI}(\text{CO})_3(\text{NCR})\{\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2\}][\text{BF}_4]$  (R = Me,  $n = 1, 2$  and 3; R = Et,  $n = 1$ ). Crystal structures of  $[\text{MoI}_2(\text{CO})_2(\text{PPh}_3)\{\text{Ph}_2\text{P}(\text{CH}_2)\text{PPh}_2\}]$ ,  $[\text{MoI}(\text{CO})_2\{\text{Ph}_2\text{P}(\text{CH}_2)\text{PPh}_2\}_2][\text{BF}_4] \cdot \text{CH}_2\text{-Cl}_2$ ,  $[\text{MoCl}_3\text{O}\{\text{Ph}_2\text{PO}(\text{CH}_2)\text{OPPh}_2\}]$  and  $[\text{Mo}_3(\mu_3\text{-I})_2\{\mu_2\text{-Ph}_2\text{P}(\text{CH}_2)\text{PPh}_2\}_3]\text{I}$ , 45

## Phosphine oxide

Polyether phosphine oxide induced phase separable homogeneous catalysis for hydroformylation of higher olefins, 1

## Phosphite

Nitrile exchange reactions of  $[\text{WI}(\text{CO})(\text{NCMe})\{\text{P}(\text{O}^i\text{Pr})_3\}_2(\eta^2\text{-MeC}_2\text{Me})][\text{BPh}_4]$ ; X-ray crystal structures of  $[\text{WI}(\text{CO})(\text{L})\{\text{P}(\text{O}^i\text{Pr})_3\}_2(\eta^2\text{-MeC}_2\text{Me})][\text{BPh}_4]$  {L = NC<sup>i</sup>Pr, NC<sup>i</sup>Bu, NCCH<sub>2</sub>Ph, 1,2-C<sub>6</sub>H<sub>4</sub>(NCCH<sub>2</sub>)<sub>2</sub>}, 37

Stabilisierung niedervalenter Ni(CO)-Bausteine durch  $[\text{Ti}](\text{C}\equiv\text{CR})_2$ ; Reaktionsverhalten von  $\{[\text{Ti}](\text{C}\equiv\text{CR})_2\}\text{Ni}(\text{CO})$  gegenüber Triphenylphosphan und Phosphiten, 150

## Phosphorus

Reactivity of *cyclo*-(PhX)<sub>6</sub> (X = As, P) towards  $[\text{M}_3\text{L}_2(\text{CO})_{10}]$  (M = Ru, L = CO or NCMc; M = Fe, L = CO), 27

## Phosphorus ligands

Facile syntheses of new multidentate (phosphino)amines: X-ray structure of 1,4- $\{(\text{OC})_4\text{Mo}(\text{Ph}_2\text{P})_2\text{NCH}_2\}_2\text{C}_6\text{H}_4$ , 294

Osmacycles containing the  $\{(\text{Ph}_2\text{PE})_2\text{N}\}^-$  ligand (E = S or Se): synthesis and crystal structures, 85

## Platinum

Oxidative insertion into the N–H bond of carbazole, indole and pyrrole with zerovalent metals, 170

## Polyamines

Hemilability of the primary amine–metal bond in polyamine–(Group 6) metal carbonyl complexes, 214

## Polymeric superstructures

Crystal and molecular structure of ( $\alpha$ -naphthyl)TeI<sub>3</sub>, 306

## Potassium

Sodium and potassium derivatives of diphenyl(trimethylsilyl)methane, 182

## Pyrazolylborate

The reaction of samarium(II) with manganese carbonyl: unexpected conversion of CO to formate. X-ray crystal structures of  $[\text{Sm}(\text{Tp}^{\text{Me}_2})_2]\text{Mn}(\text{CO})_5$  and  $\{[\text{Sm}(\text{Tp}^{\text{Me}_2})_2](\mu\text{-HCO}_2)\}\text{Mn}(\text{CO})_5$  ( $\text{Tp}^{\text{Me}_2} = \text{HB}(3,5\text{-dimethylpyrazolyl})$ ), 298

## Pyrrole

Oxidative insertion into the N–H bond of carbazole, indole and pyrrole with zerovalent metals, 170

## Quaternisation

Lithiation of ferrocenylamines and vanadium dinitrogen chemistry, 245

## Radiopharmaceuticals

Oxorhenium(V) '3+2' mixed-ligand complexes carrying the SNO/SN donor and C<sub>60</sub>, 208

## Reactivity studies

Synthesis and reactivity of organochalogen ester substituted  $\eta^3$ -butadienyl complexes of Mo(II): crystal structure of  $[\text{MoCl}(\text{CO})_2(\eta^3\text{-CH}_2\text{C}(\text{COSePh})\text{C}=\text{CH}_2)(1,10\text{-phenanthroline})] \cdot 0.5\text{CH}_2\text{Cl}_2$ , 277

## Rhenium

Oxorhenium(V) '3+2' mixed-ligand complexes carrying the SNO/SN donor and C<sub>60</sub>, 208

## Ring-opening metathesis polymerization

Steric isomerization of alkyne–dialkyltungsten complexes with a chelating diaryloxo ligand: crystal structures of C<sub>s</sub>- and C<sub>1</sub>- $W(\eta^2\text{-RC}\equiv\text{CR})[2,2' - \text{S}(4 - \text{Me} - 6 - \text{R}'\text{C}_6\text{H}_2\text{O})_2](\text{CH}_2\text{SiMe}_3)_2$ , 234

## Ruthenium

Cyclic thioether and acyclic thioether–thiolate complexes of pentamethylcyclopentadienyl ruthenium(II, III), 161

Reactivity of *cyclo*-(PhX)<sub>6</sub> (X = As, P) towards [M<sub>3</sub>L<sub>2</sub>(CO)<sub>10</sub>] (M = Ru, L = CO or NCMc; M = Fe, L = CO), 27

## Rydberg transitions

Gas-phase electronic absorption spectra of metallocene dihydrides M(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>H<sub>2</sub> (M = Mo, W), 106

## Salen and Salan complexes with tin(IV)

Preparation and structural characterization of six new diorganotin(IV) complexes of the R<sub>2</sub>Sn(SaleanH<sub>2</sub>) and R<sub>2</sub>Sn(SalanH<sub>2</sub>) type (R = Me, <sup>n</sup>Bu, Ph), 188

## Samarium

The reaction of samarium(II) with manganese carbonyl: unexpected conversion of CO to formate. X-ray crystal structures of [Sm(Tp<sup>Me2</sup>)<sub>2</sub>]Mn(CO)<sub>5</sub> and {[Sm(Tp<sup>Me2</sup>)<sub>2</sub>](μ-HCO<sub>2</sub>)}Mn(CO)<sub>5</sub> (Tp<sup>Me2</sup> = HB(3,5-dimethylpyrazolyl)), 298

## Schiff bases

Oxorhenium(V) '3 + 2' mixed-ligand complexes carrying the SNO/SN donor and C<sub>60</sub>, 208

## Selenoester

Synthesis and reactivity of organochalogen ester substituted η<sup>3</sup>-butadienyl complexes of Mo(II): crystal structure of [MoCl(CO)<sub>2</sub>(η<sup>3</sup>-CH<sub>2</sub>C(COSePh)C=CH<sub>2</sub>)(1,10-phenanthroline)]·0.5 CH<sub>2</sub>Cl<sub>2</sub>, 277

## Seven-coordinate

Synthesis and reactions of the cationic seven-coordinate complexes, [MoI(CO)<sub>3</sub>(NCR){Ph<sub>2</sub>P(CH<sub>2</sub>)<sub>n</sub>PPh<sub>2</sub>}] [BF<sub>4</sub>] (R = Me, n = 1, 2 and 3; R = Et, n = 1). Crystal structures of [MoI<sub>2</sub>(CO)<sub>2</sub>(PPh<sub>3</sub>){Ph<sub>2</sub>P(CH<sub>2</sub>)PPh<sub>2</sub>}], [MoI(CO)<sub>2</sub>{Ph<sub>2</sub>P(CH<sub>2</sub>)PPh<sub>2</sub>}] [BF<sub>4</sub>]·CH<sub>2</sub>Cl<sub>2</sub>, [MoCl<sub>3</sub>O{Ph<sub>2</sub>PO(CH<sub>2</sub>)OPPh<sub>2</sub>}] and [Mo<sub>3</sub>(μ<sub>3</sub>-I)<sub>2</sub>{μ<sub>2</sub>-Ph<sub>2</sub>P(CH<sub>2</sub>)PPh<sub>2</sub>}]<sub>3</sub>I, 45

## Sodium

Sodium and potassium derivatives of diphenyl(trimethylsilyl)methane, 182

## Solid-state structure

Synthesis and electrochemical behaviour of σ,η<sup>2</sup>-acetylide-bridged early–late complexes; The solid-state structure of [(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>SiMe<sub>3</sub>)<sub>2</sub>Ti(C≡CSiMe<sub>3</sub>)<sub>2</sub>]Pd(PPh<sub>3</sub>), 123

The crystal structure of [Mo(NCS)(CO)<sub>2</sub>(η<sup>3</sup>-C<sub>3</sub>H<sub>5</sub>)(NCMe<sub>2</sub>)]·MeCN and the reactions of {Mo(CO)<sub>2</sub>(η<sup>3</sup>-C<sub>3</sub>H<sub>5</sub>)<sup>+</sup>} containing species with symmetric alkynes, 176

## Solvent effects

Specific solvent effect on R<sub>2</sub>ZrCl<sub>2</sub> (R = butyl, ethyl) reactivity, a density functional study, 268

## Steric isomerization

Steric isomerization of alkyne–dialkyltungsten complexes with a chelating diaryloxo ligand: crystal structures of C<sub>s</sub>- and C<sub>1</sub>-W(η<sup>2</sup>-RC≡CR)[2,2'-S(4-Me-6-R'C<sub>6</sub>H<sub>2</sub>O)<sub>2</sub>](CH<sub>2</sub>SiMe<sub>3</sub>)<sub>2</sub>, 234

## Structures

Cyclic thioether and acyclic thioether–thiolate complexes of pentamethylcyclopentadienyl ruthenium(II, III), 161

Preparation and organic transformation reactions of CpFe(CO)<sub>2</sub>-SCO-3-C<sub>6</sub>H<sub>4</sub>-SO<sub>2</sub>Cl, 228

## Styrene

New complexes of palladium(II) with chelating heterocyclic nitrogen ligands. Characterization and catalytic activity in CO–styrene copolymerization. Crystal structure of [Pd(bipy)<sub>2</sub>][BAR'<sub>4</sub>], Ar' = [3,5-(CF<sub>3</sub>)<sub>2</sub>C<sub>6</sub>H<sub>3</sub>], 77

## Sulfonates

Preparation and organic transformation reactions of CpFe(CO)<sub>2</sub>-SCO-3-C<sub>6</sub>H<sub>4</sub>-SO<sub>2</sub>Cl, 228

## Supramolecular self-assembly

Crystal and molecular structure of (α-naphthyl)TeI<sub>3</sub>, 306

## Synthesis

Synthesis and structure of 1,2,4-triazol-2-ium-5-ylidene complexes of Hg(II), Pd(II), Ni(II), Ni(0), Rh(I) and Ir(I), 70

## Tellurium

Syntheses and structures of (R<sub>2</sub>Bi)<sub>2</sub>E (E = S, Te) and *cyclo*-(RSbSe)<sub>2</sub>[W(CO)<sub>5</sub>]<sub>2</sub> [R = CH(SiMe<sub>3</sub>)<sub>2</sub>], 130

## Tertiary phosphines

One-pot palladium-catalyzed phosphination of aryl iodides with Ph<sub>2</sub>PSnR<sub>3</sub>, 223

## Tetrahedron

Synthesis, reaction and enantiomeric resolution of chiral clusters containing SRuNiM (M = Mo, W) core, 201

## Tetraorganodistannoxanes

Synthesis, spectroscopic and structural aspects of some tetraorganodistannoxanes with internally functionalized oxime. 2. Crystal and molecular structure of {[Me<sub>2</sub>Sn(ON=C(Me)C<sub>5</sub>H<sub>4</sub>N)<sub>2</sub>O]<sub>2</sub>·2[2-NC<sub>5</sub>H<sub>4</sub>(Me)C=NOH]}, 66

## Tetraphenylborate

Nitrile exchange reactions of [WI(CO)(NCMe){P(O<sup>i</sup>Pr)<sub>3</sub>]<sub>2</sub>(η<sup>2</sup>-MeC<sub>2</sub>Me)][BPh<sub>4</sub>]; X-ray crystal structures of [WI(CO)(L){P(O<sup>i</sup>Pr)<sub>3</sub>]<sub>2</sub>(η<sup>2</sup>-MeC<sub>2</sub>Me)][BPh<sub>4</sub>] {L = NC<sup>i</sup>Pr, NC<sup>i</sup>Bu, NCCH<sub>2</sub>Ph, 1,2-C<sub>6</sub>H<sub>4</sub>(NCCH<sub>2</sub>)<sub>2</sub>}, 37

## 3-Thiapentane-1,5-dithiolate

Cyclic thioether and acyclic thioether–thiolate complexes of pentamethylcyclopentadienyl ruthenium(II, III), 161

## Thiocarboxylate

Preparation and organic transformation reactions of CpFe(CO)<sub>2</sub>-SCO-3-C<sub>6</sub>H<sub>4</sub>-SO<sub>2</sub>Cl, 228

## Thiolate–thioether

Cyclic thioether and acyclic thioether–thiolate complexes of pentamethylcyclopentadienyl ruthenium(II, III), 161

## Tin

Alkylderivatives of Stanna-*closa*-dodecaborate, 116

## Titanium

Monomere Kupfer(I)-Alkyle mit β-Wasserstoffatomen und Kupfer(I)-Aryle mit kondensierten Aromaten; die Festkörperstruktur von [(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>SiMe<sub>3</sub>)<sub>2</sub>Ti(C≡CSiMe<sub>3</sub>)<sub>2</sub>]Cu<sup>n</sup>C<sub>4</sub>H<sub>9</sub>, 142

Synthesis and electrochemical behaviour of σ,η<sup>2</sup>-acetylide-bridged early–late complexes; The solid-state structure of [(η<sup>5</sup>-C<sub>5</sub>H<sub>4</sub>SiMe<sub>3</sub>)<sub>2</sub>Ti(C≡CSiMe<sub>3</sub>)<sub>2</sub>]Pd(PPh<sub>3</sub>), 123

## Titanocene

Stabilisierung niedervalenter Ni(CO)-Bausteine durch [Ti](C≡CR)<sub>2</sub>; Reaktionsverhalten von {[Ti](C≡CR)<sub>2</sub>}Ni(CO) gegenüber Triphenylphosphan und Phosphiten, 150

## 1,2,4-Triazol-2-ium-5-ylidene complexes

Synthesis and structure of 1,2,4-triazol-2-ium-5-ylidene complexes of Hg(II), Pd(II), Ni(II), Ni(0), Rh(I) and Ir(I), 70

## (2,4,6-Triisopropylphenyl) methyl ketone

The factors controlling the reaction of (2,4,6-triisopropylphenyl) methyl ketone with Ph<sub>3</sub>Al and structure of [Ph<sub>2</sub>AlO(2,4,6-tri-<sup>i</sup>Pr-C<sub>6</sub>H<sub>2</sub>)C=CH<sub>2</sub>]<sub>2</sub>, 136

## Tri-phenylaluminium

The factors controlling the reaction of (2,4,6-triisopropylphenyl) methyl ketone with Ph<sub>3</sub>Al and structure of [Ph<sub>2</sub>AlO(2,4,6-tri-<sup>i</sup>Pr-C<sub>6</sub>H<sub>2</sub>)C=CH<sub>2</sub>]<sub>2</sub>, 136

## Trithiocarbonate

Reactions of cyclomanganated complexes with carbon disulfide: routes to η<sup>2</sup>-aryldithiocarboxylate-Mn(CO)<sub>4</sub> complexes and to the trithiocarbonate complex (μ<sub>3</sub>-CS<sub>3</sub>)<sub>2</sub>Mn<sub>4</sub>(CO)<sub>16</sub>, 288

## Tungsten

Gas-phase electronic absorption spectra of metallocene dihydrides M(η<sup>5</sup>-C<sub>5</sub>H<sub>5</sub>)<sub>2</sub>H<sub>2</sub> (M = Mo, W), 106

## Tungsten complex

Syntheses and structures of (R<sub>2</sub>Bi)<sub>2</sub>E (E = S, Te) and *cyclo*-(RSbSe)<sub>2</sub>[W(CO)<sub>5</sub>]<sub>2</sub> [R = CH(SiMe<sub>3</sub>)<sub>2</sub>], 130

## Tungsten(II)

Nitrile exchange reactions of  $[\text{W}(\text{CO})(\text{NCMe})\{\text{P}(\text{O}^i\text{Pr})_3\}_2(\eta^2\text{-MeC}_2\text{Me})][\text{BPh}_4]$ ; X-ray crystal structures of  $[\text{W}(\text{CO})(\text{L})\{\text{P}(\text{O}^i\text{Pr})_3\}_2(\eta^2\text{-MeC}_2\text{Me})][\text{BPh}_4]$  {L = NC<sup>i</sup>Pr, NC<sup>t</sup>Bu, NCCH<sub>2</sub>Ph, 1,2-C<sub>6</sub>H<sub>4</sub>(NCCH<sub>2</sub>)<sub>2</sub>}, 37

 $\pi$ -Tweezer

Synthesis and electrochemical behaviour of  $\sigma,\eta^2$ -acetylide-bridged early-late complexes; The solid-state structure of  $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2]\text{Pd}(\text{PPh}_3)$ , 123

## Vanadium(II)

Lithiation of ferrocenylamines and vanadium dinitrogen chemistry, 245

## X-ray

New complexes of palladium(II) with chelating heterocyclic nitrogen ligands. Characterization and catalytic activity in CO-styrene copolymerization. Crystal structure of  $[\text{Pd}(\text{bipy})_2][\text{BAr}'_4]_2$ , Ar' = [3,5-(CF<sub>3</sub>)<sub>2</sub>C<sub>6</sub>H<sub>3</sub>], 77

Stabilisierung niedervalenter Ni(CO)-Bausteine durch  $[\text{Ti}](\text{C}\equiv\text{CR})_2$ ; Reaktionsverhalten von  $\{[\text{Ti}](\text{C}\equiv\text{CR})_2\}\text{Ni}(\text{CO})$  gegenüber Triphenylphosphan und Phosphiten, 150

## X-ray crystallography

Preparation and structural characterization of six new diorgano-

tin(IV) complexes of the R<sub>2</sub>Sn(SaleanH<sub>2</sub>) and R<sub>2</sub>Sn(SalceanH<sub>2</sub>) type (R = Me, <sup>n</sup>Bu, Ph), 188

Stepwise assembly and structural characterization of oligonuclear ferrocene aggregates with boron-nitrogen backbone, 94

## X-ray diffraction study

The factors controlling the reaction of (2,4,6-triisopropylphenyl) methyl ketone with Ph<sub>3</sub>Al and structure of  $[\text{Ph}_2\text{AlO}(2,4,6\text{-tri-}^i\text{Pr-C}_6\text{H}_2)\text{C}=\text{CH}_2]_2$ , 136

## X-ray structure

Monomere Kupfer(I)-Alkyle mit  $\beta$ -Wasserstoffatomen und Kupfer(I)-Aryle mit kondensierten Aromaten; die Festkörperstruktur von  $[(\eta^5\text{-C}_5\text{H}_4\text{SiMe}_3)_2\text{Ti}(\text{C}\equiv\text{CSiMe}_3)_2]\text{Cu}^n\text{C}_4\text{H}_9$ , 142

Syntheses and structures of (R<sub>2</sub>Bi)<sub>2</sub>E (E = S, Te) and *cyclo*-(RSbSe)<sub>2</sub>[W(CO)<sub>5</sub>]<sub>2</sub> [R = CH(SiMe<sub>3</sub>)<sub>2</sub>], 130

## X-ray structures

Lithiation of ferrocenylamines and vanadium dinitrogen chemistry, 245

## Zirconium

Specific solvent effect on R<sub>2</sub>ZrCl<sub>2</sub> (R = butyl, ethyl) reactivity, a density functional study, 268

## Zwitterions

Alkylderivatives of Stanna-*closo*-dodecaborate, 116