



Contents of Volume 664

Communication

Polyether phosphine oxide induced phase separable homogeneous catalysis for hydroformylation of higher olefins

L. Xiaozhong, L. Hongmei, K. Fanzhi 1

Regular Papers

Indenyl-amido titanium and zirconium dimethyl complexes: improved synthesis and use in propylene polymerization

L. Resconi, I. Camurati, C. Grandini, M. Rinaldi, N. Mascellani, O. Traverso 5

Reactivity of *cyclo*-(PhX)₆ (X = As, P) towards [M₃L₂(CO)₁₀] (M = Ru, L = CO or NCMe; M = Fe, L = CO)

R.M. De Silva, M.J. Mays, G.A. Solan 27

Nitrile exchange reactions of [WI(CO)(NCMe){P(O*i*Pr)₃}₂(η²-MeC₂Me)][BPh₄]; X-ray crystal structures of [WI(CO)(L){P(O*i*Pr)₃}₂(η²-MeC₂Me)][BPh₄]} {L = NC*i*Pr, NC*t*Bu, NCCH₂Ph, 1,2-C₆H₄(NCCH₂)₂}

P.K. Baker, M.G.B. Drew, D.S. Moore 37

Synthesis and reactions of the cationic seven-coordinate complexes, [MoI(CO)₃(NCR){Ph₂P(CH₂)_nPPh₂}][BF₄] (R = Me, n = 1, 2 and 3; R = Et, n = 1). Crystal structures of [MoI₂(CO)₂(PPh₃)₂{Ph₂P(CH₂)PPh₂}], [MoI(CO)₂{Ph₂P(CH₂)PPh₂}₂][BF₄]·CH₂Cl₂, [MoCl₃O{Ph₂PO(CH₂)OPPh₂}] and [Mo₃(μ₃-I)₂{μ₂-Ph₂P(CH₂)PPh₂}₃]I

P.K. Baker, M.G.B. Drew, D.S. Moore 45

Highly selective synthesis of biphenyl by the Pd(OAc)₂/HPA/O₂/AcOH catalyst system

M. Okamoto, M. Watanabe, T. Yamaji 59

Synthesis, spectroscopic and structural aspects of some tetraorganodistannoxanes with internally functionalized oxime. 2. Crystal and molecular structure of [{Me₂Sn(ON=C(Me)C₅H₄N)₂O}]₂·2[2-NC₅H₄(Me)C=NOH]

V. Sharma, R.K. Sharma, R. Bohra, V.K. Jain, J.E. Drake, M.E. Light, M.B. Hursthouse 66

Synthesis and structure of 1,2,4-triazol-2-ium-5-ylidene complexes of Hg(II), Pd(II), Ni(II), Ni(0), Rh(I) and Ir(I)

C. Buron, L. Stelzig, O. Guerret, H. Gornitzka, V. Romanenko, G. Bertrand 70

New complexes of palladium(II) with chelating heterocyclic nitrogen ligands. Characterization and catalytic activity in CO-styrene copolymerization. Crystal structure of [Pd(bipy)₂][BAr'₄]₂, Ar' = [3,5-(CF₃)₂C₆H₃]

S. Stoccoro, G. Alesso, M.A. Cinello, G. Minghetti, A. Zucca, A. Bastero, C. Claver, M. Manassero 77

Osmacycles containing the {(Ph₂PE)₂N}⁻ ligand (E = S or Se): synthesis and crystal structures

J. Parr, M.B. Smith, M.R.J. Elsegood 85

Stepwise assembly and structural characterization of oligonuclear ferrocene aggregates with boron–nitrogen backbone

K. Ma, H.-W. Lerner, S. Scholz, J.W. Bats, M. Bolte, M. Wagner 94

Gas-phase electronic absorption spectra of metallocene dihydrides M(η⁵-C₅H₅)₂H₂ (M = Mo, W)

S.Y. Ketkov, V.V. Kutyreva, A.M. Ob'edkov, G.A. Domrachev 106

Syntheses of polyaluminium compounds by hydroalumination reactions, chelating Lewis acids possessing two and four coordinatively unsaturated aluminium atoms

W. Uhl, M. Matar 110

Alkylderivatives of Stanna-*clos*-dodecaborate

T. Marx, B. Ronig, H. Schulze, I. Pantenburg, L. Wesemann 116

Synthesis and electrochemical behaviour of σ,η²-acetylide-bridged early-late complexes: The solid-state structure of [(η⁵-C₅H₄SiMe₃)₂Ti(C≡CSiMe₃)₂]Pd(PPh₃)

S. Back, T. Stein, J. Kralik, C. Weber, G. Rheinwald, L. Zsolnai, G. Huttner, H. Lang 123

Syntheses and structures of $(R_2Bi)_2E$ ($E = S, Te$) and <i>cyclo</i> -(RSbSe) ₂ [W(CO) ₅] ₂ [$R = CH(SiMe_3)_2$] H.J. Breunig, I. Ghesner, E. Lork	130
The factors controlling the reaction of (2,4,6-triisopropylphenyl) methyl ketone with Ph ₃ Al and structure of [Ph ₂ AlO(2,4,6-tri- ⁱ Pr-C ₆ H ₂)C=CH ₂] A.R. Kunicki, A. Chojecki, J. Zachara, M. Gliński	136
Monomere Kupfer(I)-Alkyle mit β -Wasserstoffatomen und Kupfer(I)-Aryle mit kondensierten Aromaten; die Festkörperstruktur von $[(\eta^5-C_5H_4SiMe_3)_2Ti(C\equiv CSiMe_3)_2]Cu^nC_4H_9$ T. Stein, H. Lang	142
Stabilisierung niedervalenter Ni(CO)-Bausteine durch [Ti](C≡CR) ₂ ; Reaktionsverhalten von {[Ti](C≡CR) ₂ }Ni(CO) gegenüber Triphenylphosphitan und Phosphiten H. Lang, E. Meichel, T. Stein, C. Weber, J. Kralik, G. Rheinwald, H. Pritzkow	150
Cyclic thioether and acyclic thioether–thiolate complexes of pentamethylcyclopentadienyl ruthenium(II, III) L.Y. Goh, M.E. Teo, S.B. Khoo, W.K. Leong, J.J. Vittal	161
Oxidative insertion into the N–H bond of carbazole, indole and pyrrole with zerovalent metals C. López, G. Barón, A. Arévalo, M.A. Muñoz-Hernández, J.J. García	170
The crystal structure of [Mo(NCS)(CO) ₂ (η^3 -C ₃ H ₅)(NCMe) ₂]·MeCN and the reactions of {Mo(CO) ₂ (η^3 -C ₃ H ₅) ⁺ } containing species with symmetric alkynes J.W. Goodyear, C.W. Hemingway, R.W. Harrington, M.R. Wiseman, B.J. Brisdon	176
Sodium and potassium derivatives of diphenyl(trimethylsilyl)methane M.S. Hill, P.B. Hitchcock	182
Preparation and structural characterization of six new diorganotin(IV) complexes of the R ₂ Sn(SaleanH ₂) and R ₂ Sn(SalceanH ₂) type (R = Me, ⁿ Bu, Ph) R. García-Zarracino, J. Ramos-Quiñones, H. Höpfl	188
Synthesis, reaction and enantiomeric resolution of chiral clusters containing SRuNiM (M = Mo, W) core Z.-G. Bian, W.-Q. Zhang, H.-X. Guan, Y.-Q. Yin, Q.-S. Li, J. Sun	201
Oxorhenium(V) ‘3+2’ mixed-ligand complexes carrying the SNO/SN donor and C ₆₀ S. Ballot, N. Noiret	208
Hemilability of the primary amine–metal bond in polyamine-(Group 6) metal carbonyl complexes A.-C. Hervé, J.-J. Yaouanc, J.-C. Clément, H. des Abbayes, L. Toupet	214
One-pot palladium-catalyzed phosphination of aryl iodides with Ph ₂ PSnR ₃ S.E. Martín, M. Bonaterra, R.A. Rossi	223
Preparation and organic transformation reactions of CpFe(CO) ₂ SCO–3-C ₆ H ₄ –SO ₂ Cl M. El-khateeb, A. Younes	228
Steric isomerization of alkyne–dialkyltungsten complexes with a chelating diaryloxo ligand: crystal structures of C _s - and C ₁ -W(η^2 -RC≡CR)[2,2'-S(4-Me-6-R'C ₆ H ₂ O ₂) ₂](CH ₂ SiMe ₃) ₂ Y. Takashima, Y. Nakayama, H. Yasuda, A. Nakamura, A. Harada	234
Lithiation of ferrocenylamines and vanadium dinitrogen chemistry P.B. Hitchcock, G.J. Leigh, M. Togrou	245
Arylimido niobium(V) complexes: mononuclear and dendritic derivatives J.M. Benito, E. de Jesús, F. Javier de la Mata, J.C. Flores, R. Gómez, P. Gómez-Sal	258
Specific solvent effect on R ₂ ZrCl ₂ (R = butyl, ethyl) reactivity, a density functional study E. Derat, J. Bouquant, P. Bertus, J. Szymoniak, S. Humbel	268
Synthesis and reactivity of organochalogen ester substituted η^3 -butadienyl complexes of Mo(II): crystal structure of [MoCl(CO) ₂ (η^3 -CH ₂ C(COSePh)C=CH ₂)(1,10-phenanthroline)]·0.5 CH ₂ Cl ₂ A.G.W. Hodson, R.K. Thind, M. McPartlin	277
Reactions of cyclomanganated complexes with carbon disulfide: routes to η^2 -aryldithiocarboxylate-Mn(CO) ₄ complexes and to the trithiocarbonate complex (μ_3 -CS ₃) ₂ Mn ₄ (CO) ₁₆ W.J. Mace, L. Main, B.K. Nicholson, M. Hagyard	288

Facile syntheses of new multidentate (phosphino)amines: X-ray structure of 1,4-{(OC) ₄ Mo(Ph ₂ P) ₂ NCH ₂ } ₂ C ₆ H ₄	321
K.G. Gaw, M.B. Smith, J.W. Steed	294
The reaction of samarium(II) with manganese carbonyl: unexpected conversion of CO to formate. X-ray crystal structures of [Sm(Tp ^{Me²}) ₂]Mn(CO) ₅ and {[Sm(Tp ^{Me²}) ₂] ₂ (μ-HCO ₂)}Mn(CO) ₅ (Tp ^{Me²} = HB(3,5-dimethylpyrazolyl)	298
A.C. Hillier, A. Sella, M.R.J. Elsegood	298
Note	
Crystal and molecular structure of (α -naphthyl)TeI ₃	
E.S. Lang, G. Manzoni de Oliveira, E.T. Silveira, R.A. Burrow, E.M. Vázquez-López	306
Author Index of Volume 664	311
Subject Index of Volume 664	313
Contents of Volume 664	319