

Contents of Volume 669

Organotransition metal modified sugars: Part 23. Synthesis of vinylcarbene chromium complexes containing a C-monosaccharide ligand E. Janes, K.H. Dötz	1
Addition of diazoalkanes to alkyne(pentacarbonyl)–chromium and –tungsten — formation of 3 <i>H</i> -pyrazole complexes M.M. Abd-Elzaher, B. Weibert, H. Fischer	6
Borohydride reduction of a rhenium-bound acetonitrile: an example of a chelating iminoborane ligand at a low valent metal center A.C. Hillier, T. Fox, H.W. Schmalle, H. Berke	14
Phosphan-stabilisierte Silber(I)–Carboxylate: Synthese und Verwendung in der Silber-CVD H. Schmidt, Y. Shen, M. Leschke, T. Haase, K. Kohse-Höinghaus, H. Lang	25
Synthesis and X-ray structure of palladium dichloride complexed with THF and 2,6,7-trioxa-3,5,8-tris(trichloromethyl)-1,4-diphosphabicyclo[2.2.2]octane Y. Kim, J.G. Verkade	32
Synthesis and characterization of cyclopentadienyl thiolato complexes of magnesium A. Xia, M.J. Heeg, C.H. Winter	37
Reproducible high-yield syntheses of [Ru ₃ (CO) ₁₂], [H ₄ Ru ₄ (CO) ₁₂], and [Ru ₆ C(CO) ₁₆] ²⁻ by a convenient two-step methodology involving controlled reduction in ethylene glycol of RuCl ₃ · <i>n</i> H ₂ O E. Lucenti, E. Cariati, C. Dragonetti, D. Roberto	44
Conformationally rigid diphosphine arene–ruthenium(II) complexes as catalysts for transfer hydrogenation of ketones K.Y. Ghebreyessus, J.H. Nelson	48
Tetranuclear heterodimetallc metallamacrocycles with M–Sn(IV) (M = Mo or W) bonds. Crystal structures of { <i>p</i> -[(CO) ₃ MoC ₅ H ₄ C(O)] ₂ C ₆ H ₄ }{(Ph ₂ Sn) ₂ S} and { <i>p</i> -[(CO) ₃ MoC ₅ H ₄ C(O)] ₂ C ₆ H ₄ }{(Ph ₂ Sn) ₂ CH ₂ } L.-F. Tang, J.-F. Chai, S.-B. Zhao, J.-T. Wang	57
Influence of alkoxy ligands on the Cp–Al bonding mode in [Cp ₂ Al–μ-OR] ₂ from X-ray crystallographic and ²⁷ Al-NMR spectroscopic solution studies S. Szumacher, I. Madura, J. Zachara, A.R. Kunicki, A. Cebulski, L. Jerzykiewicz	64
Hydroboration of alkyne-1-yl(methyl)silanes bearing functional substituents at silicon B. Wrackmeyer, H.E. Maisel, W. Milius, M.H. Bhatti, S. Ali	72
Late transition metal complexes derived from diphosphazane monosulfide ligands: X-ray crystal structures of [Ru ₃ (μ-CO)(CO) ₇ (μ ₃ -S){Ph ₂ PN((<i>S</i>)-*CHMePh)PPh ₂ -κ ² P, <i>P</i> }] and [Rh(CO)Cl{Ph ₂ PN((<i>S</i>)-*CHMePh)P(S)Ph ₂ }-κ ² P, <i>S</i>]}: Part 16. Organometallic chemistry of diphosphazanes K. Raghuraman, S.S. Krishnamurthy, M. Nethaji	79
The lithiation of arylamines and the preparation of cyclopentadienyltitanium(IV) arylaminato complexes A.G. Avent, P.B. Hitchcock, G.J. Leigh, M. Togrou	87
Synthesis of titanium(IV) (cyclopentadienylalkyl)aminato complexes P.B. Hitchcock, G.J. Leigh, M. Togrou	101
Anomalous reaction of an aryl silane with Co ₂ (CO) ₈ ; characterisation of Me ₂ NC ₆ H ₄ Si[Co(CO) ₄][OCCO ₃ (CO) ₉] ₂ G.J. Harfoot, B.K. Nicholson	106
Comparative study of structure–activity relationship of di- and tri-organotin(IV) derivatives of amino acid and peptides M. Nath, S. Pokharia, G. Eng, X. Song, A. Kumar	109

Syntheses of a series of trinuclear $M\text{Ir}_2$ or pentanuclear $M\text{Ir}_4$ bimetallic bis(selenido) and selenido–sulfido clusters ($M = \text{Pd}, \text{Pt}, \text{Fe}, \text{Co}$) from diiridium μ -bis(hydroselenido) and μ -hydroselenido–hydrosulfido complexes $\{[(\eta^5\text{-C}_5\text{Me}_5)\text{IrCl}]_2(\mu\text{-SeH})(\mu\text{-EH})\}$ ($E = \text{Se}, \text{S}$) S. Nagao, H. Seino, M. Hidai, Y. Mizobe	124
Synthesis and reactivity of a new Fe(II) 5-(4-pyridyl)-tetrazolate complex and X-ray structure of its doubly protonated derivative. A. Palazzi, S. Stagni, S. Selva, M. Monari	135
Preparation and oxidation of polarized Au(III) complexes having both the <i>C</i> -deprotonated-2-phenylpyridine (ppy) and a sulfur-rich dithiolate ligand and X-ray crystal structure of $[\text{Au}(\eta^2\text{-C}, N\text{-ppy})(\eta^2\text{-S}, S\text{-C}_8\text{H}_4\text{S}_8)] \cdot 0.5\text{DMF}$ K. Kubo, M. Nakano, H. Tamura, G.-e. Matsubayashi, M. Nakamoto	141
Secondary bonding in <i>para</i> -substituted diphenyltellurium dichlorides ($p\text{-XC}_6\text{H}_4)_2\text{TeCl}_2$ ($X = \text{H}, \text{Me}, \text{MeO}$) probed by ^{125}Te MAS NMR spectroscopy. Crystal and molecular structure of ($p\text{-MeC}_6\text{H}_4)_2\text{TeCl}_2$ J. Beckmann, D. Dakternieks, A. Duthie, N.A. Smith	149
Modelling nucleophilic substitution at silicon using solution ^{19}F -NMR chemical shift, $^1J_{\text{Si-F}}$ and $^2J_{\text{C-F}}$ coupling constant data of pentacoordinate silicon compounds. Correlation with other magnetic nuclei and X-ray structures A.R. Bassindale, Y.I. Baukov, M. Borbaruah, S.J. Glynn, V.V. Negrebetsky, D.J. Parker, P.G. Taylor, R. Turtle	154
Synthesis, characterisation and study of the first luminescent platinum(II) compound with a $[\text{C}, \text{N}, \text{S}]^-$ terdentate ligand. X-ray crystal structure of $[\text{Pt}\{\text{C}_6\text{H}_4\text{-CH=N-(C}_6\text{H}_4\text{-2-SMe)\Cl}]$ A. Caubet, C. López, X. Solans, M. Font-Bardía	164
Hydridorhodium(I) complexes with amphiphilic polyether phosphines: NMR study and biphasic hydroformylation of 1-octene A. Solsona, J. Suades, R. Mathieu	172
Synthesis and X-ray structures of some metal carbonyl complexes containing dipyridylformamidine ligands H.-C. Liang, Y.-Y. Wu, F.-C. Chang, P.-Y. Yang, J.-D. Chen, J.-C. Wang	182
Synthesis, structure, and reactivity of novel iron(II) complexes with a five-membered chelate ligand $\kappa^2(\text{Si}, N)\text{-SiMe}_2\text{O}(2\text{-C}_5\text{H}_4\text{N})$ T. Sato, M. Okazaki, H. Tobita, H. Ogino	189
The first example of a structurally characterized octahedral hydrotris(pyrazolyl)borate iron methyl complex O. Graziani, L. Toupet, J.-R. Hamon, M. Tilset	200
Book reviews	207
Erratum	210
Author Index of Volume 669	211
Subject Index of Volume 669	213
Contents Index of Volume 669	217