



ELSEVIER

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SCIENCE @ DIRECT®

Journal of Organometallic Chemistry 676 (2003) 95–96

Journal  
of Organo  
metallic  
Chemistry[www.elsevier.com/locate/jorgchem](http://www.elsevier.com/locate/jorgchem)

## Subject Index of Volume 676

 $\sigma$ -Acceptor

The polar effect on the regiochemistry of nucleophilic substitution of trifluoromethylated  $\pi$ -allylpalladium complex, 43

## Alkyne

Stereocontrolled synthesis of (*E*)- and (*Z*)-poly(*p*-phenylenevinylene)s via ruthenium-catalyzed hydrosilylation of *p*-diethynylbenzene, 49

## Allyl complexes

Diastereomerism in palladium(II) allyl complexes of P,P-, P,S- and S,S-donor ligands,  $\text{Ph}_2\text{P}(\text{E})\text{N}(\text{R})\text{P}(\text{E}')\text{Ph}_2$  [R = CHMe<sub>2</sub> or (*S*)-\*CHMePh; E = E' = lone pair or S]: solution behaviour, X-ray crystal structure and catalytic allylic alkylation reactions, 22

## Allylic alkylation

Diastereomerism in palladium(II) allyl complexes of P,P-, P,S- and S,S-donor ligands,  $\text{Ph}_2\text{P}(\text{E})\text{N}(\text{R})\text{P}(\text{E}')\text{Ph}_2$  [R = CHMe<sub>2</sub> or (*S*)-\*CHMePh; E = E' = lone pair or S]: solution behaviour, X-ray crystal structure and catalytic allylic alkylation reactions, 22

 $\pi$ -Allylpalladium complexes

The polar effect on the regiochemistry of nucleophilic substitution of trifluoromethylated  $\pi$ -allylpalladium complex, 43

*ansa*-Calcocene

Calcium-mediated fulvene couplings. 3. Reductive coupling of guaiazulene with activated calcium to give a mixture of 8,6' and 8,8' (diguaiazulenide)calcium isomers. Thermal rearrangement of the 8,6' isomer to the 8,8' isomer and X-ray crystal structure of the 8,8' isomer, 73

## Bent metallocenes

Synthesis and structural features of [(C<sub>5</sub>H<sub>4</sub>-CMe<sub>2</sub>NMe<sub>2</sub>)<sub>2</sub>TiCl], a bis(cyclopentadienyl)Ti(III) complex with internal dimethylamino coordination, 1

## Bridged carbyne

Synthesis of bridging carbyne butterfly Fe/Se cluster complexes via reaction of complex anions  $[(\mu\text{-RSe})(\mu\text{-CO})\text{Fe}_2(\text{CO})_6]^-$ : Crystal structure of ( $\mu\text{-EtOC})(\mu\text{-}p\text{-MeC}_6\text{H}_4\text{Se})\text{Fe}_2(\text{CO})_6$ , 80

*cis*-Dihydride complexes

Neutron diffraction study of the highly distorted octahedral complex  $\text{FeH}_2(\text{CO})_2[\text{P}(\text{OPh})_3]_2$ , 85

## Cobalt

Study of metal exchange reactions in cobalt containing clusters, 55

## CO-bridged anions

Synthesis of bridging carbyne butterfly Fe/Se cluster complexes via reaction of complex anions  $[(\mu\text{-RSe})(\mu\text{-CO})\text{Fe}_2(\text{CO})_6]^-$ : Crystal structure of ( $\mu\text{-EtOC})(\mu\text{-}p\text{-MeC}_6\text{H}_4\text{Se})\text{Fe}_2(\text{CO})_6$ , 80

## Crystal structures

New stable cyclopentadienyl iron(II) complexes containing 3,5-dimethylpyrazolyl phosphine: X-ray structure of  $[\text{FeCp}(\text{CO})_2\text{-PPh}_2(\text{Me}_2\text{Pz})]\text{BF}_4$ , 38

Synthesis and characterization of 3- and 4-phenylgermatranes: X-ray crystal structures of  $\text{N}(\text{CH}_2\text{CH}_2\text{O})_2(\text{CH}_2\text{CHPhO})\text{GeZ}$  (Z = F, OSiMe<sub>3</sub>, C≡CPh) and  $\text{N}(\text{CH}_2\text{CH}_2\text{O})_2(\text{CHPh-CH}_2\text{O})\text{GeOH}$ , 8

## Electrochemistry

Electrochemistry of ferrocenylphosphines  $\text{FcCH}_2\text{PR}_2$  (Fc = ( $\eta^5\text{-C}_5\text{H}_5$ )Fe( $\eta^5\text{-C}_5\text{H}_4$ ); R = Ph, CH<sub>2</sub>OH and CH<sub>2</sub>CH<sub>2</sub>CN), and some phosphine oxide, phosphine sulfide, phosphonium and metal complex derivatives, 62

## Ferrocenes

Synthesis and structural features of [(C<sub>5</sub>H<sub>4</sub>-CMe<sub>2</sub>NMe<sub>2</sub>)<sub>2</sub>TiCl], a bis(cyclopentadienyl)Ti(III) complex with internal dimethylamino coordination, 1

## Ferrocenium reaction

Electrochemistry of ferrocenylphosphines  $\text{FcCH}_2\text{PR}_2$  (Fc = ( $\eta^5\text{-C}_5\text{H}_5$ )Fe( $\eta^5\text{-C}_5\text{H}_4$ ); R = Ph, CH<sub>2</sub>OH and CH<sub>2</sub>CH<sub>2</sub>CN), and some phosphine oxide, phosphine sulfide, phosphonium and metal complex derivatives, 62

## Ferrocenyl group

Electrochemistry of ferrocenylphosphines  $\text{FcCH}_2\text{PR}_2$  (Fc = ( $\eta^5\text{-C}_5\text{H}_5$ )Fe( $\eta^5\text{-C}_5\text{H}_4$ ); R = Ph, CH<sub>2</sub>OH and CH<sub>2</sub>CH<sub>2</sub>CN), and some phosphine oxide, phosphine sulfide, phosphonium and metal complex derivatives, 62

## Fulvene coupling

Calcium-mediated fulvene couplings. 3. Reductive coupling of guaiazulene with activated calcium to give a mixture of 8,6' and 8,8' (diguaiazulenide)calcium isomers. Thermal rearrangement of the 8,6' isomer to the 8,8' isomer and X-ray crystal structure of the 8,8' isomer, 73

## Functionalized cyclopentadienides

Synthesis and structural features of [(C<sub>5</sub>H<sub>4</sub>-CMe<sub>2</sub>NMe<sub>2</sub>)<sub>2</sub>TiCl], a bis(cyclopentadienyl)Ti(III) complex with internal dimethylamino coordination, 1

## Germatrane

Synthesis and characterization of 3- and 4-phenylgermatranes: X-ray crystal structures of  $\text{N}(\text{CH}_2\text{CH}_2\text{O})_2(\text{CH}_2\text{CHPhO})\text{GeZ}$  (Z = F, OSiMe<sub>3</sub>, C≡CPh) and  $\text{N}(\text{CH}_2\text{CH}_2\text{O})_2(\text{CHPh-CH}_2\text{O})\text{GeOH}$ , 8

## Guaiazulene

Calcium-mediated fulvene couplings. 3. Reductive coupling of guaiazulene with activated calcium to give a mixture of 8,6' and 8,8' (diguaiazulenide)calcium isomers. Thermal rearrangement of the 8,6' isomer to the 8,8' isomer and X-ray crystal structure of the 8,8' isomer, 73

## Hydrosilylation

Stereocontrolled synthesis of (*E*)- and (*Z*)-poly(*p*-phenylenevinylene)s via ruthenium-catalyzed hydrosilylation of *p*-diethynylbenzene, 49

## Intramolecular coordination

Synthesis and structural features of [(C<sub>5</sub>H<sub>4</sub>-CMe<sub>2</sub>NMe<sub>2</sub>)<sub>2</sub>TiCl], a bis(cyclopentadienyl)Ti(III) complex with internal dimethylamino coordination, 1

- Iron**  
 Synthesis of bridging carbyne butterfly Fe/Se cluster complexes via reaction of complex anions  $[(\mu\text{-RSe})(\mu\text{-CO})\text{Fe}_2(\text{CO})_6]^-$ : Crystal structure of  $(\mu\text{-EtOC})(\mu\text{-}p\text{-MeC}_6\text{H}_4\text{Se})\text{Fe}_2(\text{CO})_6$ , 80
- Iron complexes**  
 New stable cyclopentadienyl iron(II) complexes containing 3,5-dimethylpyrazolyl phosphine: X-ray structure of  $[\text{FeCp}(\text{CO})_2\text{-PPh}_2(\text{Me}_2\text{Pz})]\text{BF}_4$ , 38
- Iron hydride**  
 Neutron diffraction study of the highly distorted octahedral complex  $\text{FeH}_2(\text{CO})_2[\text{P}(\text{OPh})_3]_2$ , 85
- Metal exchange**  
 Study of metal exchange reactions in cobalt containing clusters, 55
- Molybdenum**  
 Study of metal exchange reactions in cobalt containing clusters, 55
- Neutron diffraction**  
 Neutron diffraction study of the highly distorted octahedral complex  $\text{FeH}_2(\text{CO})_2[\text{P}(\text{OPh})_3]_2$ , 85
- Nickel**  
 Nickel(II) and nickel(0) derivatives of bis(diphenylphosphino)amine:  $[\text{N}(\text{PPh}_2)_2]_2\text{Ni}$ ,  $(\text{Ph}_3\text{P})_2\text{Ni}[(\text{Ph}_2\text{P})_2\text{NH}]$ . Synthesis, characterization, and some properties, 89
- NMR spectroscopy**  
 Electrochemistry of ferrocenylphosphines  $\text{FcCH}_2\text{PR}_2$  ( $\text{Fc} = (\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\eta^5\text{-C}_5\text{H}_4)$ ;  $\text{R} = \text{Ph}$ ,  $\text{CH}_2\text{OH}$  and  $\text{CH}_2\text{CH}_2\text{CN}$ ), and some phosphine oxide, phosphine sulfide, phosphonium and metal complex derivatives, 62  
 Synthesis and characterization of 3- and 4-phenylgermatranes: X-ray crystal structures of  $\text{N}(\text{CH}_2\text{CH}_2\text{O})_2(\text{CH}_2\text{CHPhO})\text{GeZ}$  ( $\text{Z} = \text{F}$ ,  $\text{OSiMe}_3$ ,  $\text{C}\equiv\text{CPh}$ ) and  $\text{N}(\text{CH}_2\text{CH}_2\text{O})_2(\text{CHPhCH}_2\text{O})\text{GeOH}$ , 8
- Palladium**  
 Diastereomerism in palladium(II) allyl complexes of P,P-, P,S- and S,S-donor ligands,  $\text{Ph}_2\text{P}(\text{E})\text{N}(\text{R})\text{P}(\text{E}')\text{Ph}_2$  [ $\text{R} = \text{CHMe}_2$  or (*S*)-\*CHMePh;  $\text{E} = \text{E}' = \text{lone pair or S}$ ]; solution behaviour, X-ray crystal structure and catalytic allylic alkylation reactions, 22
- Phosphazanes**  
 Nickel(II) and nickel(0) derivatives of bis(diphenylphosphino)amine:  $[\text{N}(\text{PPh}_2)_2]_2\text{Ni}$ ,  $(\text{Ph}_3\text{P})_2\text{Ni}[(\text{Ph}_2\text{P})_2\text{NH}]$ . Synthesis, characterization, and some properties, 89
- Phosphine**  
 Electrochemistry of ferrocenylphosphines  $\text{FcCH}_2\text{PR}_2$  ( $\text{Fc} = (\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\eta^5\text{-C}_5\text{H}_4)$ ;  $\text{R} = \text{Ph}$ ,  $\text{CH}_2\text{OH}$  and  $\text{CH}_2\text{CH}_2\text{CN}$ ), and some phosphine oxide, phosphine sulfide, phosphonium and metal complex derivatives, 62
- Phosphinoamides**  
 Nickel(II) and nickel(0) derivatives of bis(diphenylphosphino)amine:  $[\text{N}(\text{PPh}_2)_2]_2\text{Ni}$ ,  $(\text{Ph}_3\text{P})_2\text{Ni}[(\text{Ph}_2\text{P})_2\text{NH}]$ . Synthesis, characterization, and some properties, 89
- P ligands**  
 Diastereomerism in palladium(II) allyl complexes of P,P-, P,S- and S,S-donor ligands,  $\text{Ph}_2\text{P}(\text{E})\text{N}(\text{R})\text{P}(\text{E}')\text{Ph}_2$  [ $\text{R} = \text{CHMe}_2$  or (*S*)-\*CHMePh;  $\text{E} = \text{E}' = \text{lone pair or S}$ ]; solution behaviour, X-ray crystal structure and catalytic allylic alkylation reactions, 22
- Polycondensation**  
 Stereocontrolled synthesis of (*E*)- and (*Z*)-poly(*p*-phenylenevinylene)s via ruthenium-catalyzed hydrosilylation of *p*-diethynylbenzene, 49
- Poly(phenylenevinylene)**  
 Stereocontrolled synthesis of (*E*)- and (*Z*)-poly(*p*-phenylenevinylene)s via ruthenium-catalyzed hydrosilylation of *p*-diethynylbenzene, 49
- Pyrazole complexes**  
 New stable cyclopentadienyl iron(II) complexes containing 3,5-dimethylpyrazolyl phosphine: X-ray structure of  $[\text{FeCp}(\text{CO})_2\text{-PPh}_2(\text{Me}_2\text{Pz})]\text{BF}_4$ , 38
- Regiochemistry**  
 The polar effect on the regiochemistry of nucleophilic substitution of trifluoromethylated  $\pi$ -allylpalladium complex, 43
- Selenium**  
 Synthesis of bridging carbyne butterfly Fe/Se cluster complexes via reaction of complex anions  $[(\mu\text{-RSe})(\mu\text{-CO})\text{Fe}_2(\text{CO})_6]^-$ : Crystal structure of  $(\mu\text{-EtOC})(\mu\text{-}p\text{-MeC}_6\text{H}_4\text{Se})\text{Fe}_2(\text{CO})_6$ , 80
- S ligands**  
 Diastereomerism in palladium(II) allyl complexes of P,P-, P,S- and S,S-donor ligands,  $\text{Ph}_2\text{P}(\text{E})\text{N}(\text{R})\text{P}(\text{E}')\text{Ph}_2$  [ $\text{R} = \text{CHMe}_2$  or (*S*)-\*CHMePh;  $\text{E} = \text{E}' = \text{lone pair or S}$ ]; solution behaviour, X-ray crystal structure and catalytic allylic alkylation reactions, 22
- Stereocontrolled synthesis**  
 Stereocontrolled synthesis of (*E*)- and (*Z*)-poly(*p*-phenylenevinylene)s via ruthenium-catalyzed hydrosilylation of *p*-diethynylbenzene, 49
- Stereodynamics by NMR spectroscopy**  
 Diastereomerism in palladium(II) allyl complexes of P,P-, P,S- and S,S-donor ligands,  $\text{Ph}_2\text{P}(\text{E})\text{N}(\text{R})\text{P}(\text{E}')\text{Ph}_2$  [ $\text{R} = \text{CHMe}_2$  or (*S*)-\*CHMePh;  $\text{E} = \text{E}' = \text{lone pair or S}$ ]; solution behaviour, X-ray crystal structure and catalytic allylic alkylation reactions, 22
- Tetrahedral cluster**  
 Study of metal exchange reactions in cobalt containing clusters, 55
- Transition metal**  
 Study of metal exchange reactions in cobalt containing clusters, 55
- Transition metals**  
 Nickel(II) and nickel(0) derivatives of bis(diphenylphosphino)amine:  $[\text{N}(\text{PPh}_2)_2]_2\text{Ni}$ ,  $(\text{Ph}_3\text{P})_2\text{Ni}[(\text{Ph}_2\text{P})_2\text{NH}]$ . Synthesis, characterization, and some properties, 89
- Trifluoromethyl group**  
 The polar effect on the regiochemistry of nucleophilic substitution of trifluoromethylated  $\pi$ -allylpalladium complex, 43
- X-ray diffraction**  
 Nickel(II) and nickel(0) derivatives of bis(diphenylphosphino)amine:  $[\text{N}(\text{PPh}_2)_2]_2\text{Ni}$ ,  $(\text{Ph}_3\text{P})_2\text{Ni}[(\text{Ph}_2\text{P})_2\text{NH}]$ . Synthesis, characterization, and some properties, 89
- X-ray structures**  
 Synthesis of bridging carbyne butterfly Fe/Se cluster complexes via reaction of complex anions  $[(\mu\text{-RSe})(\mu\text{-CO})\text{Fe}_2(\text{CO})_6]^-$ : Crystal structure of  $(\mu\text{-EtOC})(\mu\text{-}p\text{-MeC}_6\text{H}_4\text{Se})\text{Fe}_2(\text{CO})_6$ , 80