



## Contents of Volume 680

## Special Issue: Frontiers in Boron Chemistry Dedicated to Professor M. Frederick Hawthorne on his 75th Birthday

Preface . . . . .	1
<b>Reviews</b>	
Transition metal-catalyzed borylation of alkanes and arenes via C–H activation T. Ishiyama, N. Miyaura . . . . .	3
The use of organoboron chlorides and bromides in organic synthesis G.W. Kabalka, Z. Wu, Y. Ju . . . . .	12
The chemistry of the azanonaborane cluster $\text{RNH}_2\text{--B}_8\text{H}_{11}\text{NHR}$ D. Gabel, C. Bauer, M.E. El-Zaria, A.R. Genady, U. Dörfler . . . . .	23
Poly(cyclodiborazane)s F. Matsumoto, Y. Chujo . . . . .	27
The chemistry of [1]borametallocenophanes and related compounds H. Braunschweig, F.M. Breitling, E. Gullo, M. Kraft . . . . .	31
Regio- and stereoselective synthesis of boryl-substituted allylsilanes via transition metal-catalyzed silaboration M. Sugimoto, Y. Ito . . . . .	43
<b>Regular Papers</b>	
Small cobaltacborane clusters in synthesis. Peralkylation, perhalogenation, and macrocycle construction H. Yao, R.N. Grimes . . . . .	51
Synthesis and structural characterization of an 11-vertex <i>nido</i> -diphosphaborane, 7,9-Ph <sub>2</sub> - <i>nido</i> -7,9-P <sub>2</sub> B <sub>9</sub> H <sub>9</sub> D. Hong, P.J. Carroll, L.G. Sneddon . . . . .	61
Reactions of <i>nido</i> -1,2-(Cp <sup>*</sup> RuH) <sub>2</sub> B <sub>3</sub> H <sub>7</sub> with RC≡CR' (R, R' = H, Ph; Me, Me) to yield novel metallacarboranes H. Yan, A.M. Beatty, T.P. Fehlner . . . . .	66
Reactions of the dichloroboryl complex of osmium, Os(BCl <sub>2</sub> )Cl(CO)(PPh <sub>3</sub> ) <sub>2</sub> , with water, alcohols, and amines: Crystal structures of Os[B(OH) <sub>2</sub> ]Cl(CO)(PPh <sub>3</sub> ) <sub>2</sub> , Os[B(OEt) <sub>2</sub> ]Cl(CO)(PPh <sub>3</sub> ) <sub>2</sub> , and Os[BN(CH <sub>3</sub> )C <sub>6</sub> H <sub>4</sub> N(CH <sub>3</sub> ) <sub>2</sub> ]Cl(CO)(PPh <sub>3</sub> ) <sub>2</sub> G.R. Clark, G.J. Irvine, W.R. Roper, L.J. Wright . . . . .	81
A versatile rigid binucleating ligand for Rh <sub>2</sub> (μ-Cl) <sub>2</sub> moieties: its application as a catalyst in hydrogenation and cyclopropanation: Crystal structure of [Rh(C <sub>5</sub> Me <sub>5</sub> )Cl{7,8-μ-S(4'-C <sub>6</sub> H <sub>3</sub> (CH <sub>3</sub> )S-C <sub>2</sub> B <sub>9</sub> H <sub>10</sub> }][7,8-μ-S(4'-C <sub>6</sub> H <sub>3</sub> (CH <sub>3</sub> )S-C <sub>2</sub> B <sub>9</sub> H <sub>10}]·THF F. Teixidor, M.R. Cirera, C. Viñas, R. Kivekäs, R. Sillanpää, A. Demonceau . . . . .</sub>	89
Bromination of tri(isopropyl)boroxine and asymmetric synthesis of (2-cyano-3,3-dimethylcyclopropyl)boronic esters D.S. Matteson, D. Fernando . . . . .	100
[C <sub>2</sub> B <sub>10</sub> ]–[B <sub>12</sub> ] double cage boron compounds—a new approach to the synthesis of water-soluble boron-rich compounds for BNCT I.B. Sivaev, S. Sjöberg, V.I. Bregadze . . . . .	106
First agostic <i>closo</i> -metallacarboranes with η <sup>3</sup> -cyclooctenyl type ligand: synthesis and structural characterization of <i>closo</i> -3-[η <sup>3</sup> -( <i>endo</i> -1,5-dimethylcycloocten-1-yl)]-1,2-μ-(1',2'-xylylene)-3,1,2-IrC <sub>2</sub> B <sub>9</sub> H <sub>9</sub> and its isomerization to <i>closo</i> -3-[η <sup>3</sup> -( <i>exo</i> -1-methylene-5-methylcyclooctene-1-yl)]-1,2-μ-[η <sup>2</sup> -(1',2'-xylylene)]-3,1,2-IrC <sub>2</sub> B <sub>9</sub> H <sub>9</sub> A.V. Safronov, T.V. Zinevich, F.M. Dolgushin, E.V. Vorontsov, O.L. Tok, I.T. Chizhevsky . . . . .	111

Synthesis of adenosine containing carborane modification A.B. Olejniczak, A. Semenuk, M. Kwiatkowski, Z.J. Lesnikowski . . . . .	124
<i>exo</i> -closo-Rhodacarboranes: synthesis and characterisation of [{ <i>exo</i> -(R <sub>3</sub> P) <sub>2</sub> Rh}(closo-CB <sub>11</sub> H <sub>12</sub> )] [R <sub>3</sub> P = P(OMe) <sub>3</sub> , PCy <sub>3</sub> , 1/2dppe] A. Rifat, V.E. Laing, G. Kociok-Köhne, M.F. Mahon, G.D. Ruggiero, A.S. Weller . . . . .	127
Synthesis and reaction of secondary and primary diorganozinc reagents using a boron–zinc exchange reaction: A useful method for the stereo- and regioselective formation of new carbon–carbon bonds E. Hupe, M.I. Calaza, P. Knochel . . . . .	136
Catalyzed hydroboration of allyl sulfonamides M.G. Hamilton, C.E. Hughes, A.M. Irving, C.M. Vogels, S.A. Westcott . . . . .	143
Density functional theory studies on structural isomers and bonding of catecholborane adducts of Group 5 metallocene (Nb, Ta) hydride complexes D. Liu, K.-C. Lam, Z. Lin . . . . .	148
Big macrocyclic assemblies of carboranes (big MACs): synthesis and crystal structure of a macrocyclic assembly of four carboranes containing alternate <i>ortho</i> - and <i>meta</i> -carborane icosahedra linked by <i>para</i> -phenylene units M.A. Fox, J.A.K. Howard, J.A. Hugh MacBride, A. Mackinnon, K. Wade . . . . .	155
Dimesitylborane monomer-dimer equilibrium in solution, and the solid-state structure of the dimer by single crystal neutron and X-ray diffraction C.D. Entwistle, T.B. Marder, P.S. Smith, J.A.K. Howard, M.A. Fox, S.A. Mason . . . . .	165
Group 4 metallacarboranes of constrained geometries derived from B <sub>(cage)</sub> - and C <sub>(cage)</sub> -silylamido-substituted carborane ligands: a synthetic and structural investigation J. Wang, Y. Zhu, S. Li, C. Zheng, J.A. Maguire, N.S. Hosmane . . . . .	173
9-BBN activation. Synthesis, crystal structure and theoretical characterization of the ruthenium complex Ru[(μ-H) <sub>2</sub> BC <sub>8</sub> H <sub>14</sub> ] <sub>2</sub> (PCy <sub>3</sub> ) K. Essalah, J.-C. Barthelat, V. Montiel, S. Lachaize, B. Donnadieu, B. Chaudret, S. Sabo-Etienne . . . . .	182
High yield [ <sup>125</sup> I]iodide-labeling of iodinated carboranes by palladium-catalyzed isotopic exchange K.J. Winberg, G. Barberà, L. Eriksson, F. Teixidor, V. Tolmachev, C. Viñas, S. Sjöberg . . . . .	188
Ansa-metallocenes with B–N and B–P linkages: the importance of N–H···F–C hydrogen bonding in pentafluorophenyl boron compounds S.J. Lancaster, A.J. Mountford, D.L. Hughes, M. Schormann, M. Bochmann . . . . .	193
Enantioselective rhodium-catalysed addition of boronic acids using C <sub>2</sub> -symmetric aryl diphosphite ligands C.J. Chapman, K.J. Wadsworth, C.G. Frost . . . . .	206
Functionalization of the macropolyhedral borate anion cluster [B <sub>22</sub> H <sub>22</sub> ] <sup>2-</sup> : isolation and characterization of the OH and OEt derivatives O. Volkov, N.P. Rath, L. Barton . . . . .	212
Synthesis, structural characterization and reactivity of the amino borane 1-(NPh <sub>2</sub> )-2-[B(C <sub>6</sub> F <sub>5</sub> ) <sub>2</sub> ]C <sub>6</sub> H <sub>4</sub> R. Roesler, W.E. Piers, M. Parvez . . . . .	218
A new strategy for molecular modeling and receptor-based design of carborane containing compounds J. Johnsma, Y. Byun, T.P. Jones, Y. Endo, W. Tjarks . . . . .	223
Synthesis of ether- and carbon-linked polycarboranyl porphyrin dimers for cancer therapies M.F. Isaac, S.B. Kahl . . . . .	232
Two-electron homoaromatics with heteroatom bridges Y. Sahin, A. Ziegler, T. Happel, H. Meyer, M.J. Bayer, H. Pritzkow, W. Massa, M. Hofmann, P.R. Schleyer, W. Siebert, A. Berndt .	244
Synthesis and structure of potential Lewis acid–Lewis base bifunctional catalysts: 1- <i>N,N</i> -dimethylamino-8-borononaphthalene derivatives R.L. Giles, J.A.K. Howard, L.G.F. Patrick, M.R. Probert, G.E. Smith, A. Whiting . . . . .	257
Design of chiral boronate-substituted acrylanilides. Self-activation and boron-transmitted 1,8-stereoinduction in [4+2] cycloaddition J.W.J. Kennedy, D.G. Hall . . . . .	263
1,3,2-Diazastanna-[3]ferrocenophanes bearing alkyn-1-yl groups at tin and their 1,1-organoboration with triethylborane—molecular structure of a novel spirotin compound B. Wrackmeyer, H.E. Maisel, W. Milius, M. Herberhold . . . . .	271

Synthesis of enantiomerically pure vinylcyclopropylboronic esters via cross-metathesis P.G. Garcia, E. Hohn, J. Pietruszka . . . . .	281
Platination of [3-X-7,8-Ph <sub>2</sub> -7,8-nido-C <sub>2</sub> B <sub>9</sub> H <sub>8</sub> ] <sup>2-</sup> (X = Et, F): Synthesis and characterisation of slipped and 1,2 → 1,7 isomerised products S. Robertson, D. Ellis, G.M. Rosair, A.J. Welch . . . . .	286
Synthesis, structures and reactivity of macrocyclic imidazolylboranes A. Weiss, V. Barba, H. Pritzkow, W. Siebert . . . . .	294
The chemistry of the undecaborates O. Volkov, P. Paetzold . . . . .	301
Macropolyhedral boron-containing cluster chemistry. Aspects of the S <sub>2</sub> B <sub>16</sub> H <sub>16</sub> system. Preparation, structure, NMR spectroscopy and isomerism P.K. Dosangh, J. Bould, M.G.S. Londesborough, T. Jelínek, M. Thornton-Pett, B. Štíbr, J.D. Kennedy . . . . .	312
Synthesis of homoleptic Re(I) complexes of isocyano-carboranes and the X-ray structure of hexakis( <i>p</i> -carboran-1-yl-isocyanide)Re(I) P. Schaffer, J.F. Britten, A. Davison, A.G. Jones, J.F. Valliant . . . . .	323
Mechanistic and synthetic aspects of hydroboration with a simple atropisomeric ligand prepared from 1-(1'-(isoquinolyl)-2-naphthol) A. Korostylev, I. Gridnev, J.M. Brown . . . . .	329
Author Index of Volume 680 . . . . .	335
Subject Index of Volume 680 . . . . .	337
Contents of Volume 680 . . . . .	343