

Syntaxonomy and Syncology of *Quercus coccifera* Mediterranean Shrublands in Greece

Georgios Tsiorliris · Pavlos Konstantinidis · Pantelis Xofis

Received: 3 March 2009 / Accepted: 2 July 2009 / Published online: 22 July 2009
© The Botanical Society of Korea 2009

Abstract *Quercus coccifera* (kermes oak) is the most common species of the Mediterranean maquis with a wide distribution across the Mediterranean Basin. This paper presents a syntaxonomic overview of the *Q. coccifera* plant communities in the Mediterranean zone of Greece (*Quercetea ilicis*) based on the classification of 221 relevés from 34 (17 continental and 17 insular) mountainous areas throughout Greece. Two associations and eight sub-associations are described and presented in a synoptic constancy table. *Querco cocciferae*–*Pistacietaum lentisci* is the most widespread, is found in the entire continental Greece and most islands, and is further subdivided into five sub-associations reflecting primarily local peculiarities in the disturbance regime and the influence of local floristic elements. *Rhamno lycoidis*–*Cocciferetum* (Rivas Goday & Rivas-Martínez 1954), on the other hand, is geographically confined on the island of Crete and is further subdivided into three sub-associations, reflecting differences in the annual precipitation, and they are characterized by the presence of many phryganic and grazing-resistant species. Climate and the anthropogenic pressure have been identified to be the most important factors determining the

structure and the floristic composition of *Q. coccifera* Mediterranean shrublands of Greece.

Keywords Kermes oak · Mediterranean shrublands · Phytosociology · *Quercetea ilicis* · *Quercus coccifera* · Vegetation classification

Introduction

Quercus coccifera (kermes oak) is the most common species of the Mediterranean maquis, thriving in a wide variety of often contrasting environments (Balaguer et al. 2004). Characteristics, such as frugality, resprouting ability, and resilience to coppicing allow it to grow in unfavorable soil conditions and under severe human pressure (Liakos and Mouloupolous 1967). Overgrazing, overexploitation by humans and wildfires which have altered the physiognomy and composition of natural vegetation in the Mediterranean Basin, is thought to be one of the main factors responsible for the predominance of the species across the region (Mavrommatis 1980; Raus 1980; Trabaud 1987). Its prodigious regenerative capacity is supported by stored energy reserves in the roots (Konstantinidis et al. 2005) and by supplies of water and nutrients made available by an extensive and deeply penetrating rooting system (Tsiorliris 1992) that allows it to extract water reserves which are inaccessible to other species during the summer draught (Rambal 1984). It can, thus, maintain a high transpiration level for this period necessary to develop its young leaves. This gives it a competitive advantage over grasses and may explain the difficulties of establishing and maintaining perennial grasses after a brush-to-grass conversion (Poissonet et al. 1981).

Although *Q. coccifera* shrub formations do not usually exceed a height of 2 m, in the absence of grazing and fire, it

G. Tsiorliris (✉) · P. Konstantinidis
National Agricultural Research Foundation,
Forest Research Institute,
57006 Vassilika,
Thessaloniki, Greece
e-mail: gmtsiou@fri.gr
P. Konstantinidis
e-mail: pavkon@fri.gr

P. Xofis
Faculty of Natural Sciences, Imperial College London,
Wye campus, Wye, Ashford,
Kent TN25 5AH, UK

can form maquis formations up to 4–5 m high (Konstantinidis and Tsiorliris 2003) or even forests with a height of about 15 m, such as the ones observed mostly in Crete (Barbéro and Quézel 1980; Kypriotakis et al. 1996).

Q. coccifera is found across the entire Mediterranean region apart from Egypt (Ozenda 1964). In the Middle East, *Q. coccifera* is traditionally named *Quercus calliprinos*, which is a plain synonym of *Q. coccifera* (Greuter et al. 1986). It is distributed from 45°30' N in the Peninsula of Istria to 31°00' N in Negev (Israel) and from 9°30' E in Cascais (Portugal) to 37°00' E on the Mountains of Alaouites in Syria (Cañelas 1993). It dominates low and high shrub formations, such as the garrigue in Southern France (Lossaint and Rapp 1971; Poissonet et al. 1978), “coscojares” in Spain (Terradas 1999; Loidi et al. 1994), and “prinones” in continental Greece (Mavrommatis 1982; Papachristou 1998).

In Greece, *Q. coccifera* dominated vegetation formations occupy an area of more than 1.5 million hectares (Platis and Papanastasis 2003). It can be found within all life zones influenced by the Mediterranean climatic conditions (Konstantinidis 1990; Dimopoulos 1993), and its altitudinal distribution in Southern and Western Greece reaches 1,200 m asl, in Central Greece 1,000 m asl, and in Northern Greece 700 m asl (Mavrommatis 1980), while in some areas, such as in Crete, it can be found even higher (1,600 m) as recorded by Tsiorliris et al. (2001). It is most abundant in the continental part of Greece, apart from the northwest and northeastern parts, but it is also widespread in most islands apart from Cyclades. In the Thermo-Mediterranean life zone in particular, it codominates alongside with *Pistacia lentiscus* and in its drier parts with *Ceratonia siliqua* which are both character species of this life zone (Debazac 1969).

Several regional classifications were conducted across the Mediterranean region regarding the shrublands communities dominated by *Q. coccifera*. Many different associations and nomenclatures have been adopted while they often describe in fact the same plant communities, even at the level of sub-association. The most characteristic of those are Cocciferetum Braun-Blanquet (1924, 1936), Quercetum cocciferae Braun-Blanquet and Bolòs 1957, and Rhamneto-Cocciferetum Braun-Blanquet and Bolòs 1957 in western Mediterranean, as well as, Quercetum calliprini Boyko 1954 (Boyko 1954; Zohary 1973) in Eastern Mediterranean.

Only few studies, however, refer to the synecology and syntaxonomy of *Q. coccifera* shrublands in Greece (Adamović 1906, 1909 cited by Bergmeier 1990 and Theodoropoulos 1991; Horvat et al. 1974; Barbero and Quézel 1976; Mavrommatis 1980; Raus 1980, 1982; Karagiannakidou-Iatropoulou 1983; Konstantinidis 1990; Bergmeier 1990; Dimopoulos 1993; Chasapis et al. 2004).

The aim of the current study is to present a systematic classification of the *Q. coccifera* dominated Mediterranean shrubland formations of the Quercetea ilicis vegetation zone, which is its natural distribution area, throughout Greece, and to attempt to identify the ecological parameters affecting their distribution.

The prevalence of *Q. coccifera*-dominated plant communities in the Eastern Mediterranean landscape is reciprocal to the scarce syntaxonomic knowledge about this vegetation type. In this respect, the vegetation data provided by this study are generally worth being communicated to the scientific community.

Materials and Methods

Study Area

Phytosociological data across continental and insular Greece were collected on areas that were considered representative of the main Mediterranean shrubland types of *Q. coccifera*. The areas represent a variety of environmental conditions since they are distributed from north to south and east to west of Greece, covering almost the entire country (Fig. 1).

The study area includes 34 areas. Seventeen across continental Greece: Verticos and Cholomon (Northern Greece), Mavrovouni, Telethrio, and Ochi (Eastern Greece), Erimanthos, Mani, Chelmos, and Taigetos (Southern Greece), Acherontas, Akarnanika, Arakinthos, Arta, and Parga (Western Greece), Oiti, Sounio, and Parnitha (Central Greece). Three areas across Ionian Sea, namely: Ainos and Kalo Oros (Kefalonia island) and Lefkada. Five areas across Aegean Sea: Lesvos, Ampelos, and Kerkis (Samos Island) and Andros and Kithira. And nine areas across Crete: SE Crete, Moni Kapsa, and Thryptis (Eastern Crete), Dikti-Omalos, Dikti-Lasithi, Giouchtas, and Idi Oros (Central Crete) and Rodopos and Lefka Ori (Western Crete).

Table 1 shows the geological and climatic conditions and the number of relevés of the various study areas. The selected study areas cover most of the southeastern part of the Balkan Peninsula, located between the 34th and 40th parallel, with a meridional extent from 19° to 28° E, including Aegean Sea, Ionian Sea, and the East Mediterranean Sea.

The climate of the Quercetea ilicis vegetation zone is typical Mediterranean with mild and rainy winters, warm and dry summers, and an extended period of sunshine throughout the year. However, the climatic conditions vary from the semidry conditions of Attica, Crete, and Eastern Greece to the relatively subhumid to humid conditions of Northern and Western Greece.

Fig. 1 Map of Greece with the mountainous study areas



A common characteristic of all the above areas is their mountainous character and their sharp relief, independently of their size. The geologic structure is quite similar in most areas, where a calcareous substratum is dominant with parts of recently formed (Miocene and Pliocene) sea sediments (mostly in islands), while other geological formations are only occasionally met. The dominant soil type in the islands is terra rossa followed by rendzina and alluvial soils (Kagiabaki 2003). The calcareous substratum consists mainly of limestone, dolomite, and marls, while parts of flysch are found in between limestone (Vardinogianni 1994; Trichas 1996), resulting in soils with an alkaline reaction. The steep relief as well as the overgrazing and often occurrence of wildfires result in increased soil erosion and subsequent degradation of the soil chemical and physical properties (Vardinogianni 1994) leading often to desertification (Tsiourlis et al. 2001).

Data Collection and Analysis

From 1996 to 2006, 221 relevés were collected during May and June using the Braun-Blanquet (1964) approach. The

plots were located at various altitudes, expositions, inclinations, and relief. Plot size was set at 300 m², and an effort was made to achieve high ecological and physiognomic homogeneity within each plot (Dafis 1970; Athanasiadis 1986). The following parameters were recorded for each relevés: elevation, inclination, exposition, and species abundance for each vegetation layer. Vascular plants were identified using “Flora Europaea” (Tutin et al. 1964–1980; Diapoulis 1939–1949; Kavvadas 1956–1964). The nomenclature of taxa follows “Flora Europaea” (Tutin et al. 1964–1980).

The identification of vegetation units was done using the polythetic-divisive method TWINSPAN (Hill et al. 1975; Hill 1979). The original Braun-Blanquet scale of abundance was converted into an ordinal scale (Bergmeier and Dimopoulos 2001), with values ranging from 1 to 7 corresponding to the original seven classes of abundance. The pseudospecies cut levels used were 0.1, 3.1, 4.1, 5.1, and 6.1 ordinal scale units. Classification in five levels was performed, and the minimum number of samples to justify further division was set at 5. The maximum number of indicator pseudospecies was set at 7, and all pseudospecies

Table 1 Study areas of *Quercus coccifera* shrublands in Greece: location, main climatic parameters, and number of relevés

Areas	Geographic coordinates	Altitude (m)	Exposition (degree)	Slope (%)	Substrate	Climate	Number of relevés
Vertiskos	41°00', 23°20'	50–200	0–360	10–50	Limestone	Subhumid Mediter.	5
Cholomon	40°32', 23°15'	70–250	0–360	15–45	Limestone	Subhumid Mediter.	5
Mavrovouni	22°20', 36°55'	400–750	0–180	10–50	Limestone, schists	Subhumid Mediter.	7
Telethrio	23°09', 38°54'	50–450	0–360	20–70	Limestone	Subhumid Mediter.	5
Ochi	24°28', 38°07'	350–600	90–270	15–40	Limestone	Subhumid Mediter.	5
Erimanthos	22°00', 38°00'	400–700	270–360	35–55	Limestone, flysch	Subhumid Mediter.	5
Mani	22°26', 36°40'	180–360	180–270	10–40	Limestone	Subhumid Mediter.	9
Chelmos	22°15', 38°00'	400–850	360–45	20–35	Limestone	Subhumid Mediter.	5
Taigetos	22°20', 36°55'	400–900	135–315	20–40	Limestone	Subhumid Mediter.	9
Acherontas	20°30', 39°15'	30–450	90–270	20–40	Limestone	Humid Mediterranean	9
Akarnanika	21°26', 38°28'	160–370	45–220	5–50	Limestone	Subhumid Mediter.	5
Arakinthos	21°25', 38°30'	400–600	50–200	15–40	Limestone, flysch	Subhumid Mediter.	6
Arta	20°57', 39°13'	290–500	0–180	20–45	Limestone	Subhumid Mediter.	8
Parga	20°23', 39°18'	150–350	90–180	20–30	Limestone	Subhumid Mediter.	8
Oiti	22°19', 38°30'	400–600	150–360	10–60	Limestone	Subhumid Mediter.	5
Sounio	24°02', 37°41'	20–250	189–360	10–30	Limestone	Semiarid Mediter.	10
Parnitha	23°43', 38°10'	600–800	90–360	20–50	Limestone, schists	Humid Mediterranean	5
Ainos	20°43', 38°12'	90–350	45–220	5–40	Limestone	Humid Mediterranean	5
Kalo Oros	20°35', 38°20'	350–500	0–360	5–45	Limestone	Humid Mediterranean	10
Lefkada	20°36', 38°43'	90–270	0–360	5–25	Limestone	Humid Mediterranean	5
Lesvos	26°45', 39°20'	650–850	0–360	10–25	Limestone, schists	Subhumid Mediter.	6
Ampelos	26°50', 37°07'	250–550	0–360	10–40	Limestone	Subhumid Mediter.	5
Kerkis	26°40', 37°40'	400–750	150–270	20–60	Limestone	Subhumid Mediter.	5
Andros	24°50', 37°55'	200–550	90–270	15–50	Limestone	Subhumid Mediter.	5
Kithira	23°00', 36°20'	120–450	25–340	5–50	Limestone	Subhumid Mediter.	7
SE Crete	26°14', 35°04'	200–800	0–360	20–80	Limestone	Semiarid Mediter.	5
Moni Kapsa	26°03', 35°02'	400–550	0–360	50–90	Limestone	Semiarid Mediter.	5
Thryptis	25°52', 35°05'	200–1350	0–360	30–90	Limestone	Semiarid Mediter.	6
Dikti-Omalos	25°26', 35°03'	600–1400	0–360	0–80	Limestone	Semiarid Mediter.	7
Dikti-Lasithi	25°30', 35°08'	450–1600	0–360	0–90	Limestone	Semiarid Mediter.	11
Giouchtas	25°08', 35°14'	400–700	0–360	30–90	Limestone	Semiarid Mediter.	6
Idi Oros	24°50', 35°15'	450–1600	0–360	10–90	Limestone	Semiarid Mediter.	12
Rodopos	23°45', 35°36'	200–400	0–360	20–60	Limestone	Subhumid Mediter.	5
Lefka Ori	24°02', 35°17'	600–1300	45–220	20–60	Limestone	Subhumid Mediter.	5

were available as indicator species. The initial classification resulted in a large number of groups, but not all of them represented distinctive vegetation units, so some of the groups were merged at higher division levels. This process was aided by the results of a Detrended Correspondence Analysis (DCA; Hill 1973, 1974). TWINSPAN classification is based on a single-axis ordination for the differentiation of the groups and provides relatively little information on the floristic relationships between the groups and the degree of floristic overlapping between them (Gauch 1982). Ordination analysis using DCA, on the other hand, uses all the information contained in the samples by species data matrix, and in two-dimensional samples, ordination dia-

gram is more informative and allows the identification of both the floristic consistency of the identified groups and the floristic relationships between groups.

After construction of a synoptic table, the table was reordered into a diagnostic table clearly showing the differences between the plant communities. This table was used for the identification of diagnostic species and for denomination of the different plant communities.

The nomenclature of the plant communities was typified with the International Code of Phytosociological Nomenclature (ICPN; Weber et al. 2000). However, referred syntaxa names before 2002 are literally cited and not obligatory in accordance with ICPN. The definition of

“vegetation zones” and “subzones” follows Dafis (1973) and Athanasiadis (1986) for Greece.

Results

Based on TWINSPAN and DCA results, eight distinctive vegetation units were identified (Fig. 2) and, following the Braun-Blanquet approach, were classified under two associations and eight sub-associations. Table 2 shows the frequencies of the diagnostic species for the corresponding vegetation units. According to these results, the syntaxonomic synopsis of the *Q. coccifera* shrublands in the mountains of Greece is:

Quercetea ilicis Braun-Blanquet 1947

Quercetalia ilicis Braun-Blanquet 1947

Quercion ilicis Braun-Blanquet 1934 em. Rivas-Martínez 1975

1. *Querco cocciferae*–*Lentiscetum* Braun-Blanquet et al. 1936

Or typified with ICPN (Weber et al. 2000) as *Querco cocciferae*–*Pistacietaum lentisci*

- 1.1 *Querco cocciferae*–*Pistacietaum lentisci typicum*
- 1.2 *Querco cocciferae*–*Pistacietaum lentisci euphorbiotosum rigidae*
- 1.3 *Querco cocciferae*–*Pistacietaum lentisci arbutotosum andrachni*

- 1.4 *Querco cocciferae*–*Pistacietaum lentisci fraxinetosum orni*
- 1.5 *Querco cocciferae*–*Pistacietaum lentisci anthemidetosum tinctoriae*
- 2. *Rhamno lycoidis*–*Quercetum cocciferae* Braun-Blanquet and Bolòs 1954
 - 2.1 *Rhamno lycoidis*–*Quercetum cocciferae phagnaletosum graeci*
 - 2.2 *Rhamno lycoidis*–*Quercetum cocciferae berberidetosum cretiae*
 - 2.3 *Rhamno lycoidis*–*Quercetum cocciferae cerastietosum brachypetali*

- 1. *Querco cocciferae*–*Pistacietaum lentisci* (Table 2, Group 1; Fig. 2)

Character species: *P. lentiscus*, *Phillyrea media*, *Asparagus acutifolius*, *Geranium lucidum*, *Ruscus aculeatus*, *Medicago minima*, *Clematis vitalba*, *Teucrium capitatum*, *Pyrus spinosa*, *Teucrium chamaedrys*, *Smilax aspera*, *Quercus pubescens*, *Medicago lupulina*, *Anthyllis hermanniae*, *Rosa canina*, *Melica ciliata*, *Paliurus spina-christi*.

Ecology: The climate on the areas where this association is found is characterized by warm winters (mean temperature 0–5°C) with precipitations that can reach 600–1,000 mm. Altitudinal range is between 20 and 900 m. All the expositions are encountered while the slopes vary from relatively gentle to very steep. Substrates are mainly limestone with several areas on flysch and schist, while the

Fig. 2 DCA samples ordination diagram. The codes in the legend correspond to the group numbers assigned to the sub-associations as following: 1.1 *Querco cocciferae*–*Pistacietaum lentisci typicum*, 1.2 *Querco cocciferae*–*Pistacietaum lentisci euphorbiotosum rigidae*, 1.3 *Querco cocciferae*–*Pistacietaum lentisci arbutotosum andrachni*, 1.4 *Querco cocciferae*–*Pistacietaum lentisci fraxinetosum orni*, 1.5 *Querco cocciferae*–*Pistacietaum lentisci anthemidetosum tinctoriae*, 2.1 *Rhamno lycoidis*–*Quercetum cocciferae phagnaletosum graeci*, 2.2 *Rhamno lycoidis*–*Quercetum cocciferae berberidetosum cretiae*, 2.3 *Rhamno lycoidis*–*Quercetum cocciferae cerastietosum brachypetali*




Table 2 Synoptic table of the *Quercus coccifera* Mediterranean shrublands showing the frequencies of species in each of the identified associations and sub-associations

Group number	1.	2.	3.	4.	5.	1.	2.	3.	4.	Group number	1.	2.	3.	4.	5.	1.	2.	3.
Number of relevés	86	6	41	28	6	27	23	23	4	Number of relevés	86	6	41	28	6	27	23	23
Character species of Quercetea, Quercetalia, Quercion ilicis																		
<i>Quercus coccifera</i>	V	V	V	V	V	V	V	V	V	<i>Fumana thymifolia</i>	III	I	I	I	I			
<i>Pistacia lentiscus</i>	II	II	IV	III	III	I	I	III	Carex distachya	III								
<i>Lonicera implexa</i>	II	III	I	II	III		I		<i>Cistus monspeliensis</i>	III								
<i>Rubia peregrina</i>	II	IV	III	II	II	IV			<i>Vicia cretica</i>	III								
<i>Calicotome villosa</i>	III	I	III	IV	IV	III	I		<i>Brachypodium sylvaticum</i>	III	II	I						
<i>Quercus ilex</i>	I	V	I	I					<i>Tamus communis</i>	III								
<i>Phlomis fruticosa</i>	I	II	IV	IV	II	I	I		<i>Lagurus ovatus</i>	III	I							
<i>Cistus creticus</i>	IV	I	IV	IV	IV	III	I	II	<i>Campanula glomerata</i>	III								
<i>Cistus salvifolius</i>	II		III	II	II	II			<i>Helictotrichon convolutum</i>	II		I						
<i>Euphorbia acanthothamnus</i>	I			II	I	III	II		<i>Lonicera etrusca</i>	II			I					
<i>Erica manipuliflora</i>	I		IV	III	I				<i>Piptatherum miliaceum</i>	II								
<i>Juniperus oxycedrus</i>	II		III	I	I				<i>Lonicera xylosteum</i>	II								
<i>Myrtus communis</i>	III		IV	II	I	I			<i>Centaurium tenuiflorum</i>	II								
<i>Olea europaea</i>	III	V	IV	III	II	III			<i>Psoralea bituminosa</i>	II								
<i>Urginea maritima</i>	I		II	IV		IV	III		<i>Melica minuta</i>	II								
<i>Spartium junceum</i>	I		I	III	II	I			<i>Micromeria graeca</i>	II								
<i>Genista acanthoclada</i>	I			III		III			<i>Muscaris comosum</i>	II								
<i>Cercis siliquastrum</i>	I	I	IV	I	II				<i>Scorzonera crocifolia</i>	II								
<i>Micromeria juliana</i>	I	IV	V	III	I	I			<i>Petrorhagia dubia</i>	II	I							
<i>Arbutus unedo</i>	II		V	III	II				<i>Lavandula stoechas</i>	II								
<i>Erica arborea</i>	II		IV	II	II		I		<i>Valantia muralis</i>	II								
<i>Clematis flammula</i>	I		IV	III	I				<i>Asterolinon linum-stellatum</i>	II								
1. Character species of Querco cocciferae-Pistacietum																		
1.1 Character species of Querco cocciferae-Pistacietum lentisci typicum																		
<i>Pistacia lentiscus</i>	IV	II	IV	IV	IV				<i>Bromus fasciculatus</i>	II								
<i>Phillyrea media</i>	IV	V	V	IV	IV	II			<i>Teucrium alpestre</i>	II								
<i>Asparagus acutifolius</i>	V	V	IV	V	IV				<i>Papaver rhoes</i>	II								
<i>Geranium lucidum</i>	II	V	IV		I				<i>Bupleurum fruticosum</i>	II								
<i>Ruscus aculeatus</i>	I		III	IV	I				<i>Anemone coronaria</i>	II						II	II	
<i>Medicago minima</i>	I	V	IV	I	I				<i>Melittis melissophyllum</i>	II								
<i>Clematis vitalba</i>	II		IV	III			I		<i>Silene gigantea</i>	II								
<i>Teucrium capitatum</i>	III		III	III	I				<i>Astragalus monspessulanum</i>	II								
<i>Pyrus spinosa</i>	II	III	III	I	I	I			<i>Euphorbia cyparissias</i>	II								
<i>Teucrium chamaedrys</i>	I	III	I	I	I	II			<i>Scutellaria rupestris</i>	II								
<i>Smilax aspera</i>	II		III	III	II	I			<i>Trifolium hirtum</i>	II								
<i>Quercus pubescens</i>			II	IV	I				<i>Potentilla micrantha</i>	IV	I	I						
<i>Medicago lupulina</i>			II	II	I	I			<i>Helichrysum microphyllum</i>	III			I					
<i>Anthyllis hermanniae</i>	I	IV	IV	I	I				<i>Thlaspi perfoliatum</i>	II		I	I	I				
<i>Rosa canina</i>	I	II	III	I	I				<i>Piptatherum coeruleescens</i>	IV				II				
<i>Melica ciliata</i>	I		IV	II		I			<i>Gastridium ventricosum</i>	III								
<i>Paliurus spina-christi</i>	III	I	III	II					<i>Crepis cretica</i>	III								
1.2 Character and diagnostic species of Querco cocciferae-Pistacietum lentisci euphorbietsosum rigidae																		
<i>Euphorbia rigida</i>	I	II							<i>Muscaris spreitzenhoferi</i>	III								
<i>Juniperus phoenicea</i>	IV																	
<i>Thymus sibthorpii</i>	V																	
<i>Vicia parviflora</i>	V																	
<i>Fumana arabica</i>	III	I																
<i>Arisarum vulgare</i>	III																	
<i>Prasium majus</i>	III																	

Table 2 (continued)

Group number	1.1	1.2	1.3	1.4	1.5	2.1	2.2	2.3	Group number	1.1	1.2	1.3	1.4	1.5	2.1	2.2	2.3
Number of relevés	86	6	41	28	6	27	23	4	Number of relevés	86	6	41	28	6	27	23	4
Anemone pavonina	III	I							Orlaya daucoides					IV			
Silene cretica	III	I	I	I	I				Avena sterilis	I	I			III			
Trifolium grandiflorum	III								Ballota acetabulosa		I	I		III			
Veronica arvensis	I	III	I						Bromus squarrosus					III			
Arabis verna		III				I	II		Sonchus oleraceus		I	I	III				
Cardamine hirsuta		III	I			II			Carex halleriana	I		I	II				
Torilis nodosa		II	I	I					Convolvulus althaeoides		I	II	I				
Vicia cuspidata		II		I					Dorycnium pentaphyllum	I		I	II				
Sherardia arvensis		II		I					Galium laconicum		I	II					
Aubrieta deltoidea		II	I	I					Milium vernale		I	I					
Cerastium comatum	I	II															
Minuartia hybrida	I		I														
1.3 Character and diagnostic species of Quercococciferae-									2. Character species Rhamno lycioidis-Quercetum cocciferae Braun-Blanquet & Bolòs 1954								
Pistaciëtum lentisci arbuetosum andrachni									Rhamnus lycioides	I	I		IV	II	IV		
Arbutus andrachne	V	III							Asparagus aphyllus		I		V	IV			
Melica rectiflora	V	III	II						Daphne sericea	I		II	III	III	V		
Colutea arborescens	IV								Centaurea raphanina		II		II	IV	II		
Stipa capensis	IV		I						Sarcopoterium spinosum	II	II		III	I			
Trifolium squamosum	I	IV							Lamyropsis cynaroides		I	II	III	II			
Fumana procumbens	IV	III							Reichardia picroides		II		II	I	II		
Origanum vulgare	IV	I							Origanum onites	I		I	II	I			
Hippocrepis emerus	IV	III		II					Orchis quadripunctata		I	II	I	II			
Ditrichia viscosa	IV								Cyclamen graecum		I	I	III				
Trifolium scabrum	IV	I							Biscutella didyma		I	I	I	II			
Agrostis capillaris	IV								Satureja thymbra	I	I	I	I	II			
Laurus nobilis	IV		I						Phlomis lanata		I		IV	II			
Convolvulus elegantissima	III	I							Acer sempervirens		II		II	V	II		
Vicia tenuifolia	IV								Asteriscus spinosus	I		II	III	I			
Silene atropurpurea	III	II		I					Euphorbia characias		I	I	III	II			
Bromus arvensis	III	I	I						Lathyrus aphaca		I		I	III			
Carpinus orientalis	I		III														
Crepis neglecta		II															
Rhus coraria		II	V														
Bromus intermedius		III	I	I	I												
1.4 Character and diagnostic species of Quercococciferae-									2.1 Character and diagnostic species of Rhamno lycioidis - Quercetum cocciferae phagnaletosum graeci								
Pistaciëtum lentisci fraxinetosum ornii									Phagnalon graecum		II		IV	I			
Fraxinus ornus	I								Linum strictum		I		II		II		
Crataegus monogyna	I	III		IV					Gagea graeca		I		II				
Crepis hellenica		III	I						Cyclamen creticum		I		II	I			
Hedera helix	I		III		I				Rumex tuberosus		I		II	I			
Abies cephalonica	I		I	III					Anemone hortensis			I	III	I			
Hypocharaeris achyrophorus	I			III	I	I			Anthyllis vulneraria		II	I	III				
Campanula ramosissima			II						Ricotia cretica	I			II				
Clinopodium vulgare			II		I				Asphodeline lutea			III	I				
Lathyrus laxiflorus		I		II	I				Sedum litoreum	I			I				
Adiantum capillus-veneris			I														
Centaurea cyanus		I															
1.5 Character and diagnostic sp. of Quercococciferae-									2.2 Character and diagnostic sp. of Rhamno lycioidis - Quercetum cocciferae berberidetosum cretiae								
Pistaciëtum lentisci anthemidetosum tinctoriae									Berberis cretica								
Anthemis tinctoria	II		IV						Daucus guttatus		I		IV	II			
Aegilops triuncialis	I		IV						Arum idaeum			I	III				

Table 2 (continued)

Group number	1.1	1.2	1.3	1.4	1.5	2.1	2.2	2.3	Group number	1.1	1.2	1.3	1.4	1.5	2.1	2.2	2.3
Number of relevés	86	6	41	28	6	27	23	4	Number of relevés	86	6	41	28	6	27	23	4
2.3 Charact. and diff. species of Rhamno lycoidis -																	
Quercetum cocciferae cerastietosum brachypetalii																	
Cerastium brachypetalum	II			I	V				Geranium molle		II	I	I	I			
Polygala venulosa	I		I						Campanula spatulata		III	I		II	III	II	
Arenaria leptoclados									Blackstonia perfoliata		I	I					
Alyssum minutum	I								Cyclamen hederifolium	I	II		II	I			
Rhamnus alaternus			I						Teucrium divaricatum			I		I			
Companion species																	
Bromus sterilis	II	V	III	I	III	II			Ceratonia siliqua			I	I				
Salvia fruticosa	I			II	I	III	I	II	Daucus carota		I	I					
Sanguisorba minor	I	I	I	I	II		II		Dorycnium hirsutum		II	II					
Anthoxanthum odoratum	I	IV	IV	I	I		II	II	Echinops ritro		I		I		II		
Scaligeria napiformis	I	I	I	III	I	II	I	II	Erodium cicutarium		I	I		I			
Lamium amplexicaule		IV		I			III	II	Galactites tomentosa	I		I		I			
Stellaria media		II		III	I	I			Galium monachinii				I	I			
Stipa bromoides		I	IV	III		II	I		Helichrysum conglobatum		I			I			
Pteridium aquilinum		III	I	I	III				Hordeum bulbosum		I	I	I	I			
Carlina corymbosa	II	II	I	III		II	III	II	Hordeum murinum		I		I	I			
Cynosurus effusus		II	I	I		II	I		Hypericum perforatum			II	I				
Poa bulbosa		II	II		II	I			Iris unguicularis		I	II		II	I		
Dactylis glomerata	II	IV	IV	IV		IV			Knautia integrifolia		I	I					
Galium aparine		V	I	III	III		I		Legousia hybrida					I			
Galium murale		V	I	I	I	III	II		Leontodon cichoriaceus	I	I				II		
Leontodon tuberosus		IV		II		III	I		Lolium rigidum		I	I		I			
Trifolium angustifolius	I		IV	II					Micromeria nervosa			I		I			
Trifolium campestre		II	IV	III	II	II	I		Myosotis ramosissima	I		I		I			
Trifolium fragiferum			IV	II		II			Ononis spinosa		I		I		II		
Trifolium physodes	V	III	I			II	III		Picnomon acarna	I	I		I	I			
Trifolium stellatum		II	II	III	II	II			Scorzonera cretica		I		I				
Trifolium tomentosum			I		II	I	II		Silene vulgaris		I	I	I		II		
Trifolium uniflorum		II		I		II	I	IV	Stellaria pallida			I		I			
Aira elegantissima		II		I	II			I	Tordylium apulum			II					
Allium subhirsutum				III		II			Urospermum picroides		I	I					
Cynosurus echinatus	I		IV	I		I	III		Vicia grandiflora		I				II		
Geranium robertianum			II	I		III	I		Anthemis chia	I		I	I	I	I		
Eryngium campestre	II		II	I		II	III	II									
Geranium rotundifolium			II	I	I		II										
Osyris alba			I	I		III											
Asplenium ceterach		I	II	I													
Avena barbata			IV	III		III											
Brachypodium distachyon			I	II	I												
Briza maxima			I	II			I										
Coridothymus capitatus	IV		IV	III		III											
Carex flacca	I		III	II		II											
Hypericum empetrifolium	II		III	III	II	III											
Crepis fraasii			III	II		II	III										
Brachypodium retusum			IV		II	II	II	II									
Asphodelus ramosus			III			IV	III	II									
Bromus madritensis			I	I		I	II										
Catapodium rigidum			I	I		I	I	II									
Crucianella latifolia	I		I	II			I	II									

Species with a single appearance have been omitted. Character and diagnostic species for each vegetation unit are shown in rectangles. Frequency classes are: I=species recorded in 1–20% of all relevés of each vegetation unit, II=21–40%, III=41–60%, IV=61–80%, V=81–100%

soil properties vary from relatively deep and productive to shallow and infertile. The shrub layer cover varies from 20% to 90%, and the understory is rather dense reaching up to 70%.

Distribution records: Cholomon, Lesvos, Mavrovouni, Akarnanika, Oiti, Telethrio, Ainos, Taigetos, Aherontas, Arakinthos, Erimanthos, Chelmos, Arta, Parga, Andros, Kithira, Sounio, Mani, Ochi, Dikti-Lasithi, Parnitha, Lefkada, Kalo Oros, Rodopos, Ampelos, Kerkis, and Lefka Ori.

Nomenclatural and syntaxonomic notes: This association is widespread in the Mediterranean Basin and it is one of the first associations described by Braun-Blanquet et al. (1936), in Southern France and Western Mediterranean. As a result of its wide distribution the same association has been described by many researchers, while in some cases, the same formations are classified under different associations. We adopted the same typification as Rivas-Martínez et al. (2001, 2002), which renamed it as Querco cocciferae–Pistacietum lentisci Br.-Bl., Font Quer, G. Br.-Bl., Frey, Jansen, & Moor 1936 nom. mut. propos (art. 45; addenda) in the checklist of plant communities of Spain. Despite its wide distribution and the subsequent variation in the local ecological conditions, there is a constant presence of diagnostic Mediterranean shrubs of higher taxa (e.g., *Quercus ilex*, *Olea europaea* var. *sylvestris*, *Pistacia terebinthus*, *Calicotome villosa*, *Erica manipuliflora*, *Spartium junceum*, *Arbutus unedo*, *Erica arborea*).

As expected, an association with such a wide distribution is further distinguished in sub-associations reflecting variation in its composition as a result of local floristic influences and variations in the ecological conditions.

1.1 Querco cocciferae–Pistacietum lentisci typicum (Table 2, Group 1.1; Fig. 2)

Character species: *Juniperus phoenicea*, *Thymus sibthorpii*, *Vicia parviflora*, *Fumana arabica*, *Arisarum vulgare*, *Prasium majus*, *Fumana thymifolia*, *Carex distachya*, *Cistus monspeliensis*, *Vicia cretica*, *Brachypodium sylvaticum*, *Tamus communis*, *Lagurus ovatus*, *Campanula glomerata*, *Potentilla micrantha*, *Helichrysum microphyllum*, *Piptatherum coerulescens*, *Gastridium ventricosum*, *Crepis cretica*, *Muscari spreitzenhoferi*.

Ecology: Mean ecological parameters of the association: subhumid Mediterranean climate, limestone (flysch and schist) substrates; mean slopes 30%, shrub cover of 60%, and herbs 40%; moderate grazing.

Distribution records: Andros, Kithira, Taigetos, Aherontas, Arakinthos, Arta, Telethrio, Mavrovouni, Sounio, Mani, Ochi, Dikti-Lasithi, Parnitha, Lefkada, Ainos, Kalo Oros, Rodopos, Verticos, Holomon, Ampelos, Kerkis, Lefka Ori.

Nomenclatural and syntaxonomic notes: The sub-association shares most of the floristic characteristics

described above for the association, including the majority of the typical Mediterranean evergreen shrubs. *J. phoenicea* is a character species of the sub-association especially on areas with a strong maritime influence. On the other hand, Mavrommatis (1980) distinguished similar communities with the abundance of *J. phoenicea* as Oleo-lentiscetum juniperetosum. There are also similarities with plant communities outside Greece including Pistacio lentisci–Juniperetum phoeniceae Trinajstić 1987 identified by Pandža (2004) in Croatia (island of Murter) and Quercetum cocciferae brachypodietosum or Cocciferetum brachypodietosum in the Quercetum cocciferae association identified by Wraber (1952) in Southern France.

1.2 Querco cocciferae–Pistacietum lentisci euphorbietosum rigidae (Table 2, Group 1.2; Fig. 2)

Character and diagnostic species: *Euphorbia rigida*, *Prunus cocomilia*, *Ornithogalum nutans*, *Lagoecia cuminoidea*, *Ranunculus rumelicus*, *Anemone pavonina*, *Silene cretica*, *Trifolium grandiflorum*, *Veronica arvensis*, *Arabis verna*, *Cardamine hirsuta*.

Ecology: Altitudinal range of 650–850 m, subhumid Mediterranean climate, limestone (schist) substrates, gentle slopes (10–25%), mean shrub cover 30% and herbs 40%; severe grazing and frequent wild fires.

Distribution records: Lesvos.

Nomenclatural and syntaxonomic notes: This sub-association is floristically the most dissimilar in comparison with Querco cocciferae–Pistacietum lentisci typicum. Due to its proximity to Turkey, it is enriched with floristic elements from Minor Asia (Rechinger 1943; Davis 1965–1985; Edmondson 1982; Hansen and Nielsen 1993). Further, its distribution in relatively higher altitudes leads to the absence of character shrubs and climbing species of Quercion ilicis, such as *Q. ilex*, *C. villosa*, *O. europaea* var. *sylvestris*, *Myrtus communis*, *Cercis siliquastrum*, *Cistus salviifolius*, *Urginea maritima* (*Drimia maritima*). The sub-association has a floristic relationship to the Siderito dichotomae–Quercetum cocciferae identified by Karaer et al. (1999) in Keltik Valley of Minor Asia.

1.3 Querco cocciferae–Pistacietum lentisci arbutetosum andrachni (Table 2, Group 1.3; Fig. 2)

Character and diagnostic species: *Arbutus andrachne*, *Melica rectiflora*, *Colutea arborescens*, *Stipa capensis*, *Trifolium squamosum*, *Fumana procumbens*, *Origanum vulgare*, *Hippocrepis emerus*, *Dittrichia viscosa*, *Trifolium scabrum*, *Agrostis capillaries*, *Laurus nobilis*, *Convolvulus elegantissima*, *Vicia tenuifolia*, *Silene atropurpurea*, *Bromus arvensis*, *Carpinus orientalis*, *Bromus intermedium*.

Ecology: Mean altitude of 400 m (90–900 m), subhumid to humid (more than 1,000 mm of precipitations) Mediterranean climate, limestone (flysch) substrates, mean slopes

30%, mean shrub cover 60% (30–80%), and herbs cover 40%; moderate grazing.

Distribution records: Western Greece and along the central axis of peninsular Greece: Akarnanika, Oiti, Telethrio, Ainos, Taigetos, Acherontas, Arakinthos, Erimanthos, Chelmos, Arta, Parga.

Nomenclatural and syntaxonomic notes: The character species are more hygrophilous including *A. andrachne*, *C. arborescens*, *L. nobilis*, *C. orientalis*, and *Rhus coriaria*.

This sub-association has similarities with plant communities beyond the Quercetea ilicis in Northern Greece including *Q. cocciferae*–*C. orientalis* described by Chasapis et al. (2004) in Chortiatis and Coccifero–Carpinetum described by Oberdorfer (1948). We noted also some similarities with plant communities outside Greece such as Melico–Quercetum cocciferae Br. Bl., Silva and Rozeira 1964 identified in Southern France.

1.4 Querco cocciferae–Pistacietum lentisci fraxinetosum orni (Table 2, Group 1.4; Fig. 2)

Character and diagnostic species: *Fraxinus ornus*, *Crataegus monogyna*, *Crepis hellenica*, *Hedera helix*, *Abies cephalonica*, *Hypochoeris achyrophorus*.

Ecology: Mean altitude of 550 m (150–900 m), subhumid to humid Mediterranean climate, limestone (flysch) substrates, mean slopes 30%, mean shrub cover 60% (20–80%), and herbs cover 50%; moderate to severe grazing.

Distribution records: Western Greece and Peloponnese: Acherontas, Arakinthos, Arta, Parga, Oiti, Erimanthos, Chelmos, Taigetos.

Nomenclatural and syntaxonomic notes: The floristic composition of the sub-association includes a large number of relatively hygrophilous species (e.g., *F. ornus*, *H. helix*, *Clinopodium vulgare*, *Lathyrus laxiflorus*, and *Adiantum capillus-veneris*) while the endemic *A. cephalonica* is present in higher altitudes. Querco cocciferae–Pistacietum lentisci fraxinetosum orni appears to have similarities with plant communities outside Greece and especially with Crataego monogynae–Quercetum cocciferae described by Martínez-Parras et al. (1985), in the medium Mediterranean subhumid bioclimatic zone, in Spain. Further, this sub-association is reported by Ladero et al. (2002) in the surroundings of Alhama de Granada Baths and Costa et al. (1998) in Portugal.

1.5 Querco cocciferae–Pistacietum lentisci anthemidetosum tinctoriae (Table 2, Group 1.5; Fig. 2)

Character and diagnostic species: *Anthemis tinctoria*, *Aegilops triuncialis*, *Orlaya daucoides*, *Avena sterilis*, *Ballota acetabulosa*, *Bromus squarrosus*, *Sonchus oleraceus*.

Ecology: Altitudinal range of 450–750 m, subhumid Mediterranean climate, schist (limestone) substrates, mean slopes 25%, northeast–south orientations, mean shrub cover

50% (30–70), and herb cover 60%; moderate to severe grazing.

Distribution records: Mavrovouni.

Nomenclatural and syntaxonomic notes: Despite the subhumid climate, we noted the presence of xerophilic species probably as a result of severe grazing. Raus (1980) has distinguished within the shrublands of Mavrovouni, a degraded low forest of the Ostryo–Carpinion orientalis (Quercetalia pubescantis, Querco–Fagetea), rich in *C. orientalis*, on areas that have sufficient water supply (*Q. cocciferae*–*C. orientalis* community).

2. Rhamno lycioidis–Quercetum cocciferae Braun–Blanquet and Bolòs 1954 (Table 2, Group 2; Fig. 2)

Character species: *Rhamnus lycioides*, *Asparagus aphyllus*, *Daphne sericea*, *Centaurea raphanina*, *Sarcopoterium spinosum*, *Lamyropsis cynaroides*, *Reichardia picroides*, *Origanum onites*, *Orchis quadripunctata*, *Cyclamen graecum*, *Biscutella didyma*, *Satureja thymbra*, *Phlomis lanata*, *Acer sempervirens*, *Asteriscus spinosus*, *Euphorbia characias*, *Lathyrus aphaca*.

Ecology: Mean ecological parameters of the association: semiarid Mediterranean climate, limestone substrates, terra rossa, rendzina, and alluvial soils; all the orientations, mean slopes 40%, shrub cover 40%, and herb cover 10%; moderate to severe grazing. Especially for Crete, there is a precipitation gradient from west (subhumid) to east (semidry) and from the sea level to the top of the three main mountainous massives (Lefka Ori, Idi, and Dikti).

Distribution records: Mainly in Crete and some other islands of the Aegean: Dikti-Lasithi, Dikti-Omalos, Thryptis, Moni Kapsa, SE Crete, Giouchtas, Idi, Lefka Ori, Sounio, Ochi, Andros.

Nomenclatural and syntaxonomic notes: The character species is *R. lycioides* (*R. graecus*), which thrives in sunny and dry environments. The abundance of *S. spinosum*, *P. lanata*, *Ononis onites*, *E. characias*, *Phagnalon graecum*, and *S. thymbra* demonstrates the important phryganic influence of the association. This association which is mainly restricted to Crete includes many local endemic species, such as *Ph. lanata*, *Ebenus cretica*, *C. raphanina* subsp. *raphanina*, *Scorzonera cretica*, *Teucrium microphyllum*, *Teucrium alpestre*, *C. cretica*, and some protected species such as *Ophrys* spp. or *Orchis* spp. There is a wide variety of shrublands, from well conserved and dense to more degraded accompanied by species like *U. maritima* (*D. maritima*), *Asphodelus aestivus*, and *Asphodeline lutea*. This association has been identified by many phytosociologists working in the western Mediterranean and particularly in the Iberian Peninsula in regions of similar bioclimatic type. The association presents similarities with Asparago–Rhamnetum oleoidis cocciferetum Rivas Goday 1959 and

Rhamno velutini–Quercetum cocciferae (Rivas-Martinez et al. 2001). Moreover, Rivas-Martinez et al. (2001) identified in the Iberian Peninsula a superior taxonomic unit the Rhamno lycoidis–Quercion cocciferae Rivas Goday ex. Rivas Martinez 1975.

2.1 Rhamno lycoidis–Quercetum cocciferae phagnaletosum graeci (Table 2, Group 2.1; Fig. 2)

Character and diagnostic species: *P. graecum*, *Linum strictum*, *Gagea graeca*, *Cyclamen creticum*, *Rumex tuberosus*, *Anemone hortensis*, *Anthyllis vulneraria*, *Ricotia cretica*, *A. lutea*, *Sedum litoreum*.

Ecology: Wide altitudinal range from 20 to 1,600 m with a mean of 700 m; mean slopes 40%, shrub cover of 30–90% (mean 40%), and herb cover 20%. The areas where the sub-association is found have the driest climate in the country and grazing is severe.

Distribution records: Dikti-Lasithi, Dikti-Omalos, Thryptis, Moni Kapsa, SE Crete, Giouchtas, Idi, Sounio, Ochi.

Nomenclatural and syntaxonomic notes: This sub-association is considered as the more xerothermic expression of the association and is mainly distributed in the eastern and central parts of Crete comprising many phryganic species, such as *P. graecum*, *A. vulneraria*, *A. lutea*.

2.2 Rhamno lycoidis–Quercetum cocciferae berberidetosum cretiae (Table 2, Group 2.2; Fig. 2)

Character and diagnostic species: *Berberis cretica*, *Daucus guttatus*, *Arum idaeum*, *Erophila verna*, *Cardamine graeca*, *Anthemis rigida*.

Ecology: Altitudinal range from 200 to 1,600 m with a mean of 800 m; mean slopes 40%, shrub cover between 40–90% (mean 50%) and herb cover 10%; Severe grazing.

Distribution records: Dikti-Lasithi, Dikti-Omalos, Thryptis, Idi, Andros.

Nomenclatural and syntaxonomic notes: The sub-association was identified in the higher altitudes and degraded areas of Eastern and Central Crete, where grazing activities are important. The character species is the well adapted to grazing spiny shrub of Cretan mountains *B. cretica*, which originates from the neighboring orophrygana. Orophrygana is the endemic habitat of the upper altitudinal zone of Cretan mountains (>1,100 m), characterized by *B. cretica*, *Astragalus angustifolius*, and *Euphorbia acanthothamnos*, which results from the broad contact between phrygana (in lower altitudes) and hedgehog heaths in supra- and oro-Mediterranean zones of Crete (Tsiourlis et al. 2001). The sub-association includes many Cretan endemic species, such as *Centaurea idaea*, which is typical in high altitudes.

2.3 Rhamno lycoidis–Quercetum cocciferae cerastietosum brachypetalii (Table 2, Group 2.3; Fig. 2)

Character and diagnostic species: *Cerastium brachypetalum*, *Polygala venulosa*, *Arenaria leptoclados*, *Alyssum minutum*, *Rhamnus alaternus*.

Ecology: Altitudinal range from 600 to 1,300 m; mean slopes 40%, shrub cover of 50–90% (mean 60%) and herbs 15%; Severe grazing. The main character of the areas where the sub-association is found is the high level of precipitations which exceeds 1,500 mm (Grove et al. 1993). Shallow calcareous lithosols and rendzina soils dominate throughout much of the massif. They often represent degraded soil profiles with limited water supply (Vogiatzakis et al. 2003); Severe grazing.

Distribution records: Lefka Ori.

Nomenclatural and syntaxonomic notes: *P. venulosa* is a species with distribution in Southern Greece, the Aegean, Anatolia, and Cyprus. Lefka Ori has a high number of endemic plants (Strid 1996).

Discussion

Q. coccifera Greek shrublands have suffered dramatic changes under anthropogenic activities (coppice, overgrazing, and deforestation) that have affected both their physiognomy as well as their floristic composition. Human pressure has also affected the distribution of these formations which often extend beyond their natural distribution limits, occupying areas abandoned by species less resistant to human activities (Dafis and Jahn 1975; Barbero and Quézel 1976; Raus 1980; Pavlides 1982; Vardakakis et al. 1987; Bergmeier 1990; Konstantinidis 1990; Theodoropoulos 1991; Dimopoulos 1993; Chasapis et al. 2004). Moreover, this highly anthropogenic distribution and composition of *Q. coccifera* shrublands led some ecologists to suggest that these ecosystems should not be classified below the level of the association (Raus 1980; Bergmeier 1990). The unwillingness, however, to classify *Q. coccifera* shrublands below the association level has resulted in the “blanket” classification of all *Q. coccifera* dominated shrublands under Quercetum cocciferae, based only on physiognomic criteria and avoiding further classification, which in reality shows important floristic variation.

Shrublands with *Q. coccifera* in continental and insular Greece, apart from Crete, are mainly distributed within the growth zone of *P. lentiscus* that characterizes the Thermo-Mediterranean bioclimate (Debazac 1969; Mavrommatis 1980). These shrubland formations consist mainly of sclerophyllous species such as *Q. coccifera*, *P. latifolia*, and *J. phoenicea*, while some thermophilous deciduous broadleaved species often participate at lower percentages.

The results of the current study suggest that the *Q. coccifera* shrublands in Greece within the Quercetea ilicis vegetation zone, the natural growing zone of *Q. coccifera*, can be divided into two broad groups defined primarily by their geographic distribution. The first group comprises the *Q. coccifera* shrublands of continental as well as insular Greece apart from Crete, while the second group comprises those of Crete.

The first group was classified under the Querco cocciferae–Pistacietum lentisci while the second group under the Rhamno lycoidis–Quercetum cocciferae. The two associations are clearly separated along the first ordination axis of the DCA analysis (Fig. 2), where the former occupies the left part and the latter the right part of the axis. The ordination diagram suggests that the relevés collected from Akarnanika, which is the region with the highest amount of annual rainfall among the study areas, are plotted on the far left part of the axes, while those collected from Crete, and especially central and eastern Crete, which are the driest areas, are plotted on the far right part of the axis. This indicates that the main variation in species composition of the *Q. coccifera* shrublands of Greece, as expressed by the identification of two different associations, is related primarily to differences in rainfall and consequently to soil moisture availability.

Querco cocciferae–Pistacietum lentisci was recorded almost in the entire continental Greece and on most islands, with a considerably lower occurrence on Crete. The association has a distribution mainly in the subhumid Mediterranean climatic zone characterized by a mean altitude of 500 m. Occasionally, it is found beyond these limits exploiting gaps created in local communities by anthropogenic activities. In those cases, the association is enriched with species of the upper vegetation of thermophilous deciduous oaks Quercetea pubescens. The percentage cover in the shrub layer is mainly a function of the local environmental conditions and especially of the local anthropogenic pressures: dense in depressions and scarce around settlements and areas where grazing is intense.

According to our results throughout Greece, *P. lentiscus* and *Ph. latifolia* coexist within the Quercetea ilicis, apart from Crete, and are among the most common accompanying species in the *Q. coccifera* shrublands. Thus, the association Querco cocciferae–Phillyreum mediae (latifoliae) described, among others, by Barbero and Quézel (1976); Bergmeier (1990) and Dimopoulos (1993) for different regions of Greece is in fact similar to the Querco cocciferae–Pistacietum lentisci association described here. Mavrommatis (1980) classifies the same formations in Greece under the Oleo–Lentisctum, while Horvat et al. (1974) identified Coccifereto–Lentisctum in the warmer areas of Eastern, Central, and Southern Greece, as well as in Southern Chalkidiki and in the island of Thassos in Northern Greece.

The wide distribution of the association results in a number of sub-associations reflecting local peculiarities. The five identified sub-associations are clearly separated along the second ordination axis of the DCA analysis (Fig. 2). The lower part of the axis is occupied by Querco cocciferae–Pistacietum lentisci typicum which, as already noted, occurs on areas which are relatively less degraded by intense human presence. On the upper part of the axis, on the other hand, the relevés from the island of Lesvos are plotted, which are the most degraded among the studied areas, forming the Querco cocciferae–Pistacietum lentisci euphorbiotsum rigidiae. The relevés collected from the rest of the areas are plotted in between the above two forming three additional sub-associations occurring on areas with intermediate level of human pressure. Thus, the second axis appears to be representing a gradient of human induced degradation.

The significant effect of human pressure led many researchers to suggest that *Q. coccifera* shrublands are of secondary nature, mainly due to grazing and coppicing for firewood production that has led to the degradation of *Pinus halepensis* or *Pinus brutia* forests (e.g., Horvat et al. 1974; Barbero and Quezel 1976; Mavrommatis 1980; Konstantinidis 1990; Bergmeier 1990). According to our results, human-induced degradation can cause differentiations in *Q. coccifera* shrublands allowing some elements of the neighboring flora to participate and often alter both the physiognomy and the floristic composition of the shrublands.

Rhamno lycoidis–Quercetum cocciferae association is found mainly on Crete as a result of the drier conditions prevailing on most part of the island compared to the rest of the country. The association is floristically enriched by a significant number of species characteristic of semidry and dry environments and also grazing resistant species, such as species characteristic of the abundant phryganic ecosystems in the island (Tsiourlis et al. 2007). An important differentiation from the widespread previous association is its altitudinal distribution: almost from the sea level to almost the timber line (1,600 m asl; Tsiourlis et al. 2001). The Cretan forests are differentiated from the neighboring areas such as Peloponnese and Middle East, by belonging exclusively to the Quercetea ilicis in all the altitudinal range (Barbero et Quézel 1980), while communities from Quercetea pubescens common in the others parts of the country, like *Q. pubescens* and *Q. frainetto* forests (e.g., Horvat et al. 1974; Raus 1980; Konstantinidis et al. 2002) or even *Castanea sativa* forests (e.g., Raus 1980; Bergmeier 1990; Konstantinidis et al. 2008) are absent in Crete (Zaffran 1990). It appears the primary forest of Crete used to be a dense sclerophyllous Mediterranean forest (Greuter 1966). So the *Q. coccifera* shrublands of Crete are degradation forms of these primary forests with *Q. coccifera* accompanied by *Q. ilex*, which currently has only limited local

distribution (Greuter 1966; Zaffran 1990; Kypriotakis et al. 1996). Quézel (1981), also believes that climax plant communities in the Mediterranean zone were not always forests, so well as Kypriotakis et al. (1996) and Tsiorliris et al. (2001) and that applies especially in Crete.

The identification of three sub-associations of the Rhamno lycoidis–Quercetum cocciferae is primarily the result of the west–east gradient of precipitation and, to a lesser extent, the result of the varying degree of degradation, which is severe almost everywhere in Crete.

As already mentioned, the current study is confined within the Quercetea ilicis vegetation zone. However, *Q. coccifera* formations often extend beyond this vegetation zone. Bergmeier (1990) stated that the classification of *Q. coccifera*-dominated vegetation units, from meso-Mediterranean levels up to montane levels, comprise at least two classes being syntaxonomically involved, i.e., Quercetea ilicis and Quercetea pubescens, or even three, if Rhamno–Prunetea is adopted as a class of its own. Karagiannakidou–Iatropoulou (1983) and Chasapis et al. (2004) adopted syntax mentioned by Raus (1980) for E. Thessaly for the pseudomaquis of Mt. Chortiatis (N Greece) and assigned them also to Quercetea pubescens (Ostryo–Carpinion orientalis and Quercion confertae). Thus, it appears that there is a need to conduct an even wider sampling and use of bibliographical relevés beyond Quercetea ilicis in order to achieve a complete classification for the entire *Q. coccifera* shrublands of Greece.

Acknowledgments This research has been carried out in the frame of the project “NATURA 2000” funded by the Hellenic Ministry of Environment, Physical Planning and Public Works, the projects “DeMon,” “GeoRange,” and “DeSurvey” funded by the European Union (DGXII, EESD, ENV4-CT95-0166, EVK2-CT2000-00091, and CT-2003-003950), the project “ATHOS” funded by the National Agricultural Research Foundation of Greece, the project “Survey of apiarian flora of Peloponnesus” funded by the Hellenic Ministry of Rural Development and Food and the project “SITHON” funded by the Greek General Secretary of Research and Development.

References

- Athanasiadis N (1986) Forest phytosociology. Giahoudi-Giapouli, Thessaloniki 110 pp (in Greek)
- Balaguer E, Martínez-Ferri F, Valladares ME, Pérez-Corona FJ, Baquedano FJ, Castillo FJ, Manrique E (2004) Population divergence in the plasticity of the response of *Quercus coccifera* to the light environment. Funct Ecol 15:124–135
- Barbero M, Quézel P (1976) Les groupements forestiers de Grèce centro-meridionale. Ecol Medit II:1–86
- Barbero M, Quézel P (1980) La végétation forestière de Crète. Ecol Medit 5:175–210
- Bergmeier E (1990) Wälder und Gebüsche des niederer Olympos (Kato Olimpos, NO-Thessalien). Ein Beitrag zur systematischen und orographischen Vegetationsgliederung Griechenlands. Phytocoenologia 18:161–342
- Bergmeier E, Dimopoulos P (2001) *Fagus sylvatica* forest vegetation in Greece: Syntaxonomy and gradient analysis. J Veg Sci 12:109–126
- Boyko H (1954) A new plant—geographical subdivision of Israel (as an example for southwest Asia). Plant Ecol 5–6(1):309–318
- Braun-Blanquet J (1924) Études sur la végétation méditerranéenne. III. Concentration en ions H et calcimétrie du sol de quelques associations de la garigue languedocienne. Bull Soc Bot France 71:639–647 879–891
- Braun-Blanquet J (1936) La chênaie d’yeuse méditerranéenne (Quercion ilicis). Mém Soc Étude Sci Nat Nîmes 5:3–147
- Braun-Blanquet J (1947) Les groupements végétaux supérieurs de la France. In: Braun-Blanquet J, Emberger L, Molinier R (eds) Instructions pour l’établissement de la Carte des Groupements Végétaux. SIGMA, Montpellier, pp 19–32
- Braun-Blanquet J (1964) Pflanzencoziologie. Springer, Heidelberg
- Braun-Blanquet J, Bolòs O (1957) Les groupements végétaux du Bassin Moyen de l’Ebre et leur dynamisme. Anales de la Estación Experimental de Aula Dei 5:1–266
- Braun-Blanquet J, de Bolòs O (1954) Datos sobre las comunidades terofíticas de las llanuras del Ebro medio. Collect Bot (Barcelona) 4 (2):235–242
- Braun-Blanquet J, Font Quer P, Braun-Blanquet G, Frey E, Jansen P, Moor M (1936) L’Excursion de la Sigma en Catalogne (Pâques 1934). Cavanillesia 7:153–167
- Cañelas I (1993) Ecología, Características y usos de los coscojares (*Quercus coccifera* L.) en España. Thesis Doctoral. Universidad Politécnica De Madrid
- Chasapis M, Karagiannakidou V, Theodoropoulos K (2004) Phyto-sociological research of *Quercus coccifera* L. pseudomaquis on Mount Chortiatis, northern Greece. Isr J Plant Sci 52, 4:357–381
- Costa JC, Aguiar C, Capelo J, Lousã M, Neto C (1998) Biogeografia de Portugal continental. Quercetea 0:5–56
- Dafis S (1970) Forest phytosociology. Aristotle University of Thessaloniki, Thessaloniki (in Greek)
- Dafis S (1973) Taxonomy of forest vegetation of Greece. Ann Fac Forest and Envir AUTH 15(2):75–88 In Greek
- Dafis S, Jahn G (1975) Zum heutigen Waldbild Griechenlands nach ökologisch-pflanzengeographischen Gesichtspunkten. Veröff Geopot Insit ETH. Stiftung Rübel Zürich 55:99–116
- Davis PH (1965–1985) Flora of Turkey and the East Aegean Islands. Vol. 1–9. Edinburgh Univ. Press, Edinburgh
- Debazac EF (1969) La végétation de certaines stations de l’Oleo-Ceratonion en Attique. Biologia Gallo-Hellenica II 1:3–12
- Diapoulis C (1939–1949) Flora of Greece, vols 1–3. Ministry of Agriculture, Athens In Greek
- Dimopoulos P (1993) Floristic and phytosociological research of Mountain Killini: an ecological approach. Ph D Thesis. Univ of Patras, Patras, 370 p. (in Greek)
- Edmondson JR (1982) *Globularia* L. In: Davis PH (ed) Flora of Turkey and the East Aegean Islands, vol 7. University Press, Edinburgh
- Gauch HG (1982) Multivariate analysis in community ecology. Cambridge University Press, London and New York, p 574
- Greuter W (1966) Kreta, landschaft und pflanzenwelt. Mitt Naturf Ges, Bern 2, 23:31–33
- Greuter W, Burdet HM, Long G (eds) (1986) Med-Checklist. A critical inventory of vascular plants of the circum-mediterranean countries. 3. Dicotyledones (Convolvulaceae-Labiatae). Conservatoire et Jardin Botaniques, Genève
- Grove AT, Ispikoudis I, Kazaklis A, Moody JA, Papanastasis V, Rackham O (1993) Threatened Mediterranean landscapes: West Crete. University of Cambridge, Cambridge
- Hansen A, Nielsen H (1993) Contribution to the Flora of Lesvos (Mitolini), Greece. Willdenowia 23:143–156
- Hill MO (1973) Reciprocal Averaging: An eigenvector method of ordination. J Ecol 61:237–249

- Hill MO (1974) Correspondence Analysis: A neglected multivariate method. *Appl Stat* 23:340–354
- Hill MO (1979) TWINSPAN—a FORTRAN program for arranging multivariate data in an ordered two way table by classification of the individuals and the attributes. Cornell University, Department of Ecology and Systematics, Ithaca
- Hill MO, Bunce RGH, Shaw MW (1975) Indicator Species Analysis, a divisive polythetic method of classification, and its application to a survey of native pinewoods in Scotland. *J Ecol* 63:597–613
- Horvat I, Clavac V, Ellenberg H (1974) Vegetation Südosteuropas. Stuttgart University Press, Stuttgart 660 p
- Kagiaiki A (2003) Fytogeografiki meleti tou toxou tou Notiou Aigaiou. University Of Crete, Department of Biology. MSc. Thesis. Iraklion. 245 p. (In Greek: Phytogeographical study of the South Aegean Island Arc)
- Karaer F, Kiliç M, Kutbay H (1999) The woody vegetation of Kelkit Valley. *Turk J Bot* 23:319–344
- Karagiannakidou-Iatropoulou V (1983) Site Research in Quercoco-Fagetea Class of the Chortiatis Mountain Range. Ph D thesis. Sci Ann Fac Phys An Math AUTH Annex 25, vol. 22. Thessaloniki. 161 p. (in Greek)
- Kavvadas D (1956–1964) Illustrated Botanical–Phytological Lexicon I–IX. Athens. (in Greek)
- Konstantinidis P (1990) The effect of physiography on the plant communities in forest of *Pinus halepensis* M. of Sithonia Peninsula in Halkidiki, N. Greece. Thesis of doctorate. Aristotelian University of Thessaloniki. Thessaloniki, Greece. (In Greek, with English summary)
- Konstantinidis P, Tsiorliris G (2003) Description—analysis and mapping of vegetation units (habitats) of Lagadas County (Thessaloniki, Greece). NAGREF-Forest Research Institute, GeoRange project [DGXII, EESD, EVK2-CT2000-0009]
- Konstantinidis P, Chatziphilippidis G, Tsiorliris GM, Tsiontis A (2002) Taxonomy and ecology of plant communities of *Quercus frainetto* Ten (*Q. conferta* Kit.) forests in Greece. *Isr J Plant Sci* 50:145–154
- Konstantinidis P, Tsiorliris G, Galatsidas S (2005) Effects of wildfire season on the resprouting of kermes oak (*Quercus coccifera* L.). *For Ecol Manag* 208:15–27
- Konstantinidis P, Tsiorliris G, Xofis P, Buckley GP (2008) Taxonomy and ecology of *Castanea sativa* Mill. forests in Greece. *Plant Ecol* 195:235–256
- Kypriotakis Z, Tzanoudakis D, Tsiorliris G (1996) Vegetation map of Crete. Proceedings of the 6th Botanical Scientific Conference, Hellenic Botanical Society and Biological Society of Cyprus Society, Paralimni, Cyprus: 301–306 (in Greek with English abstract)
- Ladero M, Asensi A, Díez-Garretas B, Valle F (2002) Estudios sobre la vegetación del entorno del Balneario de Alhama de Granada (Granada). *Anal Real Acad Nac Farm* 68:100–125
- Liakos L, Moulopoulos C (1967) Contribution to the differentiation of some Kermes oak types. Forest Research Center of N. Greece. Bulletin No 16. Thessaloniki. (in Greek)
- Loidi J, Herrera M, Olano J, Silvan F (1994) Maquis vegetation in the eastern Cantabrian coastal fringe. *J Veg Sci* 5:533–540
- Lossaint P, Rapp M (1971) Répartition de la matière organique, productivité et cycle des éléments minéraux dans des écosystèmes de climat méditerranéen. In: Duvigneaud P (ed) Production des écosystèmes forestiers, Ecologie et Conservation vol 4. UNESCO, Paris, pp 587–617
- Martínez-Parras MT, Peinado M, Alcaraz F (1985) Sobre la vegetación termófila de la cuenca mediterránea de Granada y sus áreas limítrofes. *Lazanoa* 8:251–268
- Mavrommatis G (1980) Phytogeographical, phytosociological and bioclimatological importance of the distribution area of kermes oak (*Quercus coccifera* L.) in Greece. *TO DASOS* 90: 72–77. (in Greek)
- Mavrommatis G (1982) La signification phytogéographique, phytosociologique et bioclimatique de l'aire et des limites de l'expansion du chêne Kermès (*Quercus coccifera* L.) en Grèce. Sylviculture under extreme ecological and economic conditions IUFRO. División 1:477–481
- Oberdorfer E (1948) Gliederung und Umgrenzung der Mittelmeervegetation auf der Balkanhabbinsel. *Ber. Geobot. Forschungsinst. Rübel*. Zürich: 48–111
- Ozenda P (1964) Biogéographie végétale. Ed Doins et Cie, Paris
- Pandža M (2004) Vegetation of Phoenician Juniper Macchia–Pistacio lentisci–Juniperetum phoeniceae Trinajstić 1987 (Oleo-ceratonion Br.BI.) on the island of Murter and the small surrounding islands. *Nat Croatia* 13(3):201–212
- Papachristou T (1998) Forest ecosystems of Greece and their relation to grazing animals. In J Humphrey, R Gill, J Claridge, eds, Grazing as a Management Tool in European Forest Ecosystems. Forestry Commission Technical Paper vol. 25: 72–78
- Pavlides G (1982) Geobotanical study of Verticos mountain range. Thessaloniki. 173 p
- Platis PD, Papanastasis VP (2003) Relationship between shrub cover and available forage in Mediterranean shrublands. *Agrofor Syst* 57:59–67
- Poissonet P, Romane F, Thiault M, Trabaud L (1978) Evolution d'une garrigue de *Quercus coccifera* L. soumise à divers traitements: Quelques résultats des cinq premières années. *Vegetatio* 38 (3):135–142
- Poissonet J, Poissonet P, Thiault M (1981) Development of flora, vegetation and grazing value in experimental plots of a *Quercus coccifera* garrigue. *Vegetatio* 46:93–104
- Quézel P (1981) Floristic composition and phytosociological structure of sclerophyllous matorral around the Mediterranean. In: Di Castri F, Goodall DW, Specht RL (eds) Ecosystems of the world 11. Elsevier, Amsterdam, pp 107–121
- Rambal S (1984) Water balance and pattern of root water uptake by a *Quercus coccifera* evergreen scrub. *Oecologia* (Berlin) 62:18–25
- Raus T (1980) Die Vegetation Ostthessaliens (Griechenland). II. Quercoco-Fagetea und azonale Gehölzgesellschaften. *Bot Jahrb Syst* 101:313–361
- Raus T (1982) Phytoogeographical circumscription of the Mediterranean area on the Balkan peninsula and the problem of Thessalian pseudomaquis. *Ecol. Mediterranea* VIII:197–201
- Rechinger KH (1943) Flora Aegea. Springer, Wien
- Rivas Goday S (1959) Contribución al estudio de la Quercetea ilicis hispanica. *Anal Inst Bot A J Cavanilles* 17(2):285–406
- Rivas-Martínez S (1975) La vegetación de la clase Quercetea ilicis en España y Portugal. *Anales Inst Bot Cavanilles* 31(2):205–259
- Rivas-Martínez S, Fernández-González F, Loidi J, Lousá M, Penas A (2001) Syntaxonomical checklist of vascular plant communities of Spain and Portugal to association level. *Itineraria Geobotanica* 14:5–341
- Rivas-Martínez S, Díaz ET, Fernández-González F, Izco J, Loidi J, Lousá M, Penas A (2002) Vascular plant communities of Spain and Portugal addenda to the syntaxonomical checklist of 200. *Itineraria Geobotanica* 15(1–2):5–922
- Strid A (1996) The Greek mountain flora, with special reference to the Central European element. *Bocconeia* 5:99–112
- Terradas J (1999) Holm oak and holm oak forest: an introduction. In: Rodà F, Retana J, Gracia CA, Bellot J (eds) Ecology of Mediterranean Evergreen Oak Forests. Springer, Berlin, pp 3–14
- Theodoropoulos K (1991) Classification of the phytocociological units in Taxiarchis University forest, Chalkidiki. Ph D Thesis. Sci Ann Depart Forest Nat Environ AUTH, LB'. Annex 18. Thessaloniki. 200 pp. (in Greek)

- Trabaud L (1987) Fire and survival traits of plants. In: Trabaud L (ed) The role of fire in ecological systems. SPB, The Hague, pp 65–89
- Trichas A (1996) Oikologia kai viogeographia ton edaphikon koleopteron sto Notio Aigao, me emfasi sti sinthesi, epochiaki kai viotopiki diaforopoisi kai zoogeographia ton oikogeneion *Carabidae* kai *Tenebrionidae*. Ph D Thesis, University of Crete, 477p. (In Greek with English summary: Ecology and biogeography of the ground Coleoptera in the South Aegean area (with emphasis on the composition, seasonal and biotopic differentiation and zoogeography of the families Carabidae and Tenebrionidae))
- Trinajstic I (1987) Fitocenolo kotipolo ka analiza sastojine makije somine *Juniperus phoenicea* L. u Hrvatskom primorju. Acta Bot Croat 46:115–121
- Tsiourlis GM (1992) Etude d'un écosystème de maquis à *Juniperus phoenicea* L. (Naxos, Cyclades, Grèce): structure spatiale et phytomasse des appareils racinaires. Bulletin d'Ecologie (new name: Ecologie) t 23(1-2):71–81
- Tsiourlis GM, Kasapidis P, Konstandinidis P (2001) The role of grazing on the maintain and degradation of Mediterranean Ecosystems in Central Crete, Greece. In K Radoglou, editor, Proceedings of the Int. Conf. Forest Research: a challenge for an integrated European approach, vol. 2: 715–721
- Tsiourlis G, Konstantinidis P, Xofis P (2007) Taxonomy and ecology of phryganic communities with *Sarcopoterium spinosum* (L.) Spach of the Aegean (Greece). Isr J Plant Sci 55:15–34
- Tutin TG, Heywood VH, Burges NA, Valentine DH, Walters SM, Webb DA (1964) Flora Europea vols 1–5. Cambridge University Press, Cambridge
- Vardakakis E, Pavlides G, Lavrentiades G (1987) On the vegetation of a typical xerorthent soil of Polygyros area (SE of Thessaloniki). Feddes Repertorium 98(3–4):253–264
- Vardinogianni K (1994) Biogeographia ton chersaion malakion sto nisiotiko Aigiako toxo. PhD Thesis, University of Athens, 285 p. [In Greek with English summary: Biogeography of the terrestrial mollusks in the Aegean island arc]
- Vogiatzakis IN, Griffiths GH, Mannion AM (2003) Environmental factors and vegetation composition, Lefka Ori massif, Crete. S. Aegean. Global Ecol Biogeogr 12:131–146
- Weber HE, Moravec J, Theurillat JP (2000) International Code of Phytosociological Nomenclature. 3 rd edition. J Veg Sci 11:739–768
- Wraber M (1952) Zum Wasserhaushalt von *Quercus coccifera*. Vegetatio 3:266–278
- Zaffran J (1990) Contributions à la flore et à la vegetation de la Crète. Université de Provence Aix, Marseille 615 p
- Zohary M (1973) Geobotanical foundations of the Middle East, vol 1, 2. Gustav Fisher Verlag, Stuttgart