

Development of accurate solutions for a classical oscillator

Paolo Amore^{a,*}, Nestor E. Sanchez^b

^a*Facultad de Ciencias, Universidad de Colima, Bernal Díaz del Castillo 340, Colima, Colima, Mexico*

^b*Department of Mechanical Engineering, The University of Texas at San Antonio, San Antonio, TX 78249, USA*

Received 7 November 2005; received in revised form 17 August 2006; accepted 18 August 2006

Available online 10 October 2006

Abstract

We present a method to obtain arbitrarily accurate solutions for conservative classical oscillators. The method that we propose here works both for small and large nonlinearities and provides simple analytical approximations. A comparison with the standard Lindstedt–Poincaré method is presented, from which the advantages of our method are clear.

© 2006 Elsevier Ltd. All rights reserved.

1. Introduction

In this paper, we consider the problem of calculating the periodic solutions to the differential equation

$$m \frac{d^2x}{dt^2} + f(x) = 0. \quad (1)$$

Eq. (1) is the Newton equation for a particle of mass m moving under the action of a force $f(x)$. We assume that the system is conservative and therefore we introduce the potential $V(x) = -\int f(x) dx$. The exact solution of Eq. (1) is possible only in a few cases: more often one only disposes of approximate results, which usually are obtained by applying perturbative methods. Perturbation methods are based on an expansion in some small parameter in the problem and the approximate solutions are therefore obtained in the form of a polynomial in such a parameter. This is the case of the Lindstedt–Poincaré method and of the multiple-scale method, which are widely used in the community.

Unfortunately the validity of the perturbative approaches is restricted to the domain of small parameters and the series obtained in such a manner have a finite radius of convergence: in other words these series become useless when the parameters are larger than the radius of convergence. While other techniques have been developed in the literature to deal with this problem, see for example Refs. [1,2], we wish to present a novel method which was recently devised by one of us [3–5] and which allows one to calculate accurate analytical solutions for a classical oscillator, described by Eq. (1). The method is based on the powerful ideas of the linear delta expansion (LDE) [6,7], which allows to obtain, to any given order, fully *analytical* and

*Corresponding author. Tel.: +52 312 316 100; fax: 52 312 316 1135.

E-mail address: paolo@ucol.mx (P. Amore).

extremely accurate expressions. Since our method is not based on an expansion in a small parameter, the series obtained converges regardless of the values taken by the parameter itself.

Since our method is both systematic and analytic it provides several advantages with respect to other techniques which also claim to work for arbitrary functions $f(x)$: this is the case for example of the method of He [8], whose shortcomings have been evidenced by Sanchez in Ref. [9].

2. The method

This problem has been previously studied by Amore and collaborators in Refs. [3,4], where a nonperturbative method which allows to calculate the period of the oscillations was devised.

We will now briefly review the method and then show how to generalize it to calculate the solutions. For a conservative system the total energy E is constant and the period is simply given by

$$T = \int_{x_-}^{x_+} \frac{\sqrt{2}}{\sqrt{E - V(x)}} dx, \quad (2)$$

where x_{\pm} are the inversion points, obtained by solving the equation $E = V(x_{\pm})$. Notice that $V(x) = -\int f(x) dx$ is the potential.

As discussed in Ref. [3] the problem of calculating this integral can be converted to

$$T_{\delta} = \int_{x_-}^{x_+} \frac{\sqrt{2}}{\sqrt{E_0 - V_0(x) + \delta[E - E_0 - V(x) + V_0(x)]}} dx, \quad (3)$$

where $V_0(x)$ is a potential chosen to interpolate the original one. It is expected that $V_0(x)$ is simpler than $V(x)$. For $\delta = 1$ Eq. (3) reduces to Eq. (2), whereas for $\delta = 0$ this formula yields the period of oscillation in the potential $V_0(x)$. We will treat the term proportional to δ as a perturbation and expand in powers of δ . Since $V_0(x)$ depends upon one or more arbitrary parameters (which we will indicate with λ) a residual dependence upon these parameters shows up in the period when the expansion is carried out to a finite order. In order to eliminate such unnatural dependence we impose the principle of minimal sensitivity (PMS) [10] by requiring that

$$\frac{\partial T}{\partial \lambda} = 0. \quad (4)$$

We now proceed to define

$$\Delta(x) = \frac{E - E_0 - V(x) + V_0(x)}{E_0 - V_0(x)}. \quad (5)$$

and write T_{δ} as

$$T_{\delta} = \sum_{n=0}^{\infty} \frac{(2n-1)!!}{n!2^n} (-1)^n \delta^n \int_{x_-}^{x_+} \frac{\sqrt{2}(\Delta(x))^n}{\sqrt{E_0 - V_0(x)}} dx, \quad (6)$$

provided that the series converges uniformly, which is the case if $|\Delta(x)| < 1$ for every x , $x_- \leq x \leq x_+$. We can choose $V_0(x)$ so that each integral in Eq. (6) can be performed, thus obtaining an equivalent series representation for the original integral of Eq. (2).

Amore and collaborators have applied this method to obtain very precise analytical approximations for the period of several oscillator, showing that the error decreases exponentially with the order of the approximation.¹

In particular they considered the Duffing oscillator, which corresponds to the potential $V(x) = \frac{1}{2}x^2 + (\mu/4)x^4$. The interpolating potential was chosen to be $V_0(x) = [(1 + \lambda^2)/2]x^2$, where λ is an arbitrary parameter. Working to first order they obtained the optimal value $\lambda_{\text{PMS}} = \sqrt{3\mu A/2}$ and the

¹Recently, Amore and Arceo have given in Ref. [5] a precise estimate for the rate of convergence of series (6) and they have shown that it converges in all the physical region.

simple formula:

$$T_{PMS} = \frac{4\pi}{\sqrt{4 + 3\mu A^2}}, \tag{7}$$

which provides an error less than 2.2% to the exact period for any value of μ and A .

Physically the PMS allows us to obtain the best potential around which to perform our expansion: the form of the potential depends upon the arbitrary parameter λ and will in general depend on the order to which the calculation is made.

We will now generalize the results of Refs. [3,4] by using the conservation of the energy to obtain the solution to Eq. (1). It is straightforward to see that the time spent by the body to go from x_- to a point X ($x_- \leq X \leq x_+$) is given by

$$t = \int_{x_-}^X \frac{\sqrt{2}}{\sqrt{E - V(x)}} dx. \tag{8}$$

After repeating the procedure previously explained we have

$$t = \sum_{n=0}^{\infty} \frac{(2n - 1)!!}{n!2^n} (-1)^n \delta^n \int_{x_-}^X \frac{\sqrt{2}(\Delta(x))^n}{\sqrt{E_0 - V_0(x)}} dx. \tag{9}$$

The optimal parameter λ will again be chosen according to the same criteria previously adopted. Notice that, once the integrals in Eq. (9) are calculated, one obtains a convergent series representation for the time t as a function of the position X : having proved in Ref. [3] the convergence of Eq. (9) for $X = x_+$, the convergence for $X < x_+$ follows.

As an application we consider the Duffing oscillator and working to first order we obtain

$$t = \frac{(6\mu A^2 + 8)\cos^{-1}(X/A) - A\mu X\sqrt{1 - (X^2/A^2)}}{(3\mu A^2 + 4)^{3/2}} \tag{10}$$

or equivalently

$$\Omega_{PMS}t = \cos^{-1}\left(\frac{X}{A}\right) - \frac{A\mu X\sqrt{1 - (X^2/A^2)}}{6\mu A^2 + 8}. \tag{11}$$

In Fig. 1 we have compared the numerical solution corresponding to $\mu A^2 = 10^4$ with the approximation given by Eq. (11): our simple analytical formula provides an excellent approximation, even in presence of a




Fig. 1. Solution corresponding to Eq. (10) for $\mu A^2 = 10^4$ (solid curve). The dashed curve is numerical.

huge nonlinearity. This regime is clearly outside the region of applicability of perturbative methods, such as the Lindstedt–Poincaré (LP) method.

To ease the comparison with the LP method we notice that the solution to the nonlinear equation will have the general form

$$X(t) = \sum_{n=0}^{\infty} c_n \cos[(2n + 1)\Omega t]. \tag{12}$$

We can obtain the Fourier coefficients c_n simply by using Eq. (11):

$$\begin{aligned} c_n &= -\frac{2}{\pi} \int_{-A}^A X \frac{dt}{dX} \cos[(2n + 1)\Omega_{\text{PMS}} t] dX \\ &= -\frac{2}{\pi} \int_{-A}^A X \left[\frac{-7\mu A^2 + 2\mu X^2 - 8}{2A(3\mu A^2 + 4)\sqrt{1 - (X^2/A^2)}} \right] \\ &\quad \times \cos \left\{ (2n + 1) \left[\cos^{-1} \left(\frac{X}{A} \right) - \frac{A\mu X \sqrt{1 - (X^2/A^2)}}{6\mu A^2 + 8} \right] \right\} dX. \end{aligned} \tag{13}$$

Eq. (13) cannot be evaluated analytically in its present form. However, we notice that the function

$$\xi \equiv -\frac{A\mu X \sqrt{1 - (X^2/A^2)}}{6\mu A^2 + 8} \tag{14}$$

fulfills the constraint $|\xi| \leq -\frac{1}{12}$ and therefore can be used as an expansion parameter.

Working to order ξ^3 we obtain

$$c_0^{(\text{PMS})} = \frac{A(26449\mu^3 A^6 + 107456\mu^2 A^4 + 145408\mu A^2 + 65536)}{1024(3\mu A^2 + 4)^3}, \tag{15}$$

$$c_1^{(\text{PMS})} = \frac{A^3\mu(6435\mu^3 A^6 + 26424\mu^2 A^4 + 36096\mu A^2 + 16384)}{2048(3\mu A^2 + 4)^4}, \tag{16}$$

$$c_2^{(\text{PMS})} = \frac{5A^5\mu^2(427\mu^2 A^4 + 1112\mu A^2 + 768)}{6144(3\mu A^2 + 4)^4}, \tag{17}$$

$$c_3^{(\text{PMS})} = \frac{49A^7\mu^3(5\mu A^2 + 16)}{12288(3\mu A^2 + 4)^4}, \tag{18}$$

...

Notice that the coefficients above are rational functions of μA^2 and all tend to finite values for $\mu A^2 \rightarrow \infty$. For $\mu A^2 \ll 1$ the perturbative expressions are found:

$$c_0^{(\text{PMS})} \approx A \left[1 - \frac{1}{32}(\mu A^2) + \frac{23}{1024}(\mu A^2)^2 - \frac{1055}{65536}(\mu A^2)^3 + \dots \right], \tag{19}$$

$$c_1^{(\text{PMS})} \approx A \left[\frac{\mu A^2}{32} - \frac{51\mu^2 A^4}{2048} + \frac{1287\mu^3 A^6}{65536} + \dots \right], \tag{20}$$

$$c_2^{(\text{PMS})} \approx A \left[\frac{5}{2048}(\mu A^2)^2 - \frac{745}{196608}(\mu A^2)^3 + \dots \right], \tag{21}$$

$$c_3^{(PMS)} \approx A \left[\frac{49}{196608} (\mu A^2)^3 + \dots \right], \tag{22}$$

...

which in part reproduce the results obtained with the LP method:

$$\begin{aligned} X^{(LP)}(t) = & A \left(1 - \frac{\mu A^2}{32} + \frac{23\mu^2 A^4}{1024} - \frac{547\mu^3 A^6}{32768} + \dots \right) \cos[\Omega_{LP}t] \\ & + A \left(\frac{\mu A^2}{32} - \frac{3\mu^2 A^4}{128} + \frac{29\mu^3 A^6}{16384} + \dots \right) \cos[3\Omega_{LP}t] \\ & + A \left(\frac{\mu^2 A^4}{1024} - \frac{3\mu^3 A^6}{2048} + \dots \right) \cos[5\Omega_{LP}t] \\ & + A \left(\frac{\mu^3 A^6}{32768} + \dots \right) \cos[7\Omega_{LP}t] + \dots \end{aligned} \tag{23}$$

It is useful to consider the exact solution to the Duffing equation which, given the initial conditions used can be cast in the form

$$X^{(exact)}(t) = A \operatorname{cn} \left(\sqrt{1 + \mu A^2} t \left| \frac{A^2 \mu}{2(\mu A^2 + 1)} \right. \right). \tag{24}$$

We can use Eq. (16.23.2) of Ref. [11] to write

$$\operatorname{cn}(u|m) = \frac{2\pi}{\sqrt{m}K(m)} \sum_{n=0}^{\infty} \frac{q^{n+1/2}}{1 + q^{2n+1}} \cos(2n + 1)v, \tag{25}$$

where $q \equiv e^{-\pi K(1-m)/K(m)}$ and $v \equiv \pi u/2K(m)$.

We can easily read off Eq. (25) the Fourier coefficients of the cn function and compare them with the approximations Eqs. (15)–(18): the leading Fourier coefficient is reproduced with a maximum error of 0.15%. In Fig. 2 we plot the absolute value of the error $\Xi = \left(\frac{c_0^{approx} - c_0^{exact}}{c_0^{exact}} \right) \times 100$. We notice that for $\mu A^2 < 1$ the error is quite small and dies exponentially fast as the limit $\mu A^2 \rightarrow 0$ is approached; on the other hand, in the limit $\mu A^2 \rightarrow \infty$ the error reaches a plateau. The plateau exists because our approximate coefficient c_0^{approx} has the




Fig. 2. Absolute value of the error over the Fourier coefficient c_0 using the approximate expression (15).

correct asymptotic behaviour, i.e.

$$\lim_{\mu A^2 \rightarrow \infty} \frac{c_0^{\text{approx}}}{c_0^{\text{exact}}} = \frac{26449e^{-\pi/2}(1+e^\pi)\sqrt{\pi/2}}{110592\Gamma(\frac{3}{4})^2} \approx 1.0017. \quad (26)$$

Although the remaining coefficients are reproduced with less accuracy (for the coefficient c_1 we have a maximum 10% of error) the overall solution is very accurate, since c_0 is much larger than all the remaining coefficients. To prove this statement we can read off the exact expressions for the c_n^{exact} from Eq. (25) and calculate

$$R_n = \lim_{\mu A^2 \rightarrow \infty} \frac{c_n^{\text{exact}}}{c_0^{\text{exact}}} = \frac{e^{n\pi}(1+e^\pi)}{1+e^{2\pi n+\pi}}, \quad (27)$$

which decays exponentially for large n . In the case $n = 1$, we see that, even in the asymptotic limit $\mu A^2 \rightarrow \infty$, the coefficient c_0^{exact} is about 22 times larger than c_1^{exact} , which is the key to the precision of our results.

Notice that the limit $\mu A^2 \rightarrow \infty$ corresponds to considering a purely anharmonic oscillator $V(x) = (\mu/4)x^4$: clearly, our method is capable to deal quite efficiently also with this case.

It is worth stressing that the present analysis has been carried out only to first order δ : however, since the method is geometrically convergent (see Ref. [5]) one expects that much higher precision can be obtained by applying it to higher orders, although this issue is not pursued in this paper. We also stress that our method is completely general and that it can be applied to a large class of potentials: a detailed analysis of the application of the method to calculate the period of general oscillators is given in Ref. [4].

3. Conclusions

We have presented a method to obtain arbitrarily accurate solution for a conservative oscillator. The particular technique that we used worked for small and large nonlinearities of the equations, and provides a simple but very accurate approximation. A comparison with the exact solution and with the perturbative solution obtained using the Lindstedt–Poincaré is done. It was shown that errors as small as 0.15% are recorded on the leading Fourier coefficient.

The method provides several advantages over other well established methods in the literature: first of all, to the best of our knowledge our method is the only nonperturbative method which allows to obtain fully analytical results and for which the exponential convergence of the series is proved [3–5]; secondly, previous work done in Refs. [3–5], where the method was used to calculate the period (and not the solution) has shown that our method can be used for a quite large of class of potentials, even in cases where the exact result cannot be obtained; finally it is easy to calculate higher order contributions (*never involving special functions*) with our method.

Finally we wish to point out that the results presented in this paper represent the first application of the nonperturbative method of Refs. [3–5] to calculate the solution of a nonlinear differential equation.

Acknowledgement

This work has been partially supported by CONACYT, grant 40633.

References

- [1] C.W. Lim, Modified Mickens procedure for certain non-linear oscillators, *Journal of Sound and Vibration* 257 (2002) 202–206.
- [2] R.E. Mickens, A generalized iteration procedure for calculating approximations to periodic solutions of truly nonlinear oscillators, *Journal of Sound and Vibration* 287 (2005) 1045–1051.
- [3] P. Amore, R.A. Sáenz, The period of a classical oscillator, *Europhysics Letters* 70 (2005) 425–431.
- [4] P. Amore, A. Aranda, F. Fernández, R. Sáenz, Systematic perturbation of integrals with applications to physics, *Physical Review E* 71 (2005) 016704.
- [5] P. Amore, S. Arceo, Analytical formulas for gravitational lensing, *Physical Review D* 73 (2006) 083004.

- [6] A. Okopińska, Nonstandard expansion techniques for the effective potential in $\lambda\phi^4$ quantum field theory, *Physical Review D* 35 (1987) 1835–1847.
- [7] A. Duncan, M. Moshe, Nonperturbative physics from interpolating actions, *Physics Letters B* 215 (1988) 352–358.
- [8] J.H. He, A new perturbation technique which is also valid for large parameters, *Journal of Sound and Vibration* 225 (2000) 1257–1263.
- [9] N.E. Sanchez, A view to the new perturbation technique valid for large parameters, *Journal of Sound and Vibration* 282 (2005) 1309–1316.
- [10] P.M. Stevenson, Optimized perturbation theory, *Physical Review D* 23 (1981) 2916.
- [11] M. Abramowitz, I.A. Stegun, *Handbook of Mathematical Functions*, Dover Publications, New York, 1965.