

Stieltjes Integrals of Hölder Continuous Functions with Applications to Fractional Brownian Motion

A. A. Ruzmaikina^{1,2}

Received May 17, 1999; final May 18, 2000

We give a new estimate on Stieltjes integrals of Hölder continuous functions and use it to prove an existence-uniqueness theorem for solutions of ordinary differential equations with Hölder continuous forcing. We construct stochastic integrals with respect to fractional Brownian motion, and establish sufficient conditions for its existence. We prove that stochastic differential equations with fractional Brownian motion have a unique solution with probability 1 in certain classes of Hölder-continuous functions. We give tail estimates of the maximum of stochastic integrals from tail estimates of the Hölder coefficient of fractional Brownian motion. In addition we apply the techniques used for ordinary Brownian motion to construct stochastic integrals of deterministic functions with respect to fractional Brownian motion and give tail estimates of its maximum.

KEY WORDS: Fractional Brownian motion; stochastic differential equations; Stieltjes integrals.

1. INTRODUCTION

Fractional Brownian motion (fBm) was first introduced by Kolmogorov in 1940⁽¹⁾ and later studied by Lévy and Mandelbrot.^(2,3)

Let (Ω, \mathcal{F}, P) be a probability space, and $\alpha \in \mathbb{R}$, $|\alpha| < 1$ be a parameter. fBm with exponent α is a self-similar, centered Gaussian random process $\xi_\alpha(t, \omega)$, $(t, \omega) \in [0, \infty) \times \Omega$ (often abbreviated as $\xi_\alpha(t)$) with stationary increments and the correlation function

$$E(\xi_\alpha(s), \xi_\alpha(t)) = C(s^{1+\alpha} + t^{1+\alpha} - |s-t|^{1+\alpha})$$
$$C = -\frac{\Gamma(1-\alpha)}{\alpha} \frac{\cos((1+\alpha)/2)\pi}{((1+\alpha)/2)\pi}$$

¹ Department of Physics, Princeton University, Princeton, New Jersey 08544.

² Present address: Department of Mathematics, University of Virginia, Kerchof Hall, Charlottesville, Virginia 22903; e-mail: anastasia@virginia.edu.

For $\alpha = 0$ we recover the ordinary Brownian motion (oBm). From the form of the correlation function of the increments

$$E(d\xi_\alpha(s), d\xi_\alpha(t)) = C\alpha(\alpha + 1) \frac{ds dt}{|s - t|^{1-\alpha}}$$

one can see that fBm is not Markovian for all $\alpha \neq 0$. The trajectories of fBm are almost surely Hölder-continuous with exponent less than $(1 + \alpha)/2$, and not Hölder-continuous with exponent greater or equal to $(1 + \alpha)/2$. The finite-dimensional distributions of fBm are scale-invariant:

$$\forall r, t > 0, \quad r^{-(1+\alpha)/2} \xi_\alpha(rt) \stackrel{\text{dist}}{=} \xi_\alpha(t)$$

The scale-invariance and long-range correlations make fBm important in applications.

In recent years there has been much interest in fBm, see, for example, refs. 4–14. The problem of the construction of stochastic calculus with respect to fBm has been considered in refs. 15–17. The main difficulty is that fBm fails to be a semi-martingale for all $0 < \alpha < 1$.⁽¹⁵⁾ Our goal is to construct the stochastic integrals with respect to fBm and to prove the general existence-uniqueness theorem for solutions of stochastic differential equations (SDE's) with fBm. Due to the strongly non-Markovian nature of fBm, the most natural way to define the stochastic integral is to do it pathwise, for a.e. ω . This brings us to the question of existence of Stieltjes integrals with respect to Hölder continuous functions and of the existence and uniqueness of solutions of ODE's with Hölder continuous forcing (Section 2). The existence of Stieltjes integrals with respect to Hölder continuous functions was established in refs. 18 and 19. Below we use the methods of Renormalization group to prove a formula (Theorem 2) which allows us to estimate the L^∞ -norm of the Stieltjes integral. We then use this estimate to prove the general existence-uniqueness theorem for solutions of ODE's with Hölder continuous forcing. In Section 3 we apply these results to construct the stochastic integrals with respect to fBm and show the existence and uniqueness of solutions of stochastic differential equations with fBm. Previous results in this direction were obtained by refs. 15 and 16. In Section 4 we use Theorem 2 to get estimates on the tails of the stochastic integral with respect to fBm. We also consider the question of existence of fBm stochastic integral of deterministic functions and derive the probability distribution of its maximum. Throughout the paper, $\mathcal{C}^\beta(I)$ denotes the space of Hölder-continuous functions on the interval I with exponent β . We will have many occasions to partition an interval into 2^n sub-intervals of equal size; the i th partition point of the interval under discussion is denoted by s_i^n , and we put $\Delta f(s + s_i^n) = f(s + s_{i+1}^n) - f(s + s_i^n)$ if f is a function defined on the interval.

After this paper was accepted by *Journal of Statistical Physics*, the author learned by personal communication from L. Decreusefond about refs. 26 and 27. In ref. 26 results related to Theorem 1 and Corollary 5.1 of this paper and are proved via a different approach. In ref. 27 results related to Theorem 1 and Lemma 2.1 are proved via a different approach and a solution of a linear stochastic differential equation with fBm: $dX(t) = aX(t) d\zeta_\alpha(t) + bX(t) dt$ is constructed.

2. MAIN THEOREMS

In this section we consider the question of existence of a Stieltjes integral for functions of unbounded variation, give a formula which allows to estimate its upper bound and prove the existence-uniqueness theorem for ordinary differential equations with Hölder continuous forcing.

In applications (such as the construction of the stochastic calculus for fBm), it is often interesting to consider Stieltjes integrals $\int f dg$ for functions of unbounded variation. The difficulty in constructing the integral is that the upper bounds on Riemann sums $\sum |f| |Δg|$ diverge. However this problem can be solved on certain classes of f, g , since, if g oscillates fast enough, the nearby terms in the Riemann sum $\sum f Δg$ may cancel. It is shown in refs. 18 and 19 that the Stieltjes integral exists on certain classes of Hölder continuous functions.

Theorem 1 (Young-Kondurar). Let $f \in \mathcal{C}^\beta(\mathbb{R}), g \in \mathcal{C}^\gamma(\mathbb{R})$. If $\beta + \gamma > 1$, then $\int_0^t f dg$ exists as a Stieltjes integral for all $t > 0$.

Generalizations of Theorem 1 can be found in ref. 20.

The next formula is useful in estimating the L^∞ -norm of the Stieltjes integral and is the main tool used in this paper.

Theorem 2. Let $f, g, \beta,$ and γ be as in Statement 1. Then

$$\int_s^t f(\tau) dg(\tau) = f(s)(g(t) - g(s)) + \sum_{k=1}^\infty \sum_{i=0}^{2^{k-1}-1} Δf(s + s_{2i}^k) Δg(s + s_{2i+1}^k). \quad (1)$$

As far as we know this form of the Stieltjes integral has not appeared before. The idea behind the proof of Theorem 2 is to write a recursion between the Riemann sums on finer partitions of the interval and the Riemann sum on coarser partitions of the interval, very much like in Renormalization group. The same idea is used in ref. 21 to give a new proof of Theorem 1, which amounts to showing that the right hand side of the Eq. (1) does not depend on the sequence of partitions we choose. Below we will need the change of variables formula which is a deterministic analog of the Itô's formula for Brownian motion.

Lemma 2.1. Let $u: [0, \infty) \times \mathbb{R} \rightarrow \mathbb{R}$, $e, f: [0, \infty) \rightarrow \mathbb{R}$, $1 > \gamma > \frac{1}{2}$, $\beta > 1 - \gamma$, $T > 0$. Suppose $f \in C^\beta([0, T])$, $g \in C^\gamma([0, T])$, e is continuous, u is differentiable in t with continuous $\partial u / \partial t$ and twice differentiable in x . For $0 \leq t \leq T$, consider the Stieltjes integral

$$\eta(t) = \eta(0) + \int_0^t e(s) ds + \int_0^t f(s) dg(s).$$

Then for all $0 \leq t \leq T$, $v(t) = u(t, \eta(t))$ is also a Stieltjes integral whose differential is

$$dv = \frac{\partial u}{\partial t} dt + \frac{\partial u}{\partial \eta} d\eta. \quad (2)$$

Note that for $\gamma > \frac{1}{2}$, the change of variable formula is the same as in the case of ordinary calculus. This follows from the fact that the quadratic variation of g is zero and therefore the terms of order $d\eta^2$ are negligible.

Theorems 1 and 2 can be used to prove the following general theorem on the existence and uniqueness of solutions of ordinary differential equations with Hölder continuous forcing.

Theorem 3. Let $b, \sigma: [0, \infty) \times \mathbb{R} \rightarrow \mathbb{R}$, $g \in \mathcal{C}^\gamma(\mathbb{R})$ and $1/2 < \gamma \leq 1$. Suppose b is globally Lipschitz in t and x , and $\sigma \in \mathcal{C}^1(\mathbb{R})$ with $\sigma, \sigma'_t, \sigma'_x$ globally Lipschitz in t and x . Then for every $T > 0$ and $\gamma > \beta > 1 - \gamma$, the ordinary differential equation

$$dx(t) = b(t, x(t)) dt + \sigma(t, x(t)) dg(t), \quad x(0) = x_0 \quad (3)$$

has a unique solution in $\mathcal{C}^\beta([0, T])$.

3. APPLICATIONS TO FRACTIONAL BROWNIAN MOTION

The natural way of constructing the fBm stochastic integral $\int_0^t f(\tau, \omega) d\xi_\alpha(\tau, \omega)$ is to define it as a Stieltjes integral for a.e. ω . Since $\xi_\alpha(\tau, \omega) \in \mathcal{C}^\gamma(\mathbb{R})$ for $\gamma \geq (1 + \alpha)/2$ with probability one, Theorem 1 implies that the fBm stochastic integral $\int_0^t f(\tau, \omega) d\xi_\alpha(\tau, \omega)$ exists for all $f \in \mathcal{C}^\beta(\mathbb{R})$ with $\beta > (1 - \alpha)/2$. The paper⁽¹⁵⁾ contains a special case of this result for functions $f(\xi_\alpha(\cdot, \omega)) \in \mathcal{C}^1(\mathbb{R})$ for a.e. ω , derived by expanding f in Taylor series in ξ_α and using the fact that the quadratic variation of ξ_α is zero.

Similarly, Theorem 2 holds with probability one for $\int_0^t f(\tau, \omega) d\xi_\alpha(\tau, \omega)$ for all $f \in \mathcal{C}^\beta(\mathbb{R})$ with $\beta > (1 - \alpha)/2$.

Itô's formula for fBm can be stated pathwise, as a corollary of Lemma 2.1, however it can be stated also under weaker assumptions. Itô's formula for fBm has been established under very different assumptions in ref. 16 for $0 < \alpha < 1$, and in ref. 17 for $-1 < \alpha < 1$.

Lemma 3.1. Let $u: [0, \infty) \times \mathbb{R} \rightarrow \mathbb{R}$, $e, f: [0, \infty) \times \Omega \rightarrow \mathbb{R}$, $0 < \alpha < 1$, $\beta > (1 - \alpha)/2$, $T > 0$. Suppose $f \in C^\beta([0, T])$, e is continuous, u is differentiable with continuous $\partial u/\partial t$ and $\partial u/\partial x \in C^\gamma([0, T])$. For $0 \leq t \leq T$, consider the stochastic integral

$$\eta(t, \omega) = \eta(0, \omega) + \int_0^t e(s, \omega) ds + \int_0^t f(s, \omega) d\xi_\alpha(s, \omega).$$

Suppose $\sup_{0 \leq t \leq T} E((\partial^2 u/(\partial \eta)^2)(t, \eta(t, \omega)))^2 < \infty$. Then for all $0 \leq t \leq T$, $v(t) = u(t, \eta(t, \omega))$ is also a stochastic integral whose differential is

$$dv = \frac{\partial u}{\partial t} dt + \frac{\partial u}{\partial \eta} d\eta.$$

Note that for $0 < \alpha < 1$, Itô's formula for fBm is the same as in the deterministic case. This follows from the fact that the quadratic variation of fBm is zero. Theorem 3 implies the following existence-uniqueness theorem for SDE's with fBm:

Theorem 4. Let $b, \sigma: [0, \infty) \times \mathbb{R} \rightarrow \mathbb{R}$, $Z: \Omega \rightarrow \mathbb{R}$ and $0 < \alpha < 1$, $(1 - \alpha)/2 < \beta < (1 + \alpha)/2$. Suppose b is globally Lipschitz in t and x , and $\sigma \in \mathcal{C}^1(\mathbb{R})$ with $\sigma, \sigma'_t, \sigma'_x$ globally Lipschitz in t and x . Then for every $T > 0$ the SDE

$$dX(t, \omega) = b(t, X(t, \omega)) dt + \sigma(t, X(t, \omega)) d\xi_\alpha(t, \omega), \quad X(0, \omega) = Z(\omega) \tag{4}$$

has a unique solution in $C^\beta([0, T])$ with probability 1.

In refs. 15 and 16, the existence and uniqueness theorem for solutions of stochastic differential equations was proved when the diffusion coefficient is a function of time t only; in ref. 15, the existence theorem was proved also when the drift and diffusion coefficients are functions of X only. Both papers adapt the methods used for oBm. The new idea in this paper is to use the formula in Theorem 2, which allows us to prove the existence and uniqueness theorem in the general case, when the drift and diffusion are functions of both t and X .

4. ADDITIONAL RESULTS FOR FRACTIONAL BROWNIAN MOTION

When the integrand is of the form $f(\tau, \xi_\alpha(\tau, \omega))$, we can obtain estimates of the tail of the maximum of stochastic integrals from Theorem 2 and from tail estimates of the Hölder coefficient of fBm.

Theorem 5. Let $f: [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ and $\frac{1}{2} < \gamma < (1 + \alpha)/2$, $\delta = (1 + \alpha)/2 - \gamma$. Suppose f is differentiable with bounded $|f'_t(t, x)|$, $|f'_x(t, x)|$. Write $|f'_t| = \sup_{[0, 1], \mathbb{R}} |f'_t(t, x)|$ and $|f'_x| = \sup_{[0, 1], \mathbb{R}} |f'_x(t, x)|$. Then

$$P\left(\max_{0 \leq t \leq 1} \int_0^t f(\tau, \xi_\alpha(\tau, \omega)) d\xi_\alpha(\tau, \omega) > \lambda\right) \leq \frac{2^\gamma + 1}{2^\gamma - 1} \sqrt{\frac{2}{\pi}} \sum_{n=1}^{\infty} \frac{2^{(1-\delta)n}}{v} \exp\left\{-\frac{2^\gamma - 1}{2^\gamma + 1} v^2 2^{2n\delta - 1}\right\}, \quad (5)$$

where

$$v = \frac{2^{2\gamma-1} - 1}{|f'_x|} \left[\sqrt{\left(|f(0, 0)| + \frac{|f'_t|}{2^{\gamma+1} - 1}\right)^2 + \frac{4}{2^{2\gamma-2}} |f'_x| \lambda} - |f(0, 0)| - \frac{|f'_t|}{2^{\gamma+1} - 2} \right]. \quad (6)$$

More information on the stochastic integral is available when f is a function of t only, since in this case the techniques used for oBm can be applied.

Statement 4.1. Let $0 < \alpha < 1$, and let $f(t, \omega) = f(t)$ be a function of t only. If $f \in L^{2/(1+\alpha)}([0, \infty))$, then the fBm stochastic integral $\int_0^t f(\tau) d\xi_\alpha(\tau, \omega)$ exists in $L^2([0, \infty) \times \Omega)$ for all $t \in [0, \infty)$.

The proof is based on the Hardy–Littlewood–Sobolev inequality (see ref. 22).

Since $\int_0^t f(\tau) d\xi_\alpha(\tau, \omega)$ is a Gaussian process, we can show the following properties:

Statement 4.2. Let α and f be as in Theorem 4.1, and let $0 < \beta < \alpha$. Write $q_f(s, t) = \int_s^t \int_s^t f(u) f(v) (du dv / |u - v|^{1-\alpha})$. If $f \in L^{2/(1+\beta)}([0, 1])$, then

1. for a.e. ω , $\int_0^t f(\tau) d\xi_\alpha(\tau, \omega)$ has a t -continuous version for all $t \in [0, 1]$;

2. for every real $r \in [0, 1]$, $P(\max_{0 \leq t \leq 1} \int_0^t f(\tau) d\xi_\alpha(\tau, \omega) > \lambda)$ is bounded from above by

$$\int_{\lambda r / \sqrt{q_{f_+}(0, 1)}}^{\infty} + \int_{\lambda(1-r) / \sqrt{q_{f_-}(0, 1)}}^{\infty} \sqrt{\frac{2}{\pi}} e^{-x^2/2} dx, \quad \text{where } f_{\pm} = \frac{|f| \pm f}{2};$$

3. for every integer $m \geq 2$ and real $\lambda \geq \sqrt{1 + \log m^4}$, $P(\max_{0 \leq t \leq 1} |\int_0^t f(\tau) d\zeta_\alpha(\tau, \omega)| > \lambda)$ is bounded from above by

$$\int_{\lambda/c}^\infty \frac{5}{2} m^2 e^{-x^2/2} dx$$

where $c = \sup_{0 \leq s, t \leq 1} \sqrt{q_f(s, t)} + (2 + \sqrt{2}) \int_1^\infty \sup_{|s-t| < m^{-x^2}} \sqrt{q_f(s, t)} dx < \infty$.

(1) follows from Kolmogorov’s continuity criterion, the bound (2) follows from Slepian’s lemma,⁽²³⁾ and (3) follows from Fernique’s inequality.⁽²⁴⁾

5. PROOFS OF THEOREMS

5.1. Proof of Theorem 2

Proof. The idea of the proof is to write the Riemann sums on the smaller scales in terms of the Riemann sums on the larger scales as in Renormalization Group. Denote by $S^n(f)$ the Riemann sum of f corresponding to the partition of $[0, 1]$ into 2^n equal sub-intervals. We have

$$\begin{aligned} S^n(f) &= S^{n-1}(f) + \sum_{i=0}^{2^{n-1}-1} \Delta f(s_{2i}^n) \Delta g(s_{2i+1}^n) \\ &\vdots \\ &= S^0(f) + \sum_{k=1}^n \sum_{i=0}^{2^{k-1}-1} \Delta f(s_{2i}^k) \Delta g(s_{2i+1}^k) \end{aligned}$$

As $n \rightarrow \infty$, $S^n(f)$ converges to $\int_0^t f(\tau) dg(\tau)$ by Theorem 1. The right hand side converges provided $\beta + \gamma > 1$. ■

5.2. Proof of the Change of Variables Formula

The proof given here follows ref. 25 for the most part.

Proof. We can write the integral version of Eq. (2):

$$v(t) - v(0) = \int_0^t \frac{\partial u}{\partial s} ds + \int_0^t \frac{\partial u}{\partial \eta} d\eta.$$

From Theorem 2 and from continuity of e it follows that:

$$\Delta\eta(t) = f(t) \Delta g(t) + e(t) \Delta t + O(\Delta t)^{\beta+\gamma} + o(\Delta t).$$

$$v(t) - v(0)$$

$$\begin{aligned} &= \sum_{k \leq [2^{n_t}]} \left\{ u\left(\frac{k}{2^n}, \eta\left(\frac{k}{2^n}\right)\right) - u\left(\frac{k-1}{2^n}, \eta\left(\frac{k}{2^n}\right)\right) \right\} \\ &+ \sum_{k \leq [2^{n_t}]} \left\{ u\left(\frac{k-1}{2^n}, \eta\left(\frac{k}{2^n}\right)\right) - u\left(\frac{k-1}{2^n}, \eta\left(\frac{k-1}{2^n}\right)\right) \right\} \\ &+ u(t, \eta(t)) - u\left(\frac{[2^{n_t}]}{2^n}, \eta\left(\frac{[2^{n_t}]}{2^n}\right)\right) \\ &= \sum_{k \leq [2^{n_t}]} \left\{ \frac{\partial u}{\partial t}\left(\frac{k-1}{2^n}, \eta\left(\frac{k}{2^n}\right)\right) \frac{1}{2^n} + o\left(\frac{t}{2^n}\right) \right\} \\ &+ \sum_{k \leq [2^{n_t}]} \left\{ \frac{\partial u}{\partial \eta}\left[\frac{k-1}{2^n}, \eta\left(\frac{k-1}{2^n}\right)\right] \left(\eta\left(\frac{k}{2^n}\right) - \eta\left(\frac{k-1}{2^n}\right)\right) \right. \\ &\left. + \frac{1}{2} \frac{\partial^2 u}{\partial \eta^2}\left(\frac{k-1}{2^n}, \eta\left(\frac{k-1}{2^n}\right)\right) \left(\eta\left(\frac{k}{2^n}\right) - \eta\left(\frac{k-1}{2^n}\right)\right)^2 + o(\Delta\eta^2) \right\} \\ &= \sum_{k \leq [2^{n_t}]} \left\{ \left(\frac{\partial u}{\partial t}\left(\frac{k-1}{2^n}, \eta\left(\frac{k}{2^n}\right)\right) \right. \right. \\ &\left. \left. + \frac{\partial u}{\partial \eta}\left(\frac{k-1}{2^n}, \eta\left(\frac{k-1}{2^n}\right)\right) e\left[\frac{k-1}{2^n}\right] \right) \frac{1}{2^n} + o\left(\frac{t}{2^n}\right) \right\} \\ &+ \sum_{k \leq [2^{n_t}]} \left\{ \frac{\partial u}{\partial \eta}\left(\frac{k-1}{2^n}, \eta\left(\frac{k-1}{2^n}\right)\right) f\left(\frac{k-1}{2^n}\right) \left(g\left(\frac{k}{2^n}\right) - g\left(\frac{k-1}{2^n}\right)\right) \right. \\ &\left. + \frac{1}{2} \frac{\partial^2 u}{\partial \eta^2}\left(\frac{k-1}{2^n}, \eta\left(\frac{k-1}{2^n}\right)\right) f\left(\frac{k-1}{2^n}\right)^2 \right. \\ &\left. \times \left(g\left(\frac{k}{2^n}\right) - g\left(\frac{k-1}{2^n}\right)\right)^2 + o(\Delta g^2) \right\} \\ &= \int_0^t \left(\frac{\partial u}{\partial s} + \frac{\partial u}{\partial \eta} e(s) \right) ds + o(1) + \int_0^t \frac{\partial u}{\partial \eta} f(s) dg(s) \\ &+ \frac{1}{2} \sum_{k \leq [2^{n_t}]} u_{\eta\eta} f^2 \Delta g^2 + o\left(\sum \Delta g^2\right). \end{aligned}$$

Since $u_{\eta\eta} f^2 \Delta g^2 \leq O(t/2^n)^{2\gamma}$, and $\gamma > \frac{1}{2}$, $\sum_{k \leq [2^{n_t}]} u_{\eta\eta} f^2 \Delta g^2 \rightarrow 0$ as $n \rightarrow \infty$. Similarly, $o(\sum \Delta g^2) \rightarrow 0$ as $n \rightarrow \infty$. ■

5.3. Proof of Theorem 3: Local Existence and Uniqueness

In this section we will prove the local existence and uniqueness result. We will derive the global result in Theorem 3 by showing that we can apply the local existence and uniqueness result repeatedly, taking as an initial condition the value of the solution at the end of the previous interval.

Fix $T > 0$, $s \in [0, T]$ and $a \in \mathbb{R}$. Define an integral operator

$$F_t X = \int_s^t b(\tau, X(\tau)) d\tau + \int_s^t \sigma(\tau, X(\tau)) dg(\tau) + a.$$

For $s = 0$ and $a = x_0$, solutions of the ODE (3) are exactly fixed points of F .

For $0 < \beta < 1$, and for $\varepsilon > 0$, consider the Banach space $\mathcal{C}^\beta([s, s + \varepsilon])$ with the norm $\|f\|_\beta = \max_t |f(t)| + \max_{t_1 \neq t_2} (|f(t_1) - f(t_2)|/|t_1 - t_2|^\beta)$. For $K > 0$, consider the closed subset $\mathcal{C}_K^\beta([s, s + \varepsilon]) = \{f : |f(t_2) - f(t_1)| \leq K |t_2 - t_1|^\beta, \forall t_1, t_2 \in [s, s + \varepsilon]\}$ of $\mathcal{C}^\beta([s, s + \varepsilon])$. Finally, consider the closed subset $\mathcal{C}_K^\beta([s, s + \varepsilon], a) = \{f \in \mathcal{C}_K^\beta([s, s + \varepsilon]) : f(s) = a\}$ of $\mathcal{C}^\beta([s, s + \varepsilon])$. For a given $g \in \mathcal{C}^\gamma([0, T])$, there is $L > 0$ such that $g \in \mathcal{C}_L^\gamma([0, T])$. We will show by the contraction mapping theorem that for given T, s, a, K, L , there exists $\varepsilon > 0$ such that F_t has a unique fixed point on $\mathcal{C}_K^\beta([s, s + \varepsilon], a)$. We need to establish that F_t maps $\mathcal{C}_K^\beta([s, s + \varepsilon], a)$ into itself and that it is a contraction. This will be done in Lemma 5.3. The necessary preliminary estimates are obtained in Corollaries 5.1 and 5.2 of Theorem 2. In what follows we will consider $T, K, L > 0$ to be fixed.

Corollary 5.1. Let β and γ be as in Theorem 1, $T > 0$ and let $g \in \mathcal{C}_L^\gamma([0, T])$. Then for every $s, t \in [0, T]$,

$$\left\| \int_s^t X(\tau) dg(\tau) \right\|_\infty = L \|X\|_\beta (t - s)^\gamma \left(1 + \frac{(t - s)^\beta}{2^{\beta + \gamma} - 2} \right). \tag{7}$$

Proof. By Theorem 2

$$\begin{aligned} & \left| \int_s^t X(\tau) dg(\tau) \right| \\ & \leq |X(s)| |g(t) - g(s)| + \sum_{k=1}^\infty \sum_{i=0}^{2^k - 1 - 1} |\Delta X(s + s_{2i}^k)| |\Delta g(s + s_{2i+1}^k)|. \end{aligned} \tag{8}$$

Since $|X(s)| \leq \|X\|_\beta$, $|\Delta X(s + s_{2i}^k)| \leq \|X\|_\beta ((t - s)/2^k)^\beta$ and $g \in \mathcal{C}_L^\gamma([0, T])$, we obtain after performing the sum in (8),

$$\left| \int_s^t X(\tau) dg(\tau) \right| \leq L \|X\|_\beta (t - s)^\gamma + L \|X\|_\beta \frac{(t - s)^{\beta + \gamma}}{2^{\beta + \gamma} - 2}. \blacksquare$$

Corollary 5.2. Let $\varepsilon > 0$ and let β, γ, g and T be as in Corollary 5.1. Then, for every $s \in [0, T]$ and $t \in [s, s + \varepsilon]$,

$$\left\| \int_s^t X(\tau) d\tau \right\|_{\beta} \leq \|X\|_{\infty} \varepsilon^{1-\beta} (1 + \varepsilon^{\beta}), \quad (9)$$

and

$$\left\| \int_s^t X(\tau) dg(\tau) \right\|_{\beta} \leq L \|X\|_{\beta} \varepsilon^{\gamma-\beta} (1 + \varepsilon^{\beta}) \left(1 + \frac{\varepsilon^{\beta}}{2^{\beta+\gamma}-2} \right). \quad (10)$$

Proof.

$$\begin{aligned} \left\| \int_s^t X(\tau) d\tau \right\|_{\beta} &= \max_{t \in [s, s+\varepsilon]} \left| \int_s^t X(\tau) d\tau \right| + \max_{t_1 \neq t_2 \in [s, s+\varepsilon]} \frac{|\int_{t_1}^{t_2} X(\tau) d\tau|}{|t_2 - t_1|^{\beta}} \\ &\leq \|X\|_{\infty} (\varepsilon + \varepsilon^{1-\beta}). \end{aligned}$$

From Corollary 5.1 we obtain

$$\max_{t \in [s, s+\varepsilon]} \left| \int_s^t X(\tau) dg(\tau) \right| \leq L \|X\|_{\beta} \varepsilon^{\gamma} \left(1 + \frac{\varepsilon^{\beta}}{2^{\beta+\gamma}-2} \right), \quad (11)$$

similarly we obtain

$$\max_{t_1 \neq t_2 \in [s, s+\varepsilon]} \frac{|\int_{t_1}^{t_2} X(\tau) dg(\tau)|}{|t_1 - t_2|^{\beta}} \leq L \|X\|_{\beta} \varepsilon^{\gamma-\beta} \left(1 + \frac{\varepsilon^{\beta}}{2^{\beta+\gamma}-2} \right). \quad (12)$$

The result (10) follows from (11) and (12). ■

Lemma 5.3. Let $b, \sigma: [0, \infty) \times \mathbb{R} \rightarrow \mathbb{R}$, $1 \geq \gamma > 1/2$, $T, K, L > 0$. Suppose b and σ are Lipschitz in x and t . Then there exists $\varepsilon_1 > 0$, such that for all $\gamma > \beta > 1 - \gamma$ and $t \in [s, s + \varepsilon_1]$, the operator F_t maps $\mathcal{C}_K^{\beta}([s, s + \varepsilon_1], a)$ into itself.

Notation. We shall denote the Lipschitz coefficients of b and σ by \mathbf{B} and \mathbf{S} respectively.

Proof. Let $\varepsilon > 0$ and $t_1, t_2 \in [s, s + \varepsilon]$. It is sufficient to demonstrate that $\|F_t X\|_{\beta} \leq K$ for a sufficiently small $\varepsilon > 0$.

By the triangle inequality and by Corollary 5.2,

$$\begin{aligned} \|F_t X\|_\beta &\leq \varepsilon^{\gamma-\beta}(1 + \varepsilon^\beta) \left(\|b(t, X(t))\|_\infty \varepsilon^{1-\gamma} + L \|\sigma(t, X(t))\|_\beta \right. \\ &\quad \left. \times \left(1 + \frac{\varepsilon^\beta}{2^{\beta+\gamma}-2} \right) \right). \end{aligned} \tag{13}$$

Since b and σ are Lipschitz and $X \in \mathcal{C}_K^\beta([s, s + \varepsilon], a)$, it is easy to see that

$$\|b(t, X(t))\|_\infty \leq |b(s, a)| + \mathbf{B}(\varepsilon + K\varepsilon^\beta), \tag{14}$$

and

$$\|\sigma(t, X(t))\|_\beta \leq |\sigma(s, a)| + \mathbf{S}(\varepsilon^{1-\beta} + K)(1 + \varepsilon^\beta). \tag{15}$$

Substituting (14) and (15) into (13), we obtain

$$\begin{aligned} \|F_t X\|_\beta &\leq \varepsilon^{\gamma-\beta}(1 + \varepsilon^\beta) \left[|b(s, a)| + \mathbf{B}(\varepsilon + K\varepsilon^\beta) \varepsilon^{1-\gamma} \right. \\ &\quad \left. + L \left(1 + \frac{\varepsilon^\beta}{2^{\beta+\gamma}-2} \right) (|\sigma(s, a)| + \mathbf{S}(1 + \varepsilon^\beta)(\varepsilon^{1-\beta} + K)) \right]. \end{aligned} \tag{16}$$

Since the right hand side is an increasing continuous function of ε and is 0 at $\varepsilon = 0$, it equals K at some ε_1 . For this choice of ε_1 (or any smaller ε_1), F_t maps $\mathcal{C}_K^\beta([s, s + \varepsilon_1], a)$ into itself. ■

Lemma 5.4. Assume the same hypothesis as in Lemma 5.3. Suppose b is Lipschitz in t and x and $\sigma \in \mathcal{C}^1([0, \infty) \times \mathbb{R})$ with $\sigma'_t(t, x)$, $\sigma'_x(t, x)$ Lipschitz in x . Then there exists $\varepsilon_2 > 0$ such that for all $\gamma > \beta > 1 - \gamma$ and $t \in [s, s + \varepsilon_2]$, the operator F_t is a contraction on $\mathcal{C}_K^\beta([s, s + \varepsilon_2])$.

Notation. We shall denote the Lipschitz coefficient of b by \mathbf{B} and the Lipschitz coefficient of σ , σ'_t , σ'_x by \mathbf{S} .

Proof. We need to show that there exist $\varepsilon_2 > 0$ and $\lambda < 1$ such that for all $t \in [s, s + \varepsilon_2]$ and all $X, Y \in \mathcal{C}_K^\beta([s, s + \varepsilon_2])$,

$$\|F_t X - F_t Y\|_\beta \leq \lambda \|X - Y\|_\beta.$$

By the triangle inequality and by Corollary 5.2,

$$\begin{aligned} \|F_t X - F_t Y\|_\beta &\leq \varepsilon^{\gamma-\beta}(1 + \varepsilon^\beta) \left[\|b(t, X(t)) - b(t, Y(t))\|_\infty \varepsilon^{1-\gamma} \right. \\ &\quad \left. + L \|\sigma(t, X(t)) - \sigma(t, Y(t))\|_\beta \left(1 + \frac{\varepsilon^\beta}{2^{\beta+\gamma}-2}\right) \right]. \end{aligned} \quad (17)$$

We estimate the two terms in (17) separately. Since b is Lipschitz,

$$\|b(t, X(t)) - b(t, Y(t))\|_\infty \leq \mathbf{B} |X(t) - Y(t)| \leq \mathbf{B} \|X - Y\|_\beta. \quad (18)$$

Now we will estimate $\|\sigma(t, X(t)) - \sigma(t, Y(t))\|_\beta$. Since σ is differentiable,

$$\max_t |\sigma(t, X(t)) - \sigma(t, Y(t))| \leq \mathbf{S} \|X - Y\|_\beta. \quad (19)$$

By the fundamental theorem of calculus,

$$\sigma(t, X(t)) - \sigma(t, Y(t)) = (X(t) - Y(t)) \int_0^1 \sigma'_x(t, vX(t) + (1-v)Y(t)) dv.$$

Therefore

$$\begin{aligned} &|\sigma(t_1, X(t_1)) - \sigma(t_1, Y(t_1)) - \sigma(t_2, X(t_2)) + \sigma(t_2, Y(t_2))| \\ &= \left| (X(t_1) - Y(t_1) - X(t_2) + Y(t_2)) \int_0^1 \sigma'_x(t_1, vX(t_1) + (1-v)Y(t_1)) dv \right. \\ &\quad \left. + (X(t_2) - Y(t_2)) \int_0^1 (\sigma'_x(t_1, vX(t_1) + (1-v)Y(t_1)) \right. \\ &\quad \left. - \sigma'_x(t_2, vX(t_2) + (1-v)Y(t_2))) dv \right| \\ &\leq \|X - Y\|_\beta (t_2 - t_1)^\beta \mathbf{S} + \|X - Y\|_\beta (\mathbf{S}(t_2 - t_1) \\ &\quad + \mathbf{S}(v |X(t_1) - X(t_2)| + (1-v) |Y(t_1) - Y(t_2)|)) \\ &\leq \|X - Y\|_\beta (t_2 - t_1)^\beta \mathbf{S}(1 + (t_2 - t_1)^{1-\beta} + K). \end{aligned}$$

Consequently

$$\|\sigma(t, X(t)) - \sigma(t, Y(t))\|_\beta \leq \|X - Y\|_\beta \mathbf{S}(2 + \varepsilon^{1-\beta} + K). \quad (20)$$

Substituting (19) and (20) in (17), we obtain

$$\begin{aligned} \|F_t X - F_t Y\|_\beta &\leq \varepsilon^{\gamma-\beta}(1 + \varepsilon^\beta) \left[\mathbf{B}\varepsilon^{1-\gamma} + \mathbf{L}\mathbf{S}(2 + \varepsilon^{1-\beta} + K) \right. \\ &\quad \left. \times \left(1 + \frac{\varepsilon^\beta}{2^{\beta+\gamma}-2} \right) \right] \|X - Y\|_\beta. \end{aligned} \tag{21}$$

The coefficient of $\|X - Y\|_\beta$ is an increasing function of ε and is 0 at $\varepsilon = 0$. Choose $\varepsilon_2 > 0$ small enough so that this coefficient is less than 1 and $\varepsilon_2 \leq \varepsilon_1$. Then F is a contraction on $\mathcal{C}_K^\beta([s, s + \varepsilon_2])$. ■

Combining Lemmas 5.3 and 5.4, we obtain a local existence and uniqueness result:

Corollary 5.5. Assume the same hypothesis as in Lemma 5.4. Then there exists $\varepsilon > 0$ depending on s such that for all $t \in [s, s + \varepsilon]$, ODE (3) has a unique solution in $\mathcal{C}_K^\beta([s, s + \varepsilon], a)$. In particular, if ODE (3) has a solution X on $[0, s]$, then there exists $\varepsilon > 0$ depending on s such that for all $t \in [s, s + \varepsilon]$, ODE (3) has a unique solution Y in $\mathcal{C}_K^\beta([s, s + \varepsilon], X(s))$.

Proof. Take ε to be ε_2 of Lemma 5.4. Since F_t is a contraction on the closed subset $\mathcal{C}_K^\beta([s, s + \varepsilon], a)$ of the complete metric space $\mathcal{C}^\beta([s, s + \varepsilon])$, it has a unique fixed point X in $\mathcal{C}_K^\beta([s, s + \varepsilon], a)$. From the definition of F_t it follows that X is a unique solution of ODE (3) on $[s, s + \varepsilon]$ in $\mathcal{C}_K^\beta([s, s + \varepsilon], a)$. ■

The sufficient conditions on ε in Corollary 5.5 are given by inequalities (16) and (21) with a replaced by $X(s)$:

$$\begin{aligned} \varepsilon^{\gamma-\beta}(1 + \varepsilon^\beta) \left[|b(s, X(s))| + \mathbf{B}(\varepsilon + K\varepsilon^\beta) \varepsilon^{1-\gamma} \right. \\ \left. + L \left(1 + \frac{\varepsilon^\beta}{2^{\beta+\gamma}-2} \right) (|\sigma(s, X(s))| + \mathbf{S}(1 + \varepsilon^\beta)(\varepsilon^{1-\beta} + K)) \right] \leq K, \end{aligned} \tag{22}$$

$$\varepsilon^{\gamma-\beta}(1 + \varepsilon^\beta) \left(\mathbf{B}\varepsilon^{1-\gamma} + \mathbf{L}\mathbf{S}(1 + \varepsilon^{1-\beta} + K) \left(1 + \frac{\varepsilon^\beta}{2^{\beta+\gamma}-2} \right) \right) < 1. \tag{23}$$

Inequality (23) does not depend on $X(s)$.

5.4. Proof of Theorem 3: Global Existence and Uniqueness

By Corollary 5.5 with $s = 0$ and $a = x_0$, ODE (3) has a unique solution on $[0, \varepsilon_0]$, where ε_0 satisfies (22) and (23) with $s = 0$. By using Corollary 5.5

n times we obtain that the solution exists on $[0, \varepsilon_0 + \dots + \varepsilon_{n-1}]$. ODE (3) has a solution on $[0, T]$ if there exists $m > 0$ such that $\sum_{i=0}^m \varepsilon_i \geq T$. This is true if b and σ are globally bounded, since in this case we can choose $\varepsilon_i = \varepsilon_0$ (to see this we can substitute the global bounds on $b, \sigma, \mathbf{B}, \mathbf{S}$ into (22) and (23)). In the case when b and σ grow at most linearly, we will use a change of variables to reduce it to the case of globally bounded b and σ .

Proof. Existence: Suppose that b and σ are bounded on $[0, \infty) \times \mathbb{R}$, then taking the upper bound on b and σ in (22) we get that ε_i satisfying (22) and (23) does not depend on i . In this case the global existence is established. Now suppose that b and σ satisfy the assumptions of Theorem 3. Consider the ODE

$$dy(t) = \frac{b(t, \tan y(t))}{1 + (\tan y(t))^2} dt + \frac{\sigma(t, \tan y(t))}{1 + (\tan y(t))^2} dg(t) \tag{24}$$

This ODE has globally bounded coefficients satisfying the assumptions of Theorem 3, and thus has a global solution on $[0, T]$. Now, $x(t) = \tan y(t)$ satisfies Eq. (3) (by Lemma 2.1). Thus Eq. (3) has a global solution on $[0, T]$.

Uniqueness: Let Y_1 and Y_2 be two solutions in $\mathcal{C}^\beta([0, T])$. Then there exist K_1 and K_2 such that $Y_1 \in \mathcal{C}_{K_1}^\beta([0, T])$ and $Y_2 \in \mathcal{C}_{K_2}^\beta([0, T])$, so Y_1 and Y_2 are in $\mathcal{C}_{\max\{K_1, K_2\}}^\beta([0, T])$. Y_1 and Y_2 coincide at the initial point $t = 0$. Let t_{sup} be the supremum of the set on which they coincide. Since both solutions are continuous, they coincide at t_{sup} as well. t_{sup} must equal T , for otherwise we can make Y_1 and Y_2 coincide past t_{sup} by Corollary 5.5. ■

5.5. Proof of Itô’s Formula for fBm (Lemma 3.1)

Proof. By analogy with the proof of Lemma 2.1, we obtain

$$\begin{aligned} v(t) - v(0) &= \int_0^t \left(\frac{\partial u}{\partial s} + \frac{\partial u}{\partial \eta} e(s) \right) ds + o(1) + \int_0^t \frac{\partial u}{\partial \eta} f(s) d\xi_\alpha \\ &\quad + \frac{1}{2} \sum_{k \leq [2^n t]} u_{\eta\eta} f^2 \Delta \xi_\alpha^2 + o \left(\sum \Delta \xi_\alpha^2 \right). \end{aligned}$$

Since $E(u_{\eta\eta} f^2 \Delta \xi_\alpha^2) \leq \|f\|_\infty \sqrt{E(u_{\eta\eta})^2} \sqrt{E(\Delta \xi_\alpha^2)} = O(t/2^n)^{2\gamma}$, by Chebyshev inequality

$$P \left(\sum_{k \leq [2^n t]} u_{\eta\eta} f^2 \Delta \xi_\alpha^2 \geq \frac{1}{n} \right) \leq \text{const} \frac{n}{2^{n(2\gamma-1)}},$$

and by the Borel–Cantelli lemma $\sum_{k \leq [2^n t]} u_{\eta\eta} f^2 \Delta \xi_\alpha^2 \rightarrow 0$ as $n \rightarrow \infty$ a.e. An analogous argument shows that $o(\sum \Delta \xi_\alpha^2) \rightarrow 0$ as $n \rightarrow \infty$ a.e. Thus Itô’s formula holds. ■

5.6. Proof of Theorem 5

Proof. Since f is differentiable, and since $\xi_\alpha(\cdot, \omega) \in C^\gamma([0, 1])$ with probability 1, for a.e. ω there exists $L(\omega) > 0$ such that

$$|\Delta f(s_{2i}^k, \xi_\alpha(s_{2i}^k))| \leq |f'_t| \frac{t}{2^k} + |f'_x| \frac{L(\omega) t^\gamma}{2^{k\gamma}}$$

holds. From Theorem 2 we get

$$\left| \int_0^t f(\tau, \xi_\alpha(\tau)) d\xi_\alpha(\tau) \right| \leq |f(0, 0)| L(\omega) t^\gamma + |f'_t| \frac{L(\omega) t^{\gamma+1}}{2^{\gamma+1}-2} + |f'_x| \frac{L(\omega)^2 t^{2\gamma}}{2^{2\gamma}-2}$$

and so,

$$\max_{0 \leq t \leq 1} \left| \int_0^t f(\tau, \xi_\alpha(\tau)) d\xi_\alpha(\tau) \right| \leq |f(0, 0)| L(\omega) + |f'_t| \frac{L(\omega)}{2^{\gamma+1}-2} + |f'_x| \frac{L(\omega)^2}{2^{2\gamma}-2}.$$

Therefore

$$\begin{aligned} &P \left\{ \omega : \max_{0 \leq t \leq 1} \left| \int_0^t f(\tau, \xi_\alpha(\tau)) d\xi_\alpha(\tau) \right| > \lambda \right\} \\ &\leq P \left\{ \omega : |f(0, 0)| L(\omega) + |f'_t| \frac{L(\omega)}{2^{\gamma+1}-2} + |f'_x| \frac{L(\omega)^2}{2^{2\gamma}-2} > \lambda \right\} \\ &\leq P\{ \omega : L(\omega) > \nu \} \\ &\leq P\{ \omega : \exists t_1, t_2 \in [0, 1] \text{ s.t. } |\xi_\alpha(t_2) - \xi_\alpha(t_1)| > \nu |t_2 - t_1|^\gamma \}, \end{aligned}$$

where ν is given by (6).

It is easy to see that if

$$\left| \xi_\alpha\left(\frac{k+1}{2^n}\right) - \xi_\alpha\left(\frac{k}{2^n}\right) \right| \leq \frac{2^\gamma - 1}{2^\gamma + 1} \frac{L(\omega)}{2^{n\gamma}}, \quad \forall n > 0, \quad 0 \leq k \leq 2^n - 1,$$

then

$$|\xi_\alpha(t_2) - \xi_\alpha(t_1)| \leq L(\omega) |t_2 - t_1|^\gamma, \quad \forall t_1, t_2 \in [0, 1].$$

Therefore

$$\begin{aligned}
 & P\{\omega : \exists t_1, t_2 \in [0, 1] \text{ s.t. } |\xi_\alpha(t_2) - \xi_\alpha(t_1)| > \nu |t_2 - t_1|^\gamma\} \\
 & \leq P\left\{\omega : \exists n > 0, 0 \leq k \leq 2^n - 1, \text{ s.t. } \left| \xi_\alpha\left(\frac{k+1}{2^n}\right) - \xi_\alpha\left(\frac{k}{2^n}\right) \right| > \frac{2^\gamma - 1}{2^\gamma + 1} \frac{\nu}{2^{n\gamma}}\right\} \\
 & \leq \sum_{n=1}^{\infty} \sum_{k=0}^{2^n-1} P\left\{\omega : \left| \xi_\alpha\left(\frac{k+1}{2^n}\right) - \xi_\alpha\left(\frac{k}{2^n}\right) \right| > \frac{2^\gamma - 1}{2^\gamma + 1} \frac{\nu}{2^{n\gamma}}\right\} \\
 & \leq \sum_{n=1}^{\infty} 2^n \sqrt{\frac{2}{\pi}} \int_{(2^\gamma-1)/(2^\gamma+1) \nu 2^{n\delta}}^{\infty} e^{-x^2/2} dx.
 \end{aligned}$$

Using the estimate $\int_c^\infty e^{-x^2/2} dx \leq e^{-c^2/2}/c$, we obtain the desired result. ■

5.7. Proof of Statement 4.1

Proof. The proof will be reached via the step-by-step procedure used for oBm. Fix $t > 0$.

Step 1. Let $\phi: [0, \infty) \rightarrow \mathbb{R}$, be a simple function of the form $\sum_{i=0}^{2^n-1} \phi(s_i^n) \chi_{[s_i^n, s_{i+1}^n]}$, where χ is an indicator function and $s_i^n = (i/2^n)t$. Define

$$\int_0^t \phi(\tau) d\xi_\alpha(\tau) = \sum_{i=0}^{2^n-1} \phi(s_i^n) \Delta \xi_\alpha(s_i^n)$$

Step 2. Let $g \in \mathcal{C}^\infty([0, \infty))$. Approximate g by a sequence of simple functions: $\phi_n(\tau) = \sum_{i=0}^{2^n-1} g(s_i^n) \chi_{[s_i^n, s_{i+1}^n]}(\tau)$. Then $\phi_n \rightarrow g$ uniformly on $[0, t]$ and $\int_0^t |g(\tau) - \phi_n(\tau)|^{2/(1+\alpha)} d\tau \rightarrow 0$. Therefore the sequence ϕ_n is Cauchy in $L^{2/(1+\alpha)}([0, \infty))$. To show that the sequence $\int_0^t \phi_n(\tau) d\xi_\alpha(\tau)$ is Cauchy in $L^2([0, \infty) \times \Omega)$, we use the Hardy–Littlewood–Sobolev inequality:

$$\begin{aligned}
 & E\left(\int_0^t \phi_m(\tau) d\xi_\alpha(\tau) - \int_0^t \phi_n(\tau) d\xi_\alpha(\tau)\right)^2 \\
 & = C\alpha(\alpha+1) \int_0^t \int_0^t \frac{(\phi_m(u) - \phi_n(u))(\phi_m(v) - \phi_n(v))}{|u-v|^{1-\alpha}} du dv \\
 & \leq \text{const} \cdot \|\phi_m - \phi_n\|_{2/(1+\alpha)}^2 \rightarrow 0 \quad \text{as } m, n \rightarrow \infty.
 \end{aligned}$$

Thus the integral $\int_0^t g(\tau) d\xi_\alpha(\tau)$ exists as the L^2 -limit of $\int_0^t \phi_n(\tau) d\xi_\alpha(\tau)$.

Step 3. Let $f \in L^{2/(1+\alpha)}([0, \infty))$. Let $j \in \mathcal{C}_c^\infty([0, \infty))$ with $\int_{[0, \infty)} j = 1$. Define $j_n(\tau) = (1/n) j(n\tau)$ and $g_n = j_n * f$. Then $g_n \in \mathcal{C}^\infty([0, \infty))$ and

$\int_0^t |g_n(\tau) - f(\tau)|^{2/(1+\alpha)} d\tau \rightarrow 0$. In particular, the sequence g_n is Cauchy in $L^{2/(1+\alpha)}([0, \infty))$. The Hardy–Littlewood–Sobolev inequality gives

$$E \left(\int_0^t g_m(\tau) d\xi_\alpha(\tau) - \int_0^t g_n(\tau) d\xi_\alpha(\tau) \right)^2 \leq \text{const} \cdot \|g_m - g_n\|_{2/(1+\alpha)}^2 \rightarrow 0 \quad \text{as } m, n \rightarrow \infty.$$

Thus the integral $\int_0^t f(\tau) d\xi_\alpha(\tau)$ exists as an L^2 -limit of $\int_0^t g_n(\tau) d\xi_\alpha(\tau)$.

Thus, for all $f \in L^{2/(1+\alpha)}([0, 1])$, we can choose simple functions ϕ_n converging in $L^{2/(1+\alpha)}$ to f such that the L^2 -limit of $\int_0^t \phi_n(\tau) d\xi_\alpha(\tau)$ exists. ■

5.8. Proof of Statement 4.2

5.9. Lemmas

We begin with three lemmas. The first is Slepian’s lemma.^(23, 5)

Lemma 5.6 (Slepian). Let Γ be a countable set, and let $X(t), Y(t)$ be two real zero-mean Gaussian processes indexed by $t \in \Gamma$. Suppose $EX^2(t) = EY^2(t)$ and $EX(s)X(t) \geq EY(s)Y(t)$ for all $s, t \in \Gamma$. Then, for all real λ , $P(\max_{t \in \Gamma} X(t) \geq \lambda) \leq P(\max_{t \in \Gamma} Y(t) \geq \lambda)$.

Consequently, if X and Y have continuous versions, Lemma 5.6 holds when the index set Γ is $[0, 1]$.

The next lemma gives us Markov property.

Lemma 5.7. Let α, f be as in Theorem 4.1, and let $0 < \beta < \alpha$. Suppose $f \geq 0$. Then there exists a Gaussian Markov process $Y(t)$ with independent increments and continuous paths such that $EY(t) = 0$ and $EY(s)Y(t) = q_f(0, s) = \int_0^s \int_0^s f(u) f(v) (du dv / |u - v|^{1-\alpha})$ whenever $s \leq t$.

Proof. For $f \geq 0$ the correlation function $q_f(0, s)$ is non-decreasing. It is also non-negative definite. Indeed, for any c_1, \dots, c_n , and $s_1 \leq \dots \leq s_n$,

$$\begin{aligned} E \left(\sum_{i=1}^n c_i Y(s_i) \right)^2 &= \sum_{i,j=1}^n c_i c_j q_f(0, s_i) \\ &= \sum_{i=1}^n \left(c_i^2 + 2 \sum_{i < j} c_i c_j \right) q_f(0, s_i) \\ &= \sum_{i=1}^n ((c_i + \dots + c_n)^2 - (c_{i+1} + \dots + c_n)^2) q_f(0, s_i) \\ &= (c_1 + \dots + c_n)^2 q_f(0, s_1) \\ &\quad + \sum_{i=2}^n (c_i + \dots + c_2)^2 (q_f(0, s_{i+1}) - q_f(0, s_i)) \geq 0, \end{aligned}$$

where the last inequality is strict unless f vanishes on some interval.

To show that a process is Markov, it is sufficient to show that its non-overlapping increments are independent. For a Gaussian process this amounts to checking that any two non-overlapping increments are uncorrelated: for $s_1 < t_1 < s_2 < t_2$,

$$\begin{aligned} & E(Y(t_1) - Y(s_1))(Y(t_2) - Y(s_2)) \\ &= EY(t_1) Y(t_2) - EY(t_1) Y(s_2) - EY(s_1) Y(t_2) + EY(s_1) Y(s_2) \\ &= q_f(0, t_1) - q_f(0, t_1) - q_f(0, s_1) + q_f(0, s_1) = 0 \end{aligned}$$

Since $Y(t)$ is a Gaussian Markov process with independent increments, it can be represented as a deterministic time change of a brownian motion:

$$Y(t) = B(q_f(0, t)).$$

Thus it has a continuous version. ■

Finally, we have a reflection principle:

Statement 5.8. Let $Y(t)$ be a zero-mean Gaussian Markov process with independent increments and continuous paths. Then for all $\lambda > 0$ and $T \geq 0$, $P(\max_{0 \leq t \leq T} Y(t) \geq \lambda) = 2P(Y(T) \geq \lambda)$.

The proof is exactly analogous to the oBm case, since $Y(t)$ is a deterministic time change of a Brownian motion.

Now we are ready to prove Theorem 4.2.

5.9.1. Proof of Part (1)

Proof. Choose an integer m such that $(\alpha - \beta)m > 1$. Then

$$\begin{aligned} & E \left(\int_0^t f(\tau) d\xi_\alpha(\tau) - \int_0^s f(\tau) d\xi_\alpha(\tau) \right)^{2m} \\ & \leq \text{const} \cdot \left(E \left(\int_0^t f(\tau) d\xi_\alpha(\tau) - \int_0^s f(\tau) d\xi_\alpha(\tau) \right)^2 \right)^m \\ & = \left(C\alpha(\alpha + 1) \int_s^t \int_s^t \frac{f(u)f(v)}{|u-v|^{1-\alpha}} du dv \right)^m \\ & \leq \text{const} \cdot |s-t|^{(\alpha-\beta)m}, \end{aligned}$$

where the first inequality holds because $\int_0^t f(\tau) d\zeta_\alpha(\tau)$ is a Gaussian random variable, and the second by the Hardy–Littlewood–Sobolev inequality:

$$\begin{aligned}
 q_f(s, t) &= C\alpha(\alpha + 1) \int_s^t \int_s^t \frac{f(u) f(v)}{|u - v|^{1-\alpha}} du dv \\
 &\leq C\alpha(\alpha + 1) |s - t|^{\alpha-\beta} \int_s^t \int_s^t \frac{f(u) f(v)}{|u - v|^{1-\beta}} du dv \\
 &\leq \text{const} \cdot |s - t|^{\alpha-\beta} \|f\|_{2/(1+\beta)}^2.
 \end{aligned}
 \tag{25}$$

By Kolmogorov’s criterion the process $\int_0^t f(\tau) d\zeta_\alpha(\tau)$ admits a continuous version. ■

5.9.2. Proof of Part (2)

Proof. Let $X(t) = \int_0^t f(\tau) d\zeta_\alpha(\tau)$. $X(t)$ is a Gaussian process with $EX(t) = 0$ and $EX(s) X(t) = \int_0^s \int_0^t f(u) f(v) (du dv / |u - v|^{1-\alpha})$. Define $Y(t)$ to be a Gaussian process with $EY(t) = 0$ and $EY(s) Y(t) = q_f(0, s)$ for $s \leq t$, as in Lemma 5.7. Clearly $EX(t)^2 = EY(t)^2$.

Suppose $f \geq 0$. Then $EX(s) X(t) \geq EY(s) Y(t)$ for $s, t \in [0, 1]$. Therefore the processes $X(t)$ and $Y(t)$ satisfy the assumptions of Slepian’s lemma (Lemma 5.6). By Lemma 5.7, $Y(t)$ is a Markov process with independent increments and continuous paths, and by Statement 5.8, Y obeys the reflection principle:

$$P\left(\max_{0 \leq t \leq 1} Y(t) \geq \lambda\right) = 2P(Y(1) \geq \lambda) = \int_{\lambda/\sqrt{q_f(0,1)}}^\infty \sqrt{\frac{2}{\pi}} e^{-x^2/2} dx.$$

Hence for $f \geq 0$,

$$P\left(\max_{0 \leq t \leq 1} X(t) \geq \lambda\right) \leq \int_{\lambda/\sqrt{q_f(0,1)}}^\infty \sqrt{\frac{2}{\pi}} e^{-x^2/2} dx.$$

Let $f \in L^{2/(1+\beta)}([0, \infty))$. Write $X_\pm(t) = \int_0^t f_\pm(\tau) d\zeta_\alpha(\tau)$. Define processes $Y_\pm(t)$ by replacing f by f_\pm in the definition of $Y(t)$. Since Slepian’s lemma applies also to $-X_-(t)$ and $Y_-(t)$, we have

$$P\left(\max_{0 \leq t \leq 1} \pm X_\pm(t) \geq \lambda\right) \leq \int_{\lambda/\sqrt{q_{f_\pm}(0,1)}}^\infty \sqrt{\frac{2}{\pi}} e^{-x^2/2} dx.$$

Therefore

$$\begin{aligned}
 P(\max_{0 \leq t \leq 1} X(t) \geq \lambda) &= P(\max_{0 \leq t \leq 1} X_+(t) + \max_{0 \leq t \leq 1} (-X_-(t)) \geq \lambda) \\
 &\leq P(\max_{0 \leq t \leq 1} X_+(t) \geq \lambda r) + P(\max_{0 \leq t \leq 1} (-X_-(t)) \geq \lambda(1-r)) \\
 &\leq \int_{\lambda r / \sqrt{q_{f_+}(0,1)}}^{\infty} + \int_{\lambda(1-r) / \sqrt{q_{f_-}(0,1)}}^{\infty} \sqrt{\frac{2}{\pi}} e^{-x^2/2} dx. \quad \blacksquare
 \end{aligned}$$

5.9.3. Proof of Part (3)

Proof. Inequality (25) shows that $\sqrt{q_f(s, t)} \leq \text{const} \cdot |s - t|^{(\alpha - \beta)/2}$, so

$$\int_1^{\infty} \sup_{|s-t| < m^{-x^2}} \sqrt{q_f(s, t)} dx < \infty.$$

This is the condition of applicability of Fernique's inequality,⁽²⁴⁾ of which the claim is a direct consequence. \blacksquare

ACKNOWLEDGMENTS

I would like to thank my advisor Yakov Sinai for suggesting the problem and for all his help during the course of the work, Almut Burchard for suggesting a way to simplify the argument of Section 5.3, the referee for his helpful suggestions in restructuring the paper and especially for an elegant way to simplify the argument of Section 5.4, Tadashi Tokieda for his help and comments on the text and Luc Rey-Bellet for his comments on the text.

REFERENCES

1. A. N. Kolmogorov, Wiener screw-line and other interesting curves in Hilbert space, *Doklady Akad. Nauk* **26**:115–118 (1940).
2. P. Lévy, *Processus Stochastiques et Movement Brownien* (Paris, 1948, 2nd ed., 1965).
3. B. Mandelbrot and J. W. Van Ness, Fractional Brownian motions, fractional noises and applications, *SIAM Rev.* **10**:422–437 (1968).
4. A. Goldman, Estimations analytiques concernant le mouvement brownien fractionnaire à plusieurs parametres, *C. R. Acad. Sci. Paris, Sér. I* **298**:91–93 (1984).
5. J.-P. Kahane, Some random series of functions, *Cambridge Studies in Adv. Math.* **5** (1985).
6. J.-P. Kahane, Sur les mouvements browniens fractionnaires: images, graphes, niveaux, *C. R. Acad. Sci. Paris, Sér. I* **300**:501–503 (1985).
7. A. Le Breton, Filtering and parameter estimation in a simple linear system driven by a fractional Brownian motion, *Statist. Probab. Lett.* **38**:263–274 (1998).
8. G. M. Molchan, Maximum of fractional Brownian motion: probabilities of small values (preprint).

9. Ya. G. Sinaĭ, On the distribution of the maximum of fractional Brownian motion, *Uspekhi Mat. Nauk* **52**:119–138 (1997).
10. M. Talagrand, Hausdorff measure of trajectories of multiparameter fractional Brownian motion, *Ann. Probab.* **23**:767–775 (1995).
11. M. Talagrand, Lower classes for fractional Brownian motion, *J. Theoret. Probab.* **9**:191–213 (1996).
12. M. Weber, Dimension de Hausdorff et points multiples du mouvement brownien fractionnaire dans \mathbb{R}^n , *C. R. Acad. Sci. Paris, Sér. I* **297**:357–360 (1983).
13. Y. Xiao, Hausdorff-type measures of the sample paths of fractional Brownian motion, *Stochastic Process. Appl.* **74**:251–272 (1998).
14. M. Yor, Remarques sur certaines constructions des mouvements browniens fractionnaires, *Séminaire de Probabilités, XXII*, Lecture Notes in Math., Vol. 1321 (Springer, Berlin/New York, 1988), pp. 217–224.
15. S. J. Lin, Stochastic analysis of fractional Brownian motions, *Stochastics Rep.* **55**:121–140 (1995).
16. W. Dai and C. C. Heyde, Itô's formula with respect to fractional Brownian motion and its application, *J. Appl. Math. Stochastic Anal.* **9**:439–448 (1996).
17. L. Decreusefond and Üstünel, Ali Süleyman, Application du calcul des variations stochastiques au mouvement brownien fractionnaire, *C. R. Acad. Sci. Paris, Sér. I* **321**:1605–1608 (1995).
18. L. C. Young, An inequality of the Hölder type connected with Stieltjes integration, *Acta Math.* **67**:251–282 (1936).
19. V. Kondurar, Sur l'intégrale de Stieltjes, *Mat. Sbornik* **2**:361–366 (1937).
20. A. M. Dyachkov, Conditions for the existence of Stieltjes integral of functions of bounded generalized variation, *Anal. Math.* **14**(4):295–313 (1988).
21. A. A. Ruzmaikina, *Stochastic Calculus with Fractional Brownian Motion*, Ph.D. thesis (Princeton University, 1999).
22. E. H. Lieb and M. Loss, *Analysis*, Graduate Studies in Mathematics, Vol. 14 (AMS, 1997).
23. D. Slepian, The one-sided barrier problem for Gaussian noise, *Bell System Tech. J.* **41**:463–501 (1962).
24. X. Fernique, Régularité des trajectoires des fonctions aléatoires gaussiennes, *École d'Été de Probabilités de Saint-Flour, avril 1974*, Lecture Notes in Math., Vol. 480 (Springer, Berlin, 1975), pp. 47–52.
25. H. P. McKean Jr., *Stochastic Integrals*, Probability and Mathematical Statistics, Vol. 5 (Academic Press, 1969).
26. D. Feyel and A. de La Pradelle, On fractional Brownian processes, *Potential Anal.* **10**:273–288 (1999).
27. M. Zähle, Integration with respect to fractal functions and stochastic calculus I, *Probab. Theory Related Fields* **111**:333–374 (1998).