

Linear Response Theory for Thermally Driven Quantum Open Systems

V. Jakšić,¹ Y. Ogata,^{2,3} and C.-A. Pillet²

Received September 27, 2005; accepted February 21, 2006

Published Online: June 8, 2006

This note is a continuation of our recent paper [V. Jakšić, Y. Ogata, and C.-A. Pillet, The Green-Kubo formula and Onsager reciprocity relations in quantum statistical mechanics. *Commun. Math. Phys.* in press.] where we have proven the Green-Kubo formula and the Onsager reciprocity relations for heat fluxes in thermally driven quantum open systems. In this note we extend the derivation of the Green-Kubo formula to heat and charge fluxes and discuss some other generalizations of the model and results of [V. Jakšić, Y. Ogata and C.-A. Pillet, The Green-Kubo formula and Onsager reciprocity relations in quantum statistical mechanics. *Commun. Math. Phys.* in press.].

KEY WORDS: nonequilibrium statistical mechanics, steady states, linear response, kubo formula, fluctuation-dissipation, onsager reciprocity

1. INTRODUCTION

This paper is the second in a series dedicated to linear response theory for non equilibrium steady states (NESS) of quantum open systems. The development of linear response theory is a part of a much wider research program initiated in Refs. 22–24, 29–31. This program deals with mathematical foundations of non-equilibrium thermodynamics in the framework of algebraic quantum statistical mechanics. Motivated by the developments in classical non-equilibrium statistical mechanics (see the review, Ref. 32) the program addresses the central issue of NESS in two independent steps.

¹ Department of Mathematics and Statistics, McGill University, 805 Sherbrooke Street West, Montreal, QC, H3A 2K6, Canada; e-mail: jaksic@math.mcgill.ca

² CPT-CNRS, UMR 6207, Université du Sud, Toulon-Var, B.P. 20132, F-83957 La Garde Cedex, France; e-mail: ogata@monet.phys.s.u-tokyo.ac.jp

³ Department of Mathematical Sciences, University of Tokyo, Komaba, Tokyo, 153-8914, Japan; e-mail: pillet@univ-tln.fr

(A) The existence and analytic properties of NESS are assumed as an *axiom*. On the basis of this axiom one develops the mathematical theory of non-equilibrium quantum statistical mechanics in an abstract setting. This step is primarily concerned with the mathematical structure of the theory and its relation to the fundamental physical aspects of non-equilibrium (see Refs. 13, 26).

(B) The second step concerns study of specific physically relevant models. Relaxation to a NESS and analytical properties of this NESS are detailed dynamical problems which can be answered only in the context of concrete models. Once these fundamental problems are solved, the thermodynamics and the transport theory of the model are derived from the general structural results established in (A).

So far, the main focus of the program has been the second law of thermodynamics (positivity of the entropy production). In this case the part (A) has been settled in Refs. 22, 25, 30, where the entropy production has been defined in the abstract framework of algebraic quantum statistical mechanics. In these works various structural properties of the entropy production have been established and in particular it was shown that the entropy production of any NESS is non-negative. The strict positivity of the entropy production is a problem which belongs to the category (B). At the moment there are two classes of non-trivial models whose NESS are well-understood and which have strictly positive entropy production. The first class of models describes an N -level quantum system coupled to finitely many independent free Fermi gas reservoirs.^(12,23,27) The second class describes finitely many free Fermi gas reservoirs coupled by local interactions.^(8,9,17) Some exactly solvable spin or fermion models with strictly positive entropy production have been studied in Refs. 4–7.

The natural next step in this program is the development of linear response theory and in particular the derivation of the Green-Kubo formulas (abbreviated GKF). A typical physical situation we consider concerns the steady states of a quantum device, a confined system \mathcal{S} with a finite number of degrees of freedom, coupled to M reservoirs $\mathcal{R}_1, \dots, \mathcal{R}_M$, see Figure 1 (a generalization of this setup is discussed in Sec. 5). More specifically, we are interested in situations where the system \mathcal{S} is driven out of equilibrium by thermodynamic forces, i.e., by discrepancies in the intensive thermodynamic parameters of the reservoirs around some common equilibrium values. Suppose that each reservoir \mathcal{R}_j is in a thermal equilibrium state characterized by some inverse temperature $\beta_j = \beta - X_j$ and chemical potential $\mu_j = \mu + Y_j/\beta$. If some of the forces X_j, Y_j do not vanish, then under normal conditions they induce energy and mass/charge currents across the system \mathcal{S} . Linear response theory is concerned with the calculation of these currents to first order in the forces. In⁽¹⁸⁾ we have derived the GKF for heat fluxes (the case $\mu = Y_j = 0$) in the axiomatic framework of algebraic quantum statistical mechanics. In this note we discuss a derivation which applies to both heat *and* charge fluxes and complete the step (A) of the program. Concerning (B), the




Fig. 1. An open system with M reservoirs.

examples to which our derivation directly applies include all models for which the strict positivity of the entropy production has been established. These applications are discussed in the forthcoming papers.^(19–21)

In classical mechanics there is a number of different ways to describe an open system out of thermal equilibrium. Some of these descriptions involve various kinds of thermostating devices which lead to non-Hamiltonian effective equations of motion (see Refs. 16, 28). Due to the intrinsic Hamiltonian nature of quantum dynamics, the situation is different for quantum open systems. Except in some special limiting cases (e.g., in the weak coupling limit, see Ref. 27 one is forced to consider the joint dynamics of the system S and its environment.

To describe the joint system $S + \mathcal{R}_1 + \dots + \mathcal{R}_M$ we suppose that it is initially prepared in a state where each reservoir \mathcal{R}_j is characterized by intensive thermodynamic parameters β_j and μ_j . Due to the interactions between the system S and the reservoirs this state is not stationary. We shall assume that, as $t \rightarrow +\infty$, the joint system relaxes to a steady state. Since confined quantum systems have discrete spectrum and almost periodic dynamics, a non-trivial steady state may exist only if the reservoirs are infinitely extended. Moreover, in order for this steady state to be a NESS (i.e., to avoid the joint system to relax to an equilibrium state), the reservoirs must be “ideal” in the following sense. A reservoir serves two purposes: on the one hand it is a source feeding energy/particles to the device S in a statistically controlled way. On the other hand it also works as a sink or dissipator, transporting to spatial infinity energy/particles coming out of S . In an

“ideal” reservoir the source and the sink are independent as much as quantum dynamics allows. The fact that incoming and outgoing fluxes do not interact ensures that the intensive thermodynamic parameters describing the initial state of an “ideal” reservoir still apply to the outgoing flux in a steady state. Thus, “ideal” reservoirs are able to maintain fluxes across the system \mathcal{S} over the infinite time interval needed to reach a steady state.

Linear response theory of NESS is a delicate interplay between three limits which must be taken in a definite order. First, one must perform the thermodynamic (or infinite volume) limit of the reservoirs. Then, a $t \rightarrow +\infty$ limit is necessary to reach a NESS. Final $X_j, Y_j \rightarrow 0$ limits are needed to extract the linear response proper.

For interacting quantum systems the first limit is already a difficult problem which can only be treated in a limited number of models (see e.g. Chapter 6 in Ref. 11). However, for the ideal reservoirs we are dealing with, this problem is well understood (see e.g. Sec. 5.2 in Ref. 11). An infinitely extended quantum dynamical system at non-vanishing density can be described in the universal conceptual framework of algebraic quantum statistical mechanics. It is therefore possible to decouple the thermodynamic limit from the two remaining ones. In this paper we derive the GKF under the assumption that these two limits exist and can be interchanged. The justification of this fact is a delicate dynamical problem which belongs to the category (B) and will be treated in the aforementioned companion papers.

This note is organized as follows. For notational purposes, in Sec. 2 we quickly review a few basic notions of algebraic quantum statistical mechanics. In Sec. 3 we introduce the model and review basic concepts of non-equilibrium statistical mechanics (the reader may complement this section with reviews Refs. 6, 24). Linear response theory is discussed in Sec. 4. Our main result is stated in Subsec. 4.2. Its proof follows closely the arguments in Ref. 18 and is outlined in Subsec. 4.3. Various generalizations of our model and results are discussed in Sec. 5.

2. BASIC NOTIONS

Let \mathcal{O} be a C^* -algebra with identity $\mathbb{1}$ and $\tau^t, t \in \mathbb{R}$, a strongly continuous group of $*$ -automorphisms of \mathcal{O} . The group τ and the pair (\mathcal{O}, τ) are often called C^* -dynamics and C^* -dynamical system. A state ω on \mathcal{O} is called τ -invariant if $\omega \circ \tau^t = \omega$ for all $t \in \mathbb{R}$. An anti-linear involutive $*$ -automorphism $\Theta : \mathcal{O} \rightarrow \mathcal{O}$ is called *time-reversal* of (\mathcal{O}, τ) if $\Theta \circ \tau^t = \tau^{-t} \circ \Theta$ for all $t \in \mathbb{R}$. A state ω on \mathcal{O} is called *time-reversal invariant* if $\omega(\Theta(A)) = \omega(A^*)$ for all $A \in \mathcal{O}$.

We call quantum dynamical system a triple $(\mathcal{O}, \tau, \omega)$ where ω is a given state on \mathcal{O} . The state ω describes the initial (or reference) thermodynamical state of the system and is not necessarily τ -invariant (for a discussion of this point we

refer the reader to Sec. 2 of Ref. 6). Under normal conditions, i.e., under natural ergodicity assumptions, all ω -normal states are thermodynamically equivalent reference states in the sense that they lead to the same NESS.

We denote by $\text{Ent}(\eta_1|\eta_2)$ the Araki relative entropy of two states η_1 and η_2 . We use the sign and ordering convention of^(11,14,15) (hence, $\text{Ent}(\eta_1|\eta_2) \in [-\infty, 0]$). The Araki relative entropy has played an important role in recent developments in non-equilibrium quantum statistical mechanics.

Let $\beta > 0$. A state ω is called a (τ, β) -KMS state if for all $A, B \in \mathcal{O}$ there exists a function $F_{A,B}(z)$, analytic in the strip $S_\beta = \{z \in \mathbb{C} \mid 0 < \text{Im } z < \beta\}$, bounded and continuous on its closure, and satisfying the KMS-boundary condition

$$F_{A,B}(t) = \omega(A\tau^t(B)), \quad F_{A,B}(t + i\beta) = \omega(\tau^t(B)A).$$

As usual, we write $\omega(A\tau^z(B)) = F_{A,B}(z)$ for $z \in \overline{S}_\beta$ even when $\tau^z(B)$ is not well-defined. A (τ, β) -KMS states describes a physical system in thermal equilibrium at inverse temperature β . For all practical purposes these states can be considered as thermodynamic limits of Gibbs canonical ensembles.

The general theory of chemical potential in quantum statistical mechanics is discussed in Sec. 5.4.3 of Ref. 11. In our study of linear response theory we will only consider the chemical potential associated to the usual $U(1)$ gauge invariance of quantum mechanics. We will call charge flux the current associated to the corresponding conserved charge. The extension of our results to more general gauge groups is straightforward. Since we only need a fraction of the mathematical structures commonly associated to the chemical potential we shall be brief. Let ϑ^φ be a C^* -dynamics on \mathcal{O} such that $\tau^t \circ \vartheta^\varphi = \vartheta^\varphi \circ \tau^t$ for all $t, \varphi \in \mathbb{R}$. ϑ is the gauge-group and its elements ϑ^φ are gauge transformations. Physical observables are invariant under gauge transformations and are therefore elements of

$$\mathcal{O}_\vartheta = \{A \in \mathcal{O} \mid \vartheta^\varphi(A) = A \text{ for all } \varphi \in \mathbb{R}\}. \tag{2.1}$$

Note that \mathcal{O}_ϑ is a τ -invariant C^* -subalgebra of \mathcal{O} and so $(\mathcal{O}_\vartheta, \tau)$ is a C^* -dynamical system. Let $\mu \in \mathbb{R}$ and

$$\alpha^t = \tau^t \circ \vartheta^{-\mu t}.$$

Clearly τ^t and α^t coincide on \mathcal{O}_ϑ . We say that a state ω on \mathcal{O} is a $(\tau, \vartheta, \beta, \mu)$ -KMS state if it is an (α, β) -KMS state. Although this last terminology is not common, it is convenient for our purposes. A $(\tau, \vartheta, \beta, \mu)$ -KMS state describes a physical system in equilibrium at inverse temperature β and chemical potential μ . Note that if ω is a $(\tau, \vartheta, \beta, \mu)$ -KMS state on \mathcal{O} , then its restriction to the gauge-invariant subalgebra \mathcal{O}_ϑ is a (τ, β) -KMS state on \mathcal{O}_ϑ which describes a thermodynamic limit of grand canonical ensembles associated to the parameters β, μ .

3. THE MODEL AND THE FRAMEWORK

3.1. The Model

Our starting point are two C^* -dynamical systems (\mathcal{O}_L, τ_L) and (\mathcal{O}_R, τ_R) with gauge-groups ϑ_L and ϑ_R . For convenience we shall call them the left, L, and the right, R, system. We denote the generators of $\tau_L, \tau_R, \vartheta_L$ and ϑ_R by $\delta_L, \delta_R, \xi_L$ and ξ_R . For many applications the left system can be thought of as composed of a first reservoir and a confined system, $L = \mathcal{S} + \mathcal{R}_1$, while the right system is just a second reservoir, $R = \mathcal{R}_2$. The generalizations of this setup are discussed in Sec. 5.

The C^* -algebra of the joint system $L + R$ is $\mathcal{O} = \mathcal{O}_L \otimes \mathcal{O}_R$ and its decoupled (non-interacting) dynamics is $\tau_0 = \tau_L \otimes \tau_R$. The generator of τ_0 is $\delta_0 = \delta_L \otimes I + I \otimes \delta_R$. In the sequel, whenever the meaning is clear within the context, we shall write δ_L for $\delta_L \otimes I$, δ_R for $I \otimes \delta_R$, etc.

The gauge-group of the joint system is $\vartheta = \vartheta_L \otimes \vartheta_R$ and its generator is $\xi = \xi_L + \xi_R$. We denote by \mathcal{O}_ϑ the corresponding gauge-invariant subalgebra of \mathcal{O} .

Let $V \in \mathcal{O}_\vartheta$ be a self-adjoint element describing the interaction of L and R. The interacting C^* -dynamics τ is generated by $\delta = \delta_0 + i[V, \cdot]$ and commutes with the gauge-group ϑ . The coupled (interacting) joint system $L + R$ is described by the C^* -dynamical system (\mathcal{O}, τ) .

3.2. The Reference States

We set $I_\epsilon(x) = (x - \epsilon, x + \epsilon)$ and write $I_\epsilon = I_\epsilon(0)$.

Let $\beta_{\text{eq}} > 0$ and $\mu_{\text{eq}} \in \mathbb{R}$ be given reference (equilibrium) values of the inverse temperature and the chemical potential. We make the following assumptions concerning the initial states of L and R.

(A1) ω_L is the unique $(\tau_L, \vartheta_L, \beta_{\text{eq}}, \mu_{\text{eq}})$ -KMS state on \mathcal{O}_L . The reference states of R are parametrized by $\beta \in I_{\epsilon_1}(\beta_{\text{eq}})$ and $\mu \in I_{\epsilon_2}(\mu_{\text{eq}})$ and $\omega_{R, \beta, \mu}$ is the unique $(\tau_R, \vartheta_R, \beta, \mu)$ -KMS state on \mathcal{O}_R . We shall denote $\omega_{R, \beta_{\text{eq}}, \mu_{\text{eq}}}$ by ω_R .

Throughout the paper we shall assume that (A1) holds. The reference states of our model are $\omega_L \otimes \omega_{R, \beta, \mu}$, $\beta \in I_{\epsilon_1}(\beta_{\text{eq}})$, $\mu \in I_{\epsilon_2}(\mu_{\text{eq}})$. For our purposes it is convenient to introduce the parameters (thermodynamical forces)

$$X = \beta_{\text{eq}} - \beta, \quad Y = \beta\mu - \beta_{\text{eq}}\mu_{\text{eq}},$$

and to parametrize the reference states by X and Y , i.e., we write

$$\omega_{X, Y, 0} = \omega_L \otimes \omega_{R, \beta, \mu}.$$

Since we are interested in linear response theory, without loss of generality we may restrict the values of X, Y to I_ϵ , where $\epsilon > 0$ is a small positive number. Note that $\omega_{0, 0, 0}$ is the unique $(\tau_0, \vartheta, \beta_{\text{eq}}, \mu_{\text{eq}})$ -KMS state on \mathcal{O} .

As we have already mentioned, under normal conditions all $\omega_{X,Y,0}$ -normal states are thermodynamically equivalent reference states of $L + R$. We now describe a particular $\omega_{X,Y,0}$ -normal reference state which will play an important role in our discussion of linear response theory.

Set

$$\alpha_L^t = \tau_L^t \circ \vartheta_L^{-\mu_{\text{eq}}t}, \quad \alpha_{R,\mu}^t = \tau_R^t \circ \vartheta_R^{-\mu t}.$$

Assumption (A1) implies that ω_L is the unique $(\alpha_L, \beta_{\text{eq}})$ -KMS state on \mathcal{O}_L and that $\omega_{R,\beta,\mu}$ is the unique $(\alpha_{R,\mu}, \beta)$ -KMS state on \mathcal{O}_R . Set

$$\alpha_{X,Y,0}^t = \alpha_L^t \otimes \alpha_{R,\mu}^{\beta t / \beta_{\text{eq}}}.$$

Then $\omega_{X,Y,0}$ is the unique $(\alpha_{X,Y,0}, \beta_{\text{eq}})$ -KMS state on \mathcal{O} . Let $\delta_{X,Y,0}$ be the generator of $\alpha_{X,Y,0}$ and

$$\delta_{X,Y} = \delta_{X,Y,0} + i[V, \cdot].$$

The subalgebra $\text{Dom}(\delta_L) \cap \text{Dom}(\xi_L) \cap \text{Dom}(\delta_R) \cap \text{Dom}(\xi_R)$ is a core for $\delta_{X,Y,0}$ and $\delta_{X,Y}$. On this subalgebra $\delta_{X,Y,0}$ acts as

$$\delta_{X,Y,0} = \delta_0 - \mu_{\text{eq}}\xi - \frac{X}{\beta_{\text{eq}}}\delta_R - \frac{Y}{\beta_{\text{eq}}}\xi_R. \tag{3.2}$$

Let $\alpha_{X,Y}$ be the C^* -dynamics generated by $\delta_{X,Y}$. Araki's perturbation theory yields that there exists a unique $(\alpha_{X,Y}, \beta_{\text{eq}})$ -KMS state $\omega_{X,Y}$ on \mathcal{O} .

The states $\omega_{X,Y,0}$ and $\omega_{X,Y}$ are mutually normal. The reference states $\omega_{X,Y}$ will play a central role in our study of linear response theory. Note that $\omega_{0,0}$ is the unique $(\tau, \vartheta, \beta_{\text{eq}}, \mu_{\text{eq}})$ -KMS state on \mathcal{O} . We denote this state by ω_{eq} . The next assumption concerns the $(\tau, \beta_{\text{eq}})$ -KMS state induced by ω_{eq} on the gauge invariant subalgebra \mathcal{O}_ϑ .

(A2) For all $A, B \in \mathcal{O}_\vartheta$,

$$\lim_{|t| \rightarrow \infty} \omega_{\text{eq}}(\tau^t(A)B) = \omega_{\text{eq}}(A)\omega_{\text{eq}}(B).$$

A well-known consequence of the KMS condition and Assumption (A2) is the relation

$$\lim_{t \rightarrow +\infty} \int_{-t}^t \omega_{\text{eq}}([\tau^s(A), B]) ds = 0, \tag{3.3}$$

which holds for all $A, B \in \mathcal{O}_\vartheta$ (see Theorem 5.4.12 in Ref. 11). This relation plays a key role in the derivation of the Onsager reciprocity relations.

3.3. Non-Equilibrium Steady States

We postulate relaxation to a NESS as follows:

(A3) For all $X, Y \in I_\epsilon$ there exists a state $\omega_{X,Y,+}$ on \mathcal{O}_ϑ such that for all $A \in \mathcal{O}_\vartheta$,

$$\lim_{t \rightarrow +\infty} \omega_{X,Y}(\tau^t(A)) = \omega_{X,Y,+}(A).$$

Assumptions (A2) and (A3) are strong ergodic hypotheses which are difficult to verify in concrete models. We remark that in typical physical situations one expects more, namely that

$$\lim_{t \rightarrow +\infty} \eta(\tau^t(A)) = \omega_{X,Y,+}(A),$$

for all $\omega_{X,Y,0}$ -normal states η and $A \in \mathcal{O}_\vartheta$. Indeed, such strong form of approach to NESS has been established in all examples we consider in Refs. 19–21. However, we will not need such an assumption in our axiomatic study of linear response theory.

3.4. Time-Reversal Invariance

Our next assumption concerns time-reversal.

(A4) There exists a time-reversal Θ of (\mathcal{O}, τ_0) such that $\Theta(V) = V$ and

$$\begin{aligned} \Theta \circ \tau_L^t &= \tau_L^{-t} \circ \Theta, & \Theta \circ \tau_R^t &= \tau_R^{-t} \circ \Theta, \\ \Theta \circ \vartheta_L^\varphi &= \vartheta_L^{-\varphi} \circ \Theta, & \Theta \circ \vartheta_R^\varphi &= \vartheta_R^{-\varphi} \circ \Theta, \end{aligned}$$

for all $t, \varphi \in \mathbb{R}$.

Clearly, Θ is a time-reversal of (\mathcal{O}, ϑ) and $(\mathcal{O}, \alpha_{X,Y,0})$. In particular it leaves \mathcal{O}_ϑ invariant. It is not difficult to show that Θ is also a time-reversal of (\mathcal{O}, τ) and $(\mathcal{O}, \alpha_{X,Y})$, and that the states $\omega_{X,Y,0}$ and $\omega_{X,Y}$ are time-reversal invariant. The proofs of these facts are the same as the proof of Lemma 3.1 in Ref. 18.

3.5. Fluxes

To define the flux observables we need:

(A5) $V \in \text{Dom}(\delta_R) \cap \text{Dom}(\xi_R)$.

If (A5) holds, we set

$$\Phi = \delta_R(V), \quad \mathcal{J} = \xi_R(V).$$

The observable Φ describes the heat flux out of the system R. The observable \mathcal{J} describes the charge flux out of R. Since $V \in \mathcal{O}_\vartheta$ and τ_R, ϑ_R commute with ϑ we have $\Phi, \mathcal{J} \in \mathcal{O}_\vartheta$. If the time-reversal assumption (A4) holds, then

$$\Theta(\Phi) = -\Phi, \quad \Theta(\mathcal{J}) = -\mathcal{J}.$$

3.6. Entropy Balance Equation

In the recent literature the entropy balance equation has been always discussed with respect to the product reference state $\omega_{X,Y,0}$.^(22,24,25,31) The finite time entropy balance equation w.r.t. the reference state $\omega_{X,Y}$ has the following form.

Theorem 3.1. *Assume that $V \in \text{Dom}(\delta_L) \cap \text{Dom}(\xi_L) \cap \text{Dom}(\delta_R) \cap \text{Dom}(\xi_R)$. Then*

$$\text{Ent}(\omega_{X,Y} \circ \tau^t | \omega_{X,Y}) = -X \int_0^t \omega_{X,Y}(\tau^s(\Phi)) ds - Y \int_0^t \omega_{X,Y}(\tau^s(\mathcal{J})) ds. \quad (3.4)$$

Proof. The assumptions of the theorem imply that $V \in \text{Dom}(\delta_{X,Y})$. Since $V \in \mathcal{O}_\vartheta$ implies $\xi(V) = 0$, we have

$$\beta_{\text{eq}} \delta_{X,Y}(V) = \beta_{\text{eq}} \delta(V) - X\Phi - Y\mathcal{J}. \quad (3.5)$$

The entropy balance equation of Refs. 22, 25 yields

$$\begin{aligned} \text{Ent}(\omega_{X,Y} \circ \tau^t | \omega_{X,Y,0}) &= \text{Ent}(\omega_{X,Y} | \omega_{X,Y,0}) + \beta_{\text{eq}} \int_0^t \omega_{X,Y}(\tau^s(\delta_{X,Y}(V))) ds \\ &= \text{Ent}(\omega_{X,Y} | \omega_{X,Y,0}) + \beta_{\text{eq}} \omega_{X,Y}(\tau^t(V)) - \beta_{\text{eq}} \omega_{X,Y}(V) \\ &\quad - X \int_0^t \omega_{X,Y}(\tau^s(\Phi)) ds - Y \int_0^t \omega_{X,Y}(\tau^s(\mathcal{J})) ds. \end{aligned} \quad (3.6)$$

The fundamental formula of Araki^(1,2) (see also Refs. 11, 14, 15) yields that

$$\begin{aligned} \text{Ent}(\omega_{X,Y} \circ \tau^t | \omega_{X,Y}) &= \text{Ent}(\omega_{X,Y} \circ \tau^t | \omega_{X,Y,0}) - \beta_{\text{eq}} \omega_{X,Y}(\tau^t(V)) + C, \\ \text{Ent}(\omega_{X,Y} | \omega_{X,Y,0}) &= \beta_{\text{eq}} \omega_{X,Y}(V) - C, \end{aligned} \quad (3.7)$$

where C is a constant expressible in terms of the modular structure (we do not need its explicit form here). The relations (3.6) and (3.7) yield the statement. \square

The entropy production of the NESS $\omega_{X,Y,+}$ is defined by

$$\text{Ep}(\omega_{X,Y,+}) = - \lim_{t \rightarrow +\infty} \frac{\text{Ent}(\omega_{X,Y} \circ \tau^t | \omega_{X,Y})}{t}.$$

Theorem 3.1 yields

$$\text{Ep}(\omega_{X,Y,+}) = X\omega_{X,Y,+}(\Phi) + Y\omega_{X,Y,+}(\mathcal{J}) \geq 0, \quad (3.8)$$

and this relation is the second law of thermodynamics for our model. Of course, if $(X, Y) \neq (0, 0)$, then under normal conditions one expects that $\text{Ep}(\omega_{X,Y,+}) > 0$, i.e., that the fluxes are non-vanishing. The strict positivity of entropy production

is a detailed dynamical question which can be answered only in the context of specific models.

3.7. Centered Observables

An observable $A \in \mathcal{O}$ is called *centered* if $\omega_{X,Y}(A) = 0$ for all $X, Y \in I_\epsilon$. We denote by \mathcal{C} the set of all centered observables. Obviously, \mathcal{C} is a norm-closed vector subspace of \mathcal{O} . Our derivation of the Green-Kubo formula applies only to centered observables.

If Assumption (A4) holds, then any self-adjoint observable A satisfying $\Theta(A) = -A$ is centered. Indeed, since $\omega_{X,Y}$ is time-reversal invariant,

$$\omega_{X,Y}(A) = \omega_{X,Y}(\Theta(A)) = -\omega_{X,Y}(A),$$

and so $\omega_{X,Y}(A) = 0$. In particular, if (A4) holds, then the flux observables Φ and \mathcal{J} are centered.

It is an important fact that the flux observables are centered irrespectively of the time-reversal assumption. This fact will play a central role in our discussion of the Green-Kubo formula for systems which are not time-reversal invariant.

Proposition 3.2. *Under Assumption (A5)*

$$\omega_{X,Y}(\Phi) = \omega_{X,Y}(\mathcal{J}) = 0,$$

holds for all $X, Y \in I_\epsilon$.

Proof. Assume first that

$$V \in \text{Dom}(\delta_L) \cap \text{Dom}(\xi_L) \cap \text{Dom}(\delta_R) \cap \text{Dom}(\xi_R). \tag{3.9}$$

Note that C^* -dynamics $\alpha_{X,Y}$ is well-defined for all $X, Y \in \mathbb{R}$. The following generalization of the entropy balance Eq. (3.4) holds: for all $X, Y \in I_\epsilon$ and $Z, U \in \mathbb{R}$,

$$\begin{aligned} \text{Ent}(\omega_{X,Y} \circ \alpha'_{Z,U} | \omega_{X,Y}) &= -(X - Z) \int_0^t \omega_{X,Y}(\alpha_{Z,U}^s(\Phi)) \, ds \\ &\quad - (Y - U) \int_0^t \omega_{X,Y}(\alpha_{Z,U}^s(\mathcal{J})) \, ds. \end{aligned} \tag{3.10}$$

The proof of this relation is essentially the same as the proof of (3.4). The only difference is that the relation (3.5) is now replaced with

$$\beta_{\text{eq}} \delta_{X,Y}(V) = \beta_{\text{eq}} \delta_{Z,U}(V) - (X - Z)\Phi - (Y - U)\mathcal{J}. \tag{3.11}$$

The entropy balance equation of Refs. 22, 25 yields

$$\text{Ent}(\omega_{X,Y} \circ \alpha_{Z,U}^t | \omega_{X,Y,0}) = \text{Ent}(\omega_{X,Y} | \omega_{X,Y,0}) + \beta_{\text{eq}} \int_0^t \omega_{X,Y}(\alpha_{Z,U}^s(\delta_{X,Y}(V))) \, ds,$$

and the rest of the argument follows line by line the proof of Theorem 3.1.

The Eq. (3.10) yields

$$\lim_{t \downarrow 0} \frac{\text{Ent}(\omega_{X,Y} \circ \alpha_{Z,U}^t | \omega_{X,Y})}{t} = -(X - Z)\omega_{X,Y}(\Phi) - (Y - U)\omega_{X,Y}(\mathcal{J}),$$

and so for all $X, Y \in I_\epsilon$ and $Z, U \in \mathbb{R}$,

$$(X - Z)\omega_{X,Y}(\Phi) + (Y - U)\omega_{X,Y}(\mathcal{J}) \geq 0.$$

This relation yields the statement.

To prove the general case, let $V \in \text{Dom}(\delta_R) \cap \text{Dom}(\xi_R)$ and

$$V_j = \frac{j}{\pi} \int_{\mathbb{R}^2} e^{-j(t^2+s^2)} \tau_L^t \circ \vartheta_L^s(V) \, dt \, ds, \quad j = 1, 2, \dots$$

The observables V_j satisfy (3.9). Let $\omega_{X,Y,j}$ and Φ_j, \mathcal{J}_j be the reference state and the flux observables associated to V_j . We have established that for all $X, Y \in I_\epsilon$,

$$\omega_{X,Y,j}(\Phi_j) = \omega_{X,Y,j}(\mathcal{J}_j) = 0. \tag{3.12}$$

By the properties of analytic approximations (see Ref. 11) $\|\omega_{X,Y,j} - \omega_{X,Y}\| \rightarrow 0, \|\Phi_j - \Phi\| \rightarrow 0, \|\mathcal{J}_j - \mathcal{J}\| \rightarrow 0$ as $j \rightarrow \infty$, and the statement follows from (3.12). \square

Note that we did not use the gauge invariance of V in the above proof.

3.8. Regular Observables

As mentioned in the introduction, our derivation of GKF relies on the assumption that the $t \rightarrow +\infty$ limit can be interchanged with differentiation w.r.t. X, Y . We note that if the states $\omega_L, \omega_{R,\beta,\mu}$ are ergodic for β, μ sufficiently close to $\beta_{\text{eq}}, \mu_{\text{eq}}$ then it is not difficult to show that the states $\omega_{X,Y}$ are mutually singular for distinct values of X, Y . Therefore the differentiability of the function $(X, Y) \mapsto \omega_{X,Y}(\tau^t(A))$ is an extremely delicate question, already for *finite* t . However, as we shall see in Subsec. 4.2, one can prove that this function is differentiable at $X = Y = 0$ under very mild regularity assumptions on A , provided A is centered (this is the content of our main technical result, Theorem 4.2).

The following definition encapsulates our assumption on the interchange of limits.

Definition 3.3. Assume that (A3) holds. Let $A \in \mathcal{O}_\vartheta$ be an observable such that the function

$$(X, Y) \mapsto \omega_{X,Y}(\tau^t(A)),$$

is differentiable at $(0, 0)$ for all t . We call such an observable regular if the function

$$(X, Y) \mapsto \omega_{X,Y,+}(A),$$

is also differentiable at $(0, 0)$ and

$$\begin{aligned} \lim_{t \rightarrow +\infty} \partial_X \omega_{X,Y}(\tau^t(A)) \Big|_{X=Y=0} &= \partial_X \omega_{X,Y,+}(A) \Big|_{X=Y=0}, \\ \lim_{t \rightarrow +\infty} \partial_Y \omega_{X,Y}(\tau^t(A)) \Big|_{X=Y=0} &= \partial_Y \omega_{X,Y,+}(A) \Big|_{X=Y=0}. \end{aligned} \quad (3.13)$$

4. LINEAR RESPONSE THEORY

4.1. Overview

Suppose that Assumptions (A3) and (A5) hold and that the functions

$$(X, Y) \mapsto \omega_{X,Y,+}(\Phi), \quad (X, Y) \mapsto \omega_{X,Y,+}(\mathcal{J}),$$

are differentiable at $(0, 0)$. The kinetic transport coefficients are defined by

$$\begin{aligned} L_{\text{hh}} &= \partial_X \omega_{X,Y,+}(\Phi) \Big|_{X=Y=0}, \\ L_{\text{hc}} &= \partial_Y \omega_{X,Y,+}(\Phi) \Big|_{X=Y=0}, \\ L_{\text{ch}} &= \partial_X \omega_{X,Y,+}(\mathcal{J}) \Big|_{X=Y=0}, \\ L_{\text{cc}} &= \partial_Y \omega_{X,Y,+}(\mathcal{J}) \Big|_{X=Y=0}, \end{aligned} \quad (4.14)$$

where indices h/c stand for heat/charge. Linear response theory is concerned with these coefficients. An elementary consequence of the second law (Relation (3.8)) is that the matrix

$$L = \begin{bmatrix} L_{\text{hh}} & L_{\text{hc}} \\ L_{\text{ch}} & L_{\text{cc}} \end{bmatrix},$$

is positive definite on the real vector space \mathbb{R}^2 (this of course does not imply that $L_{\text{hc}} = L_{\text{ch}}!$).

The Green-Kubo formulas are at the center of linear response theory. For $A, B \in \mathcal{O}_\vartheta$ we set

$$\mathcal{L}(A, B) = \lim_{t \rightarrow +\infty} \frac{1}{2} \int_{-t}^t \omega_{\text{eq}}(A \tau^s(B)) \, ds.$$

The GKF assert that if the system is time-reversal invariant, then

$$\begin{aligned}
 L_{hh} &= \mathcal{L}(\Phi, \Phi), \\
 L_{hc} &= \mathcal{L}(\Phi, \mathcal{J}), \\
 L_{ch} &= \mathcal{L}(\mathcal{J}, \Phi), \\
 L_{cc} &= \mathcal{L}(\mathcal{J}, \mathcal{J}).
 \end{aligned}
 \tag{4.15}$$

These formulas are mathematical expressions of the fluctuation-dissipation mechanism in statistical mechanics—they link linear response to a thermodynamical force to the equilibrium correlations w.r.t. the corresponding flux observable.

The coefficients L_{hc} and L_{ch} are of particular physical importance. In words, the chemical potential difference may cause a heat flow out of R even if L and R are at the same temperature. For Y small, this flow is equal to $YL_{hc} + o(Y)$. Similarly, the temperature difference may cause a charge flow out of R even if L and R have equal chemical potentials. For X small this flow is equal to $XL_{ch} + o(X)$. An immediate consequence of the second and third relation in (4.15) and the formula (3.3) are the Onsager reciprocity relations

$$L_{hc} = L_{ch}. \tag{4.16}$$

For $A, B \in \mathcal{O}_\vartheta$ and $t \in \mathbb{R}$ we set

$$\mathfrak{L}(A, B, t) = \frac{1}{\beta_{\text{eq}}} \int_0^t ds \int_0^{\beta_{\text{eq}}} du \omega_{\text{eq}}(\tau^s(A)\tau^{iu}(B)),$$

and

$$\mathfrak{L}(A, B) = \lim_{t \rightarrow +\infty} \mathfrak{L}(A, B, t),$$

whenever the limit exists. We remark that by the KMS condition the function

$$(s, z) \mapsto \omega_{\text{eq}}(\tau^s(A)\tau^z(B)),$$

is bounded and continuous on the set $\mathbb{R} \times \overline{S}_{\beta_{\text{eq}}}$. The central step in our derivation of (4.15) are the following formulas

$$\begin{aligned}
 L_{hh} &= \mathfrak{L}(\Phi, \Phi), \\
 L_{hc} &= \mathfrak{L}(\Phi, \mathcal{J}), \\
 L_{ch} &= \mathfrak{L}(\mathcal{J}, \Phi), \\
 L_{cc} &= \mathfrak{L}(\mathcal{J}, \mathcal{J}).
 \end{aligned}
 \tag{4.17}$$

It is an important point that these formulas hold *without the time-reversal assumption*—they are the Green-Kubo formulas for systems which are not time-reversal invariant. The Green-Kubo formulas (4.15) are an immediate consequence of (4.17) and the following result established in Ref. 18.

Proposition 4.1. *Suppose that Assumptions (A1), (A2), and (A4) hold and let $A, B \in \mathcal{O}_\vartheta$ be two self-adjoint observables which are both even or odd under Θ . Then*

$$\mathcal{L}(A, B) = \mathfrak{L}(A, B).$$

Proof. The argument follows line by line the proof of Theorem 2.3 in Ref. 18. For reader convenience we outline the main steps of the argument.

We need to prove that

$$\lim_{t \rightarrow +\infty} \frac{1}{\beta_{\text{eq}}} \int_0^{\beta_{\text{eq}}} \left[\int_0^t \omega_{\text{eq}}(\tau^s(A)\tau^{iu}(B)) ds \right] du = \lim_{t \rightarrow +\infty} \int_{-t}^t \omega_{\text{eq}}(A\tau^s(B)) ds.$$

The time-reversal invariance and the KMS-condition yield that for $s \in \mathbb{R}$ and $u \in [0, \beta]$,

$$\omega_{\text{eq}}(\tau^s(A)\tau^{iu}(B)) = \omega_{\text{eq}}(\tau^{-s}(A)\tau^{i\beta_{\text{eq}}-iu}(B)),$$

and so

$$\frac{1}{\beta_{\text{eq}}} \int_0^{\beta_{\text{eq}}} \left[\int_0^t \omega_{\text{eq}}(\tau^s(A)\tau^{iu}(B)) ds \right] du = \frac{1}{2\beta_{\text{eq}}} \int_0^{\beta_{\text{eq}}} \left[\int_{-t}^t \omega_{\text{eq}}(A\tau^{s+iu}(B)) ds \right] du.$$

Since the integral of the function $z \mapsto \omega_{\text{eq}}(A\tau^z(B))$ over the boundary of the rectangle with vertices $-t, t, t + iu, -t + iu$ is zero, we have

$$\begin{aligned} \frac{1}{\beta_{\text{eq}}} \int_0^{\beta_{\text{eq}}} \left[\int_0^t \omega_{\text{eq}}(\tau^s(A)\tau^{iu}(B)) ds \right] du &= \frac{1}{2} \int_{-t}^t \omega_{\text{eq}}(A\tau^s(B)) ds \\ &+ \frac{1}{2\beta_{\text{eq}}} \int_0^{\beta_{\text{eq}}} R(t, u) du, \end{aligned} \tag{4.18}$$

where

$$R(t, u) = i \int_0^u [\omega_{\text{eq}}(A\tau^{t+iy}(B)) - \omega_{\text{eq}}(A\tau^{-t+iy}(B))] dy.$$

Assumption (A2) implies that

$$\lim_{t \rightarrow +\infty} \omega_{\text{eq}}(A\tau^{\pm t+iy}(B)) = \omega_{\text{eq}}(A)\omega_{\text{eq}}(B),$$

and the dominated convergence theorem yields

$$\lim_{t \rightarrow +\infty} \sup_{0 \leq u \leq \beta} |R(t, u)| = 0.$$

This fact and the formula (4.18) yield the statement. □

In the next subsection we state our main results concerning the Green-Kubo formulas.

4.2. The Green-Kubo Formulas

As already mentioned, our main technical result concerns the differentiability at $(0, 0)$ and for finite t of the function $(X, Y) \mapsto \omega_{X,Y}(\tau^t(A))$. The resulting *finite time linear response formula* is the content of the next Theorem. We set

$$\begin{aligned}\mathcal{O}_{\vartheta,\mathbb{R}} &= \mathcal{O}_{\vartheta} \cap \text{Dom}(\delta_{\mathbb{R}}) \cap \text{Dom}(\xi_{\mathbb{R}}), \\ \mathcal{O}_{\vartheta,\mathbb{R},\mathfrak{C}} &= \mathcal{O}_{\vartheta,\mathbb{R}} \cap \mathfrak{C}.\end{aligned}$$

Theorem 4.2. *Suppose that Assumptions (A1) and (A5) hold and let $A \in \mathcal{O}_{\vartheta,\mathbb{R},\mathfrak{C}}$. Then for all $t \in \mathbb{R}$ the function*

$$(X, Y) \mapsto \omega_{X,Y}(\tau^t(A)),$$

is differentiable at $(0, 0)$ and

$$\begin{aligned}\partial_X \omega_{X,Y}(\tau^t(A)) \Big|_{X=Y=0} &= \mathfrak{L}(A, \Phi, t), \\ \partial_Y \omega_{X,Y}(\tau^t(A)) \Big|_{X=Y=0} &= \mathfrak{L}(A, \mathcal{J}, t).\end{aligned}$$

We will prove Theorem 4.2 in Subsec. 4.3. The next two theorems are consequence of Theorem 4.2, definition of the regular observable, and Proposition 4.1.

Theorem 4.3. *Suppose that Assumptions (A1), (A3) and (A5) hold.*

(1) *Let $A \in \mathcal{O}_{\vartheta,\mathbb{R},\mathfrak{C}}$ be a regular observable. Then*

$$\begin{aligned}\partial_X \omega_{X,Y,+}(A) \Big|_{X=Y=0} &= \mathfrak{L}(A, \Phi), \\ \partial_Y \omega_{X,Y,+}(A) \Big|_{X=Y=0} &= \mathfrak{L}(A, \mathcal{J}).\end{aligned}$$

(2) *If in addition (A2) and (A4) hold and $A \in \mathcal{O}_{\vartheta,\mathbb{R}}$ is a regular self-adjoint observable such that $\Theta(A) = -A$, then*

$$\begin{aligned}\partial_X \omega_{X,Y,+}(A) \Big|_{X=Y=0} &= \mathfrak{L}(A, \Phi), \\ \partial_Y \omega_{X,Y,+}(A) \Big|_{X=Y=0} &= \mathfrak{L}(A, \mathcal{J}).\end{aligned}$$

Theorem 4.4. *Suppose that Assumptions (A1), (A3) and (A5) hold and that Φ, \mathcal{J} are regular observables in $\text{Dom}(\delta_{\mathbb{R}}) \cap \text{Dom}(\xi_{\mathbb{R}})$. Then the formulas (4.17) hold. If in addition (A2) and (A4) hold, then the formulas (4.15) and (4.16) hold.*

Theorem 4.2 was proven in Ref. 18 in the case $\mu_{\text{eq}} = 0, Y = 0$. The technical extensions of the proofs in Ref. 18 needed to accommodate charge fluxes are relatively minor and are discussed in the next section.

4.3. Proof of Theorem 4.2

We will freely use the notation introduced in Subsec. 3.1.

Lemma 4.5.

(a) *The group $\alpha_{0,0}$ preserves $\text{Dom}(\delta_{\mathbb{R}}) \cap \text{Dom}(\xi_{\mathbb{R}})$ and for $A \in \text{Dom}(\delta_{\mathbb{R}}) \cap \text{Dom}(\xi_{\mathbb{R}})$ the functions*

$$\mathbb{R} \ni t \mapsto \delta_{\mathbb{R}}(\alpha_{0,0}^t(A)), \quad \mathbb{R} \ni t \mapsto \xi_{\mathbb{R}}(\alpha_{0,0}^t(A)),$$

are norm continuous.

(b) *For all $t \in \mathbb{R}$ and $A \in \text{Dom}(\delta_{\mathbb{R}}) \cap \text{Dom}(\xi_{\mathbb{R}})$,*

$$\begin{aligned} \alpha_{X,Y}^t(A) - \alpha_{0,0}^t(A) &= -\frac{X}{\beta_{\text{eq}}} \int_0^t \alpha_{X,Y}^{t-s}(\delta_{\mathbb{R}}(\alpha_{0,0}^s(A))) ds \\ &\quad - \frac{Y}{\beta_{\text{eq}}} \int_0^t \alpha_{X,Y}^{t-s}(\xi_{\mathbb{R}}(\alpha_{0,0}^s(A))) ds. \end{aligned}$$

(c) *For all $t \in \mathbb{R}$ and $A \in \mathcal{O}$,*

$$\lim_{(X,Y) \rightarrow (0,0)} \|\alpha_{X,Y}^t(A) - \alpha_{0,0}^t(A)\| = 0.$$

(d) *For all $A \in \mathcal{O}$,*

$$\lim_{(X,Y) \rightarrow (0,0)} \omega_{X,Y}(A) = \omega_{\text{eq}}(A).$$

Proof. To simplify notation let us set $\alpha_0 = \alpha_{0,0,0}$ and $\alpha = \alpha_{0,0}$. We shall use the identity

$$\alpha^t(A) = \Gamma_t \alpha'_0(A) \Gamma_t^*,$$

where $\Gamma_t \in \mathcal{O}$ is a family of unitary elements defined by

$$\Gamma_t = \mathbb{1} + \sum_{n \geq 1} (it)^n \int_{0 \leq s_n \leq \dots \leq s_1 \leq 1} \alpha_0^{ts_n}(V) \cdots \alpha_0^{ts_1}(V) ds_1 \cdots ds_n,$$

see Proposition 5.4.1 in Ref. 11. Since $V \in \text{Dom}(\delta_{\mathbb{R}}) \cap \text{Dom}(\xi_{\mathbb{R}})$, one easily shows that $\Gamma_t \in \text{Dom}(\delta_{\mathbb{R}}) \cap \text{Dom}(\xi_{\mathbb{R}})$ and that

$$\delta_{\mathbb{R}}(\Gamma_t) = \sum_{n \geq 1} (it)^n \int_{0 \leq s_n \leq \dots \leq s_1 \leq 1} \sum_j \alpha_0^{ts_n}(V) \cdots \alpha_0^{ts_j}(\delta_{\mathbb{R}}(V)) \cdots \alpha_0^{ts_1}(V) ds_1 \cdots ds_n,$$

$$\xi_R(\Gamma_t) = \sum_{n \geq 1} (it)^n \int_{0 \leq s_n \leq \dots \leq s_1 \leq 1} \sum_j \alpha_0^{ts_n}(V) \cdots \alpha_0^{ts_j}(\xi_R(V)) \cdots \alpha_0^{ts_1}(V) ds_1 \cdots ds_n.$$

These two formulas yield that the functions

$$t \mapsto \delta_R(\Gamma_t), \quad t \mapsto \xi_R(\Gamma_t),$$

are norm continuous. Finally, the identities

$$\delta_R(\alpha^t(A)) = \delta_R(\Gamma_t)\alpha_0^t(A)\Gamma_t^* + \Gamma_t\alpha_0^t(\delta_R(A))\Gamma_t^* + \Gamma_t\alpha_0^t(A)\delta_R(\Gamma_t^*),$$

$$\xi_R(\alpha^t(A)) = \xi_R(\Gamma_t)\alpha_0^t(A)\Gamma_t^* + \Gamma_t\alpha_0^t(\xi_R(A))\Gamma_t^* + \Gamma_t\alpha_0^t(A)\xi_R(\Gamma_t^*),$$

yield Part (a).

If $A \in \text{Dom}(\delta_L) \cap \text{Dom}(\delta_R) \cap \text{Dom}(\xi_L) \cap \text{Dom}(\xi_R)$, then

$$\frac{d}{dt} \alpha_{X,Y}^{-t} \circ \alpha^t(A) = \frac{X}{\beta_{\text{eq}}} \alpha_{X,Y}^{-t}(\delta_R(\alpha^t(A))) + \frac{Y}{\beta_{\text{eq}}} \alpha_{X,Y}^{-t}(\xi_R(\alpha^t(A))),$$

and (b) follows. The case $A \in \text{Dom}(\delta_R) \cap \text{Dom}(\xi_R)$ is handled by approximating A with the sequence

$$A_j = \frac{j}{\pi} \int_{\mathbb{R}^2} e^{-j(t^2+s^2)} \tau_L^t \circ \vartheta_L^s(A) dt ds,$$

see the proof of Lemma 3.3 in Ref. 18.

Since $\text{Dom}(\delta_R) \cap \text{Dom}(\xi_R)$ is dense in \mathcal{O} , (b) implies (c). The proof of (d) is the same as the proof of Lemma 3.4 in Ref. 18. \square

Lemma 4.6. *Let $A \in \mathcal{O}_{\vartheta,R,C}$. Then for all $t \in \mathbb{R}$ the function*

$$(X, Y) \mapsto \omega_{X,Y}(\tau^t(A)),$$

is differentiable at $(0, 0)$, and

$$\partial_X \omega_{X,Y}(\tau^t(A)) \Big|_{X=Y=0} = \frac{1}{\beta_{\text{eq}}} \int_0^t \omega_{\text{eq}}(\delta_R(\tau^s(A))) ds,$$

$$\partial_Y \omega_{X,Y}(\tau^t(A)) \Big|_{X=Y=0} = \frac{1}{\beta_{\text{eq}}} \int_0^t \omega_{\text{eq}}(\xi_R(\tau^s(A))) ds.$$

Proof. Since A is a centered observable and $\omega_{X,Y}$ is $\alpha_{X,Y}$ -invariant, we have that $\omega_{X,Y}(\alpha_{X,Y}^t(A)) = 0$ for all t . Since $\alpha_{0,0} = \tau$ on \mathcal{O}_{ϑ} , we have that $\omega_{X,Y}(\alpha_{0,0}^t(A)) = \omega_{X,Y}(\tau^t(A))$ and $\omega_{0,0}(\tau^t(A)) = \omega_{\text{eq}}(\tau^t(A)) = 0$ for all t . These observations and Part (b) of Lemma 4.5 imply

$$\begin{aligned} \omega_{X,Y}(\tau^t(A)) - \omega_{0,0}(\tau^t(A)) &= \frac{X}{\beta_{\text{eq}}} \int_0^t \omega_{X,Y}(\delta_R(\tau^s(A))) ds \\ &\quad + \frac{Y}{\beta_{\text{eq}}} \int_0^t \omega_{X,Y}(\xi_R(\tau^s(A))) ds. \end{aligned}$$

This relation, Lemma 4.5, and dominated convergence yield the statement. \square

Lemma 4.7. *Assume that $A \in \mathcal{O}_{\vartheta, \mathbb{R}}$. Then*

$$\omega_{\text{eq}}(\delta_{\mathbb{R}}(A)) = \int_0^{\beta_{\text{eq}}} \omega_{\text{eq}}(A\tau^{\text{is}}(\Phi)) \, ds,$$

$$\omega_{\text{eq}}(\xi_{\mathbb{R}}(A)) = \int_0^{\beta_{\text{eq}}} \omega_{\text{eq}}(A\tau^{\text{is}}(\mathcal{J})) \, ds.$$

Proof. This lemma is the central and technically most demanding step of the argument. Fortunately, its proof is identical to the proof of Lemma 3.6 in Ref. 18. This follows from the fact that $A, V, \Phi, \mathcal{J} \in \mathcal{O}_{\vartheta}$ and that $\overline{\omega}_{\text{eq}}\mathcal{O}_{\vartheta}$ is a $(\tau, \beta_{\text{eq}})$ -KMS state. \square

Theorem 4.2 is an immediate consequence of Lemmas 4.6 and 4.7.

5. SOME GENERALIZATIONS

Although we have restricted ourselves in this note to two coupled quantum dynamical systems, the model, the framework and all our results have a straightforward extension to the case of M systems. Let β_{eq} and μ_{eq} be the reference (equilibrium) values of the inverse temperature and chemical potential. For $j = 1, \dots, M$ let $(\mathcal{O}_j, \tau_j, \omega_{j, \beta_j \mu_j})$ be quantum dynamical systems with gauge groups ϑ_j where ω_j is a $(\tau_j, \vartheta_j, \beta_j, \mu_j)$ -KMS state. We denote by δ_j and ξ_j the generators of τ_j and ϑ_j . Assumption (A1) is replaced with

(G1) The reference states of the j -th system are parametrized by $\beta_j \in I_{\epsilon}(\beta_{\text{eq}})$ and $\mu_j \in I_{\epsilon}(\mu_{\text{eq}})$ and $\omega_{j, \beta_j \mu_j}$ is the unique $(\tau_j, \vartheta_j, \beta_j, \mu_j)$ -KMS state on \mathcal{O}_j .

Let $\mathcal{O} = \mathcal{O}_1 \otimes \dots \otimes \mathcal{O}_M$, $\tau_0 = \tau_1 \otimes \dots \otimes \tau_M$, $\vartheta = \vartheta_1 \otimes \dots \otimes \vartheta_M$. The algebra \mathcal{O}_{ϑ} is again defined by (2.1). The pair (\mathcal{O}, τ_0) describes the uncoupled joint system. Let $V \in \mathcal{O}_{\vartheta}$ be a self-adjoint perturbation and τ the perturbed C^* -dynamics. The coupled joint system is described by (\mathcal{O}, τ) . The thermodynamical forces are

$$X_j = \beta_{\text{eq}} - \beta_j, \quad Y_j = \beta_j \mu_j - \beta_{\text{eq}} \mu_{\text{eq}}.$$

We set $X = (X_1, \dots, X_M)$, $Y = (Y_1, \dots, Y_M)$. The reference state is $\omega_{X, Y, 0} = \omega_{1, \beta_1 \mu_1} \otimes \dots \otimes \omega_{M, \beta_M \mu_M}$. $\omega_{X, Y, 0}$ is the unique β_{eq} -KMS state for the C^* -dynamics

$$\alpha_{X, Y, 0}^t = [\tau_1^{\beta_1 t / \beta_{\text{eq}}} \circ \vartheta_1^{-\mu_1 \beta_1 t / \beta_{\text{eq}}}] \otimes \dots \otimes [\tau_M^{\beta_M t / \beta_{\text{eq}}} \circ \vartheta_M^{-\mu_M \beta_M t / \beta_{\text{eq}}}] .$$

Let $\delta_{X, Y, 0}$ be the generator of $\alpha_{X, Y, 0}$ and $\delta_{X, Y} = \delta_{X, Y, 0} + i[V, \cdot]$. Let $\alpha_{X, Y}$ be the C^* -dynamics generated by $\delta_{X, Y}$ and let $\omega_{X, Y}$ be the $(\alpha_{X, Y}, \beta_{\text{eq}})$ -KMS state obtained from $\omega_{X, Y, 0}$ by Araki's perturbation theory. This completes the setup

of the model. Note that the state $\omega_{\text{eq}} \equiv \omega_{0,0}$ is the unique $(\tau, \vartheta, \beta_{\text{eq}}, \mu_{\text{eq}})$ -KMS state on \mathcal{O} . Assumption (G2) has the same formulation as Assumption (A2) and Assumptions (A3), (A4) and (A5) are replaced with:

(G3) For all $X, Y \in I_\epsilon^M$ there exists a state $\omega_{X,Y,+}$ on \mathcal{O}_ϑ such that for all $A \in \mathcal{O}_\vartheta$,

$$\lim_{t \rightarrow +\infty} \omega_{X,Y}(\tau^t(A)) = \omega_{X,Y,+}(A).$$

(G4) There exists a time-reversal Θ of (\mathcal{O}, τ_0) such that $\Theta(V) = V$ and

$$\Theta \circ \tau_j^t = \tau_j^{-t} \circ \Theta, \quad \Theta \circ \vartheta_j^t = \vartheta_j^{-t} \circ \Theta,$$

for all j .

(G5) $V \in \text{Dom}(\delta_j) \cap \text{Dom}(\xi_j)$ for all j .

The observables associated to the heat and charge flux out of the j -th system are

$$\Phi_j = \delta_j(V), \quad \mathcal{J}_j = \xi_j(V).$$

It immediately follows that

$$\sum_{j=1}^M \omega_{X,Y,+}(\Phi_j) = 0 \quad \text{and} \quad \sum_{j=1}^M \omega_{X,Y,+}(\mathcal{J}_j) = 0,$$

which are respectively the first law of thermodynamics (conservation of energy) and charge conservation. The entropy balance equation reads

$$\begin{aligned} \text{Ent}(\omega_{X,Y} \circ \tau^t | \omega_{X,Y}) &= - \sum_{j=1}^M X_j \int_0^t \omega_{X,Y}(\tau^s(\Phi_j)) ds \\ &\quad - \sum_{j=1}^M Y_j \int_0^t \omega_{X,Y}(\tau^s(\mathcal{J}_j)) ds, \end{aligned}$$

and in particular the second law holds:

$$\text{Ep}(\omega_{X,Y,+}) = \sum_{j=1}^M X_j \omega_{X,Y,+}(\Phi_j) + \sum_{j=1}^M Y_j \omega_{X,Y,+}(\mathcal{J}_j) \geq 0. \tag{5.19}$$

The definition of the centered observable is the same as in Subsec. 3.7. We set

$$\begin{aligned} \hat{\mathcal{O}}_\vartheta &= (\cap_{j=1}^M \text{Dom}(\delta_j)) \cap (\cap_{j=1}^M \text{Dom}(\xi_j)) \cap \mathcal{O}_\vartheta, \\ \hat{\mathcal{O}}_{\vartheta,c} &= \hat{\mathcal{O}}_\vartheta \cap \mathcal{C}. \end{aligned}$$

If $V \in \hat{\mathcal{O}}_{\vartheta}$, then $\Phi_j, \mathcal{J}_j \in \hat{\mathcal{O}}_{\vartheta,c}$ for all j (after obvious notational changes, Proposition 3.2 applies directly to the model consider in this section).

Theorem 4.2 is replaced with:

Theorem 5.1. *Suppose that Assumptions (G1) and (G5) hold and let $A \in \hat{\mathcal{O}}_{\vartheta,c}$. Then for all $t \in \mathbb{R}$ the function*

$$(X, Y) \mapsto \omega_{X,Y}(\tau^t(A)),$$

is differentiable at $(0, 0)$ and

$$\partial_{X_j} \omega_{X,Y}(\tau^t(A)) \Big|_{X=Y=0} = \mathfrak{L}(A, \Phi_j, t),$$

$$\partial_{Y_j} \omega_{X,Y}(\tau^t(A)) \Big|_{X=Y=0} = \mathfrak{L}(A, \mathcal{J}_j, t).$$

The definition of the regular observable is the same as before, and we have:

Theorem 5.2. *Suppose that Assumptions (G1), (G3) and (G5) hold.*

(1) *Let $A \in \hat{\mathcal{O}}_{\vartheta,c}$ be a regular observable. Then*

$$\partial_{X_j} \omega_{X,Y,+}(A) \Big|_{X=Y=0} = \mathfrak{L}(A, \Phi_j),$$

$$\partial_{Y_j} \omega_{X,Y,+}(A) \Big|_{X=Y=0} = \mathfrak{L}(A, \mathcal{J}_j).$$

(2) *If in addition (G2) and (G4) hold and $A \in \hat{\mathcal{O}}_{\vartheta}$ is a regular self-adjoint observable such that $\Theta(A) = -A$, then*

$$\partial_{X_j} \omega_{X,Y,+}(A) \Big|_{X=Y=0} = \mathcal{L}(A, \Phi_j),$$

$$\partial_{Y_j} \omega_{X,Y,+}(A) \Big|_{X=Y=0} = \mathcal{L}(A, \mathcal{J}_j).$$

Theorem 5.3. *Suppose that (G1), (G3) and (G5) hold and that Φ_j, \mathcal{J}_j are regular observables in $\text{Dom}(\delta_j) \cap \text{Dom}(\xi_j)$. Then:*

(1) *The kinetic transport coefficients*

$$L_{\text{hh}}^{kj} = \partial_{X_j} \omega_{X,Y,+}(\Phi_k) \Big|_{X=Y=0},$$

$$L_{\text{hc}}^{kj} = \partial_{Y_j} \omega_{X,Y,+}(\Phi_k) \Big|_{X=Y=0},$$

$$L_{\text{ch}}^{kj} = \partial_{X_j} \omega_{X,Y,+}(\mathcal{J}_k) \Big|_{X=Y=0},$$

$$L_{\text{cc}}^{kj} = \partial_{Y_j} \omega_{X,Y,+}(\mathcal{J}_k) \Big|_{X=Y=0},$$

satisfy

$$\begin{aligned} L_{hh}^{kj} &= \mathfrak{L}(\Phi_k, \Phi_j), \\ L_{hc}^{kj} &= \mathfrak{L}(\Phi_k, \mathcal{J}_j), \\ L_{ch}^{kj} &= \mathfrak{L}(\mathcal{J}_k, \Phi_j), \\ L_{cc}^{kj} &= \mathfrak{L}(\mathcal{J}_k, \mathcal{J}_j). \end{aligned}$$

Assume in addition that (G2) and (G4) hold. Then

(2) The Green-Kubo formulas hold:

$$\begin{aligned} L_{hh}^{kj} &= \mathcal{L}(\Phi_k, \Phi_j), \\ L_{hc}^{kj} &= \mathcal{L}(\Phi_k, \mathcal{J}_j), \\ L_{ch}^{kj} &= \mathcal{L}(\mathcal{J}_k, \Phi_j), \\ L_{cc}^{kj} &= \mathcal{L}(\mathcal{J}_k, \mathcal{J}_j). \end{aligned}$$

(3) The Onsager reciprocity relations hold:

$$\begin{aligned} L_{hh}^{kj} &= L_{hh}^{jk}, \\ L_{cc}^{kj} &= L_{cc}^{jk}, \\ L_{hc}^{kj} &= L_{ch}^{jk}. \end{aligned}$$

The remark after Theorem 4.4 applies to Theorems 5.2 and 5.3.

In the literature one often considers a special case, described in the introduction, where one of the quantum dynamical systems, say $(\mathcal{O}_1, \tau_1, \omega_{1,\beta_1\mu_1})$, is finite dimensional and plays a role of a “small” quantum system \mathcal{S} coupled to reservoirs described by $(\mathcal{O}_j, \tau_j, \omega_{j,\beta_j\mu_j})$, $j \geq 2$. Such systems are one of the basic paradigms of non-equilibrium quantum statistical mechanics and have played an important role in the historical development of the subject. With regard to the algebraic approach described in this note, the only additional feature of these models is vanishing of heat and charge fluxes out of the small system: $\omega_{X,Y,+}(\Phi_1) = \omega_{X,Y,+}(\mathcal{J}_1) = 0$.

Many other generalizations are possible and it appears difficult to have a unified framework which covers all cases of physical interest. The Electronic Black Box Models studied in Refs. 6, 7, 21 are examples of open quantum systems which do not fit directly into the class of models described here (the non-interacting coupled system is not a tensor product of the individual subsystems). However,

the changes needed to apply our results to these models are elementary. One may also consider W^* -dynamical systems instead of C^* -dynamical systems and unbounded interactions which are only affiliated to the algebra of observables. The models where such generalization is necessary involve free bosonic reservoirs (a well-known example is the spin-boson model). One may also consider time-dependent interactions (see Refs. 3, 24, 25, 29). Another possible generalization involves more general gauge groups. The important point is that although all such generalizations may require some adjustment of technique and presentation, they bring nothing conceptually new.

ACKNOWLEDGMENTS

The research of V.J. was partly supported by NSERC. A part of this work has been done during the visit of V.J. to CPT-CNRS. Y.O. is supported by the Japan Society for the Promotion of Science. This work has been done during the stay of Y.O. to CPT-CNRS, partly supported by the Canon Foundation in Europe and JSPS.

REFERENCES

1. H. Araki, Relative entropy of states of von Neumann algebras, *Publ. Res. Inst. Math. Sci. Kyoto Univ.* **11**:809 (1975/76).
2. H. Araki, Relative entropy of states of von Neumann algebras II, *Publ. Res. Inst. Math. Sci. Kyoto Univ.* **13**:173 (1977/78).
3. W. K. Abou-Salem, and J. Fröhlich, Adiabatic Theorems and Reversible Isothermal Processes, *Lett. Math. Phys.*, (in press).
4. H. Araki, T. G. Ho, Asymptotic time evolution of a partitioned infinite two-sided isotropic XY -chain, *Tr. Mat. Inst. Steklova*, **228 Probl. Sovrem. Mat. Fiz.**, **203**, (2000); translation in Proc. Steklov Inst. Math. **228**, 191, (2000).
5. W. Aschbacher, and C.-A. Pillet, Non-equilibrium steady states of the XY chain *J. Stat. Phys.* **12**:1153 (2003).
6. W. Aschbacher, V. Jakšić, Y. Pautrat, and C.-A. Pillet, Topics in non-equilibrium quantum statistical mechanics. In *Open Quantum Systems III*. S. Attal, A. Joye, C.-A. Pillet editors. Lecture Notes in Mathematics **1882** (Springer, New York 2006).
7. W. Aschbacher, V. Jakšić, and Y. Pautrat, C.-A., Pillet, Transport properties of Ideal Fermi Gases (in preparation).
8. V. V. Aizenstadt, and V. A. Malyshev, Spin interaction with an Ideal fermi Gas. *J. Stat. Phys.* **48**:51 (1987).
9. D. D. Botvich, and V. A. Malyshev, Unitary equivalence of temperature dynamics for ideal and locally perturbed Fermi Gas, *Commun. Math. Phys.* **61**:209 (1978).
10. O. Bratteli, and D. W. Robinson, *Operator Algebras and Quantum Statistical Mechanics 1* (Springer-Verlag, Berlin, 1987).
11. O. Bratteli, and D. W. Robinson, *Operator Algebras and Quantum Statistical Mechanics 2nd edn* (Springer-Verlag, Berlin, 1996).
12. E. B. Davies, Markovian master equations. *Commun. Math. Phys.* **39**:91 (1974).

13. S. R. De Groot, and P. Mazur, *Non-Equilibrium Thermodynamics* (North-Holland, Amsterdam, 1969).
14. J. Dereziński, and V. Jakšić, C.-A. Pillet, Perturbation theory of W^* -dynamics, KMS-states and Liouvillean *Rev. Math. Phys.* **15**:447 (2003).
15. M. J. Donald, Relative Hamiltonians which are not bounded from above, *J. Func. Anal.* **91**:143 (1990).
16. D. J. Evans, and G. P. Morriss, *Statistical Mechanics of Nonequilibrium Liquids* (Academic Press, London, 1990).
17. J. Fröhlich, M. Merkli, and D. Ueltschi, Dissipative transport: thermal contacts and tunneling junctions, *Ann. Henri Poincaré* **4**:897 (2004).
18. V. Jakšić, Y. Ogata, and C.-A. Pillet, The Green-Kubo formula and Onsager reciprocity relations in quantum statistical mechanics. *Commun. Math. Phys.* (in press).
19. V. Jakšić, Y. Ogata, and C.-A. Pillet, The Green-Kubo Formula for the Spin-Fermion Model. Submitted.
20. V. Jakšić, Y. Ogata, and C.-A. Pillet, The Green-Kubo formula for locally interacting fermionic systems, In preparation.
21. V. Jakšić, Y. Ogata, and Y. Pautrat, C.-A. Pillet, The Green-Kubo formula for locally interacting fermionic systems, in preparation.
22. V. Jakšić, and C.-A. Pillet, On entropy production in quantum statistical mechanics, *Commun. Math. Phys.* **217**:285 (2001).
23. V. Jakšić, and C.-A. Pillet, Non-equilibrium steady states for finite quantum systems coupled to thermal reservoirs, *Commun. Math. Phys.* **226**:131 (2002).
24. V. Jakšić, and C.-A. Pillet, Mathematical theory of non-equilibrium quantum statistical mechanics, *J. Stat. Phys.* **108**:787 (2002).
25. V. Jakšić, and C.-A. Pillet, A note on the entropy production formula, *Contemp. Math.* **327**:175 (2003).
26. R. Kubo, M. Toda, and N. Hashitsune, *Statistical Physics* 2nd edition (Springer-Verlag, Berlin, 1991).
27. J. Lebowitz, and H. Spohn, Irreversible thermodynamics for quantum systems weakly coupled to thermal reservoirs, *Adv. Chem. Phys.* **39**:109 (1978).
28. L. Rey-Bellet, Open Classical Systems. In *Open Quantum Systems II*. S. Attal, A. Joye, C.-A. Pillet editors. Lecture Notes in Mathematics **1881**: (Springer, New York, 2006).
29. D. Ruelle, Natural nonequilibrium states in quantum statistical mechanics, *J. Stat. Phys.* **98**:57 (2000).
30. D. Ruelle, Entropy production in quantum spin systems, *Commun. Math. Phys.* **224**:3 (2001).
31. D. Ruelle, Topics in quantum statistical mechanics and operator algebras. Preprint, mp-arc 01-257 (2001).
32. D. Ruelle, Smooth dynamics and new theoretical ideas in nonequilibrium statistical mechanics, *J. Stat. Phys.* **95**:393 (1999).