

Sharp Lower Bounds for the Dimension of the Global Attractor of the Sabra Shell Model of Turbulence

Peter Constantin¹, Boris Levant² and Edriss S. Titi^{2,3}

Received November 1, 2006; accepted April 12, 2007

Published Online: May 5, 2007

In this work we derive lower bounds for the Hausdorff and fractal dimensions of the global attractor of the Sabra shell model of turbulence in different regimes of parameters. We show that for a particular choice of the forcing term and for sufficiently small viscosity term ν , the Sabra shell model has a global attractor of large Hausdorff and fractal dimensions proportional to $\log \nu^{-1}$ for all values of the governing parameter ε , except for $\varepsilon = 1$. The obtained lower bounds are sharp, matching the upper bounds for the dimension of the global attractor obtained in our previous work. Moreover, the complexity of the dynamics of the shell model increases as the viscosity ν tends to zero, and we describe a precise scenario of successive bifurcations for different parameters regimes. In the “three-dimensional” regime of parameters this scenario changes when the parameter ε becomes sufficiently close to 0 or to 1. We also show that in the “two-dimensional” regime of parameters, for a certain non-zero forcing term, the long-term dynamics of the model becomes trivial for every value of the viscosity.

KEY WORDS: turbulence, dynamic models, shell models, Navier–Stokes equations

AMS Subject Classifications: 76F20, 76D05, 35Q30

1. INTRODUCTION

Shell models of turbulence have attracted interest as useful phenomenological models that retain certain features of the Navier–Stokes equations (NSE). In this

¹Department of Mathematics, The University of Chicago, Chicago, IL 60637, USA; e-mail: const@math.uchicago.edu.

²Department of Computer Science and Applied Mathematics, Weizmann Institute of Science, Rehovot, 76100, Israel; e-mail: boris.levant@weizmann.ac.il.

³Department of Mathematics and, Department of Mechanical and Aerospace Engineering, University of California, Irvine, CA 92697, USA; e-mail: etiti@math.uci.edu and edriss.titi@weizmann.ac.il.

work we continue our analytical study of the Sabra shell model of turbulence that was introduced in Ref. 15. For other shell models see Refs. 4, 9, 10, 19. A recent review of the subject emphasizing the applications of the shell models to the study of the energy-cascade mechanism in turbulence can be found in Ref. 2.

The Sabra shell model of turbulence describes the evolution of complex Fourier-like components of a scalar velocity field denoted by u_n . The associated one-dimensional wave-numbers are denoted by k_n , where the discrete index n is referred to as the “shell index.” The equations of motion of the Sabra shell model of turbulence have the following form

$$\frac{du_n}{dt} = i(ak_{n+1}u_{n+2}u_{n+1}^* + bk_nu_{n+1}u_{n-1}^* - ck_{n-1}u_{n-1}u_{n-2}) - \nu k_n^2 u_n + f_n, \quad (1)$$

for $n = 1, 2, 3, \dots$, with the boundary conditions $u_{-1} = u_0 = 0$. The wave-numbers k_n are given by

$$k_n = k_0 \lambda^n, \quad (2)$$

with $\lambda > 1$ being the shell spacing parameter, and $k_0 > 0$. Although the equation does not capture any geometry, we will consider $L = k_0^{-1}$ as a fixed typical length scale of the model. In an analogy with the Navier–Stokes equations, $\nu > 0$ represents a kinematic viscosity and f_n are the Fourier components of the forcing term.

The three parameters of the model a , b and c are real. In order for the Sabra shell model to be a system of the hydrodynamic type we require that, in the inviscid ($\nu = 0$) and unforced ($f_n = 0$, $n = 1, 2, 3, \dots$) case, the model has at least one quadratic invariant. Requiring conservation of the energy

$$E = \sum_{n=1}^{\infty} |u_n|^2$$

leads to the following relation between the parameters of the model, which we will refer to as an energy conservation condition

$$a + b + c = 0. \quad (3)$$

Moreover, in the inviscid and unforced case, the model possesses another quadratic invariant

$$W = \sum_{n=1}^{\infty} \left(\frac{a}{c}\right)^n |u_n|^2.$$

The Sabra shell model (1) has the following six parameters: ν , λ , k_0 , a , b , and c . However, the “characteristic length-scale” k_0^{-1} does not appear on its own, but rather in the following combinations: $k_0 a$, $k_0 b$, and $k_0 c$. Therefore, without

loss of generality, we may assume that $k_0 = 1$. Next, by rescaling the time

$$t \rightarrow at,$$

and using the energy conservation assumption (3), we may set

$$a = 1, \quad b = -\varepsilon, \quad c = \varepsilon - 1.$$

Therefore, the Sabra shell model is in fact a three-parameter family of equations with parameters $\nu > 0$, ε , and $\lambda > 1$. In most of the numerical investigations of the shell models, the parameter λ is set to $\lambda = 2$ (see Refs. 4, 15). The physically relevant range of parameters is $|a/c| > 1$, or equivalently, $0 < \varepsilon < 2$ (see Ref. 15 for details). For $0 < \varepsilon < 1$, the quantity W is not sign-definite, and therefore it is common to associate it with the helicity—in an analogy to the 3D turbulence. The 2D parameters regime corresponds to $1 < \varepsilon < 2$, for which the quantity W becomes positive definite. In that case, the second conserved quadratic quantity W is identified with the enstrophy—in analogy to the 2D turbulence.

Classical theories of turbulence assert that the turbulent flows governed by the Navier–Stokes equations have a finite number of degrees of freedom (see, e.g., Refs. 9, 13). Arguing in the same vein, one can state that the Sabra shell model with non-zero viscosity has finitely many degrees of freedom. One of the ways to interpret such a physical statement mathematically is to assume that the number of degrees of freedom of the model corresponds to the Hausdorff or fractal dimension of its global attractor. Recall that the global attractor of the evolution equation on the space H is the maximal compact invariant subset of H , which attracts all bounded subsets of H . Thus the global attractor is unique. It can be shown that the global attractor is a nonempty connected subset of H , and in particular, it is a union of all bounded invariant attracting sets with different basins. For the precise definition and properties of the global attractor see Refs. 1, 20.

In our previous study of the Sabra shell model of turbulence,⁽⁶⁾ we proved the existence of a global attractor for the model and provided explicit upper bounds of its Hausdorff and fractal dimensions. Therefore, we proved that indeed the long-term dynamics of the Sabra shell model with non-zero viscosity has effectively finitely many degrees of freedom. The question remains, how many? The main motivation behind this work is to provide a lower bound for the Hausdorff and fractal dimensions of the global attractor. Namely, to show that for a particular choice of the forcing term, and for all $\varepsilon \in (0, 2)$, $\varepsilon \neq 1$, the Hausdorff and fractal dimensions of the global attractor are large, proportional to the upper bound obtained previously in Ref. 6. However, we also give an example of the forcing term such that for every $\varepsilon \in (1, 2)$ and for any non-zero viscosity ν the long-term dynamics of the Sabra shell model of turbulence is trivial.

In our work, we show that the Sabra shell model of turbulence possesses a global attractor of large dimension for all values of the parameter $\varepsilon \in (0, 2)$, $\varepsilon \neq 1$. In other words, we show that for every $\varepsilon \neq 1$, the Hausdorff and fractal dimensions

of the attractor are proportional to $\log_\lambda \nu^{-1}$ for small enough viscosity ν . Therefore, we extend and give a rigorous analytical justification for the numerical results observed in Refs. 21, 22 for $\varepsilon = 1/2$, and $\varepsilon = 3/2$, which corresponds respectively to the purely “three” and “two-dimensional” values of parameters.

Moreover, in Sec. 4, we obtain an estimate of the dimension of the global attractor in terms of the non-dimensional generalized Grashoff number G defined as

$$G = \frac{|\mathbf{f}|}{\nu^2 k_1^3}, \quad (4)$$

where $|\mathbf{f}|$ is an appropriate norm of the forcing term, which will be defined later. More specifically, we show that for every $\varepsilon \in (0, 2)$, $\varepsilon \neq 1$, and for a small enough viscosity ν , there exist positive constants c_1, c_2, c_3 , depending on λ, ε , and independent of the viscosity ν and the forcing term \mathbf{f} , such that

$$c_1 \log_\lambda G + c_2 \leq \dim_H(\mathcal{A}) \leq \dim_F(\mathcal{A}) \leq \frac{1}{2} \log_\lambda G + c_3. \quad (5)$$

The right-hand side of the inequality was proved in Ref. 6, and is true for every forcing term \mathbf{f} . In this work, we show that this estimate is tight in the sense that for particular choices of the forcing term \mathbf{f} , the lower bound in (5) is achieved. We would like to point out, that although our upper and lower bounds in (5) differ only by a multiplicative constant, the wave-numbers k_n (see (2)) of the Sabra shell model are exponential with n . Therefore, if we calculate the analog of an energy dissipation wave-number (see, e.g., Ref. 9) of the Sabra shell model, based on our estimation of the dimension of the global attractor, the gap between the upper and lower bounds becomes exponential.

When studying the lower-bounds for the dimension of the global attractors of the 2D NSE and related equations, one uses the force and the corresponding stationary solution concentrated on a single mode, the so-called Kolmogorov flow (see, e.g., Refs. 1, 14, 17). In the case of the shell model, the contiguous character of the interactions prevents the unstable manifold of a single mode solution from becoming large. Therefore, we constructed another kind of forcing term—the lacunary power-law forcing term—for which the lower bound in (5) is attained. However, this is not related to the power-law forcing term, used, e.g., in Ref. 18.

In Sec. 4.1, we study the linear stability of the stationary solution of the Sabra shell model, concentrated on a single mode N . We show that it becomes unstable for every N and for small enough viscosity for all $\varepsilon \in (0, 2)$, $\varepsilon \neq 1$, thus correcting a result of Ref. 12. Next, in Sec. 4.2 we construct the lacunary power-law forcing term and consider a corresponding stationary solution \mathbf{u} . We demonstrate how the size of the unstable manifold of \mathbf{u} grows both in the “two” and “three-dimensional” parameters regime, through successive bifurcations, as the viscosity ν tends to zero. In the “three-dimensional” regime $\varepsilon \in (0, 1)$, when

the parameter ε becomes close to 0 or 1, this scenario of successive bifurcations changes. More precisely, for a fixed viscosity, when ε crosses the values 0.05 and 0.97, the dimension of the unstable manifold of \mathbf{u} drops by a factor of 3. However, the attractor in those regimes is still of the size proportional to $\log_\lambda v^{-1}$.

The transition to chaos in the GOY shell model of turbulence was studied previously in Refs. 3, 11 by investigating numerically the stability properties of the special stationary solution corresponding to the single mode forcing term, which has a $k^{-1/3}$ Kolmogorov's power law in the inertial range. It was found that this solution becomes unstable at $\varepsilon = 0.3843$, and at some value of ε a phase transition occurs, when many stable directions become suddenly unstable. In this work, we find evidence that the nature of the transition to chaos might strongly depend on the type of the forcing term chosen. We expect that as ν tends to zero, the transition to chaotic behavior occurs in all regimes of the parameter ε , and this scenario changes for values of ε close to 0 and 1. This will be a subject of future investigation.

Finally in Sec. 5, we show that in the “two-dimensional” parameter regime the Sabra shell model has a trivial attractor reduced to a single equilibrium solution for any value of viscosity ν , when the forcing term is applied only to the first shell. This result is similar to the one for the 2-dimensional NSE due to Yudovich⁽²⁴⁾ and independently by Marchioro⁽¹⁶⁾ (see also Ref. 5).

First, we give a brief introduction to the mathematical formulations of the Sabra shell model problem. More details on this subject could be found in Refs. 6, 7.

2. PRELIMINARIES AND FUNCTIONAL SETTING

In this work, we will consider the real form of the Sabra model

$$\frac{du_n}{dt} = (ak_{n+1}u_{n+2}u_{n+1} + bk_nu_{n+1}u_{n-1} + ck_{n-1}u_{n-1}u_{n-2}) - \nu k_n^2 u_n + f_n,$$

for $n = 1, 2, 3, \dots$, and u_n, f_n are real for all n . This formulation is obtained from the original one by assuming that both the forcing term, f_n , and the velocity, u_n , components in Eq. (1) are purely imaginary. Our goal in this work is to show that the upper bounds of the Hausdorff and fractal dimensions of the global attractor of the Sabra shell model obtained in Ref. 6 are optimal in the sense that they can be achieved for some specific choice of the forcing term. Therefore, this formulation of the model is not restrictive, as long as we are able to show in that case the size of the global attractor matches the upper bound of Ref. 6.

Following the classical treatment of the NSE and Euler equations, and in order to simplify the notation, we write system (1) in the following functional

form

$$\frac{d\mathbf{u}}{dt} + \nu \mathbf{A}\mathbf{u} + \mathbf{B}(\mathbf{u}, \mathbf{u}) = \mathbf{f} \tag{6a}$$

$$\mathbf{u}(0) = \mathbf{u}^{\text{in}}, \tag{6b}$$

in a Hilbert space H . The linear operator \mathbf{A} as well as the bilinear operator \mathbf{B} will be defined below. In our case, the space H will be the sequences space ℓ^2 over the field of complex numbers \mathbb{C} . For every $\mathbf{u}, \mathbf{v} \in H$, the scalar product (\cdot, \cdot) and the corresponding norm $|\cdot|$ are defined as

$$(\mathbf{u}, \mathbf{v}) = \sum_{n=1}^{\infty} u_n v_n, \quad |\mathbf{u}| = \left(\sum_{n=1}^{\infty} |u_n|^2 \right)^{1/2}.$$

The linear operator $\mathbf{A} : D(\mathbf{A}) \rightarrow H$ is a positive definite, diagonal operator defined through its action on the sequence $\mathbf{u} = (u_1, u_2, \dots)$ by

$$\mathbf{A}\mathbf{u} = (k_1^2 u_1, k_2^2 u_2, \dots),$$

where the eigenvalues k_j^2 satisfy Eq. (2). Furthermore, we will need to define a space

$$V := D(\mathbf{A}^{1/2}) = \left\{ \mathbf{u} = (u_1, u_2, u_3, \dots) : \sum_{j=1}^{\infty} k_j^2 |u_j|^2 < \infty \right\}.$$

The bilinear operator $\mathbf{B}(\mathbf{u}, \mathbf{v}) = (B_1(\mathbf{u}, \mathbf{v}), B_2(\mathbf{u}, \mathbf{v}), \dots)$ will be defined formally in the following way. Let $\mathbf{u} = (u_1, u_2, \dots)$ and $\mathbf{v} = (v_1, v_2, \dots)$ be two sequences, then

$$B_n(\mathbf{u}, \mathbf{v}) = -k_n(\lambda v_{n+2} u_{n+1} - \varepsilon v_{n+1} u_{n-1} - \lambda^{-1} u_{n-1} v_{n-2} + \varepsilon \lambda^{-1} v_{n-1} u_{n-2}),$$

for $n = 1, 2, \dots$, where $u_0 = u_{-1} = v_0 = v_{-1} = 0$. It is easy to see that our definition of $\mathbf{B}(\mathbf{u}, \mathbf{v})$ is consistent with (1). In Ref. 6 we showed that indeed our definition of $\mathbf{B}(\mathbf{u}, \mathbf{v})$ makes sense as an element of H , whenever $\mathbf{u} \in H$ and $\mathbf{v} \in V$ or $\mathbf{u} \in V$ and $\mathbf{v} \in H$.

For more details on the material of this section see Refs. 6, 7.

3. LOWER BOUNDS FOR THE DIMENSION OF THE GLOBAL ATTRACTOR—THE “TWO-DIMENSIONAL” PARAMETER REGIME

The Hausdorff and fractal dimensions of the global attractor of the evolution equation are bounded from below by the dimension of the unstable manifold of every stationary solution (see, e.g., Refs. 1, 20). Therefore, in order to derive the lower bound for the Hausdorff and fractal dimensions of the global attractor of the Sabra shell model equation, we will construct a specific stationary solution of

Eq. (6a) and count the number of linearly unstable directions of that equilibrium. The same technique was first used in Ref. 17 (see also Refs. 1, 14) to obtain lower bounds for the dimension of the Navier–Stokes global attractor in 2D. In this section we will consider the “two-dimensional” parameters regime of the Sabra shell model corresponding to

$$1 < \varepsilon < 2.$$

Consider the forcing term

$$\mathbf{f} = (f_1, f_2, f_3, \dots),$$

where

$$f_n = \begin{cases} k_n^\alpha, & n = 0 \pmod 3 \\ 0, & \text{otherwise,} \end{cases} \tag{7}$$

for

$$\alpha = \frac{1}{3} \log_\lambda \frac{\varepsilon - 1}{\varepsilon} + \frac{5}{3}. \tag{8}$$

In addition, we will choose $M > 0$, large enough to be specified later in this section, such that $f_n = 0$, for all $n > M$.

The questions of global existence, uniqueness and regularity of solutions to problem (6a), with general forcing term $\mathbf{f} \in H$, are treated in Ref. 6. Specifically, for every initial condition $\mathbf{u}(0) \in H$ there is a unique solution, $u(t)$, of the Sabra shell model of turbulence which exists globally in time. Moreover, $\mathbf{u}(t) \in V$ for all $t > 0$. It was also established in Ref. 6 that for $\mathbf{f} \in H$ the Sabra shell model of turbulence has a global attractor, which is a compact subspace of the space V . In addition, in the case where the forcing term is supported on finite number of modes, such as in our case, the solution possess an exponentially decaying (with respect to k_n) dissipation range (see Ref. 6 for details).

The corresponding stationary solution of the Sabra shell model Eq. (1) or (6a) is

$$\mathbf{u} = (u_1, u_2, u_3, \dots), \tag{9}$$

where

$$u_n = \begin{cases} \frac{f_n}{\nu k_n^2}, & n = 0 \pmod 3 \\ 0, & \text{otherwise.} \end{cases} \tag{10}$$

Consider $\mathbf{v} = (v_1, v_2, v_3, \dots) \in H$ – an arbitrary perturbation of the stationary solution \mathbf{u} . Plugging $\mathbf{u} + \mathbf{v}$ into the equation of motion (6a), we find that the

perturbation \mathbf{v} satisfies the equation

$$\frac{d\mathbf{v}}{dt} + \nu \mathbf{A}\mathbf{v} + \mathbf{B}(\mathbf{u}, \mathbf{v}) + \mathbf{B}(\mathbf{v}, \mathbf{u}) + \mathbf{B}(\mathbf{v}, \mathbf{v}) = \mathbf{0}.$$

To study the linear stability of the equilibrium solution \mathbf{u} , we will consider the properties of the linearized equation

$$\frac{d\mathbf{v}}{dt} + \mathbf{L}_\mathbf{u}\mathbf{v} = \mathbf{0},$$

where the linear operator is defined as

$$\mathbf{L}_\mathbf{u}\mathbf{v} = \nu \mathbf{A}\mathbf{v} + \mathbf{B}(\mathbf{u}, \mathbf{v}) + \mathbf{B}(\mathbf{v}, \mathbf{u}). \tag{11}$$

We are looking for the solution of the eigenvalue problem

$$\mathbf{L}_\mathbf{u}\mathbf{v} = -\sigma\mathbf{v}, \tag{12}$$

for some $\sigma \in \mathbb{C}$. Our goal is to count the number of the solution of Eq. (12) with $Re(\sigma) > 0$. Equation (12) can be written componentwise as

$$\begin{aligned} \nu k_n^2 v_n - k_n(\lambda u_{n+2} v_{n+1} - \varepsilon u_{n+1} v_{n-1} + (\varepsilon - 1)\lambda^{-1} u_{n-1} v_{n-2} \\ + \lambda v_{n+2} u_{n+1} - \varepsilon v_{n+1} u_{n-1} + (\varepsilon - 1)\lambda^{-1} v_{n-1} u_{n-2}) = -\sigma v_n, \end{aligned} \tag{13}$$

where u_n is specified in (10). Note, that $u_n = 0$ for all $n \neq 0 \pmod 3$, therefore the last equation could be written in the following detailed form

- For $n = 0 \pmod 3$,

$$\nu k_n^2 v_n = -\sigma v_n. \tag{14}$$

- For $n = 1 \pmod 3$,

$$\nu k_n^2 v_n - k_n((\lambda u_{n+2} - \varepsilon u_{n-1})v_{n+1} + (\varepsilon - 1)\lambda^{-1} u_{n-1} v_{n-2}) = -\sigma v_n. \tag{15}$$

- For $n = 2 \pmod 3$,

$$\nu k_n^2 v_n - k_n(\lambda u_{n+1} v_{n+2} + ((\varepsilon - 1)\lambda^{-1} u_{n-2} - \varepsilon u_{n+1})v_{n-1}) = -\sigma v_n. \tag{16}$$

Note that from relation (14) it follows that $\sigma = -\nu k_{n_0}^2$ corresponds to the eigenvectors

$$\mathbf{v} = (0, \dots, 0, v_{n_0}, 0, \dots), \tag{17}$$

with $v_{n_0} \neq 0$ for some $n_0 = 0 \pmod 3$. However, we are only interested in the solutions of Eq. (12) with $Re(\sigma) > 0$. Based on the above, the only solution of relation (13) for which $Re(\sigma) > 0$ should satisfy $v_n = 0$, for all $n = 0 \pmod 3$. Equations (14) are not coupled with the rest of the recursive Eqs. (15) and (16).

Therefore, in looking for non-trivial solutions \mathbf{v} of Eq. (12) we can look only for the coupled recursive linear Eqs. (15) and (16), and set

$$v_n = 0, \quad \forall n = 0 \pmod 3, \tag{18}$$

as the solution of (14).

In what follows we will find sufficient conditions for the existence of non-trivial solutions for (12) with $Re(\sigma) > 0$. Let

$$b_{n,1} = k_n \frac{\lambda u_{n+2} - \varepsilon u_{n-1}}{\nu k_n^2 + \sigma},$$

$$c_{n,1} = \frac{(\varepsilon - 1)k_{n-1}u_{n-1}}{\nu k_n^2 + \sigma},$$

for all $n = 1 \pmod 3$, and

$$b_{n,2} = \frac{k_{n+1}u_{n+1}}{\nu k_n^2 + \sigma},$$

$$c_{n,2} = k_n \frac{(\varepsilon - 1)\lambda^{-1}u_{n-2} - \varepsilon u_{n+1}}{\nu k_n^2 + \sigma},$$

for all $n = 2 \pmod 3$. Then we can rewrite Eqs. (15) and (16) as a recursive relation for v_n

$$v_n - b_{n,1}v_{n+1} - c_{n,1}v_{n-2} = 0, \quad \text{for all } n = 1 \pmod 3,$$

$$v_n - b_{n,2}v_{n+2} - c_{n,2}v_{n-1} = 0, \quad \text{for all } n = 2 \pmod 3.$$

Due to our choice (8) for the value of α , one can realize from (10) that $c_{n,2} = 0$, for all $n = 2 \pmod 3$. Therefore, we can further simplify the last equations, which become

$$v_n - b_{n,1}v_{n+1} - c_{n,1}v_{n-2} = 0, \quad n = 1 \pmod 3, \tag{19}$$

$$v_n - b_{n,2}v_{n+2} = 0, \quad n = 2 \pmod 3.$$

The following result gives a sufficient condition for the last recursion to have at least one non-trivial solution.

Lemma 1. *Let M be a large positive integer. Let us fix $N < M$, and assume that $N = 1 \pmod 3$. Then the recursive Eq. (19) has a non-trivial solution of the form $v_n = 0$, for all $n > N$, and $v_n \neq 0$, for some $n \leq N$, if and only if*

$$b_{N-2,2} c_{N,1} = 1. \tag{20}$$

Proof: The proof of the Lemma 1 is simple once we observe that the recursive relation (19) for solutions of the form $v_n = 0$, for $n > N$, becomes

$$\begin{aligned} v_1 - b_{1,1}v_2 &= 0, \\ v_2 - b_{2,2}v_4 &= 0, \\ v_4 - b_{4,1}v_5 - c_{4,1}v_2 &= 0, \\ v_5 - b_{5,2}v_7 &= 0, \\ &\vdots \\ v_{N-2} - b_{N-2,2}v_N &= 0, \\ v_N - c_{N,1}v_{N-2} &= 0. \end{aligned}$$

The last two equations have a one-parameter family of nontrivial solutions if and only if the condition (20) is satisfied. □

Finally, we are ready to prove the main result of this section.

Theorem 1. *The Hausdorff and fractal dimensions of the global attractor of the Eq. (6a) in the parameter regime $1 < \varepsilon < 2$, with the forcing term f specified in (7), satisfy*

$$\dim_F \mathcal{A} \geq \dim_H \mathcal{A} \geq \frac{2}{4 - \log_\lambda \frac{\varepsilon-1}{\varepsilon}} \log_\lambda v^{-1} + \frac{1}{8 - 2 \log_\lambda \frac{\varepsilon-1}{\varepsilon}} \log_\lambda(\varepsilon - 1). \tag{21}$$

Proof: Fix M to be large enough, and let $N < M$ be such that $N = 1 \pmod 3$. Suppose, that for such N the condition (20) is satisfied for certain $\sigma = \sigma(N)$, depending on N , for which $Re(\sigma) > 0$. Then, for such σ , there exists a solution of Eq. (19), and in particular, there exists a solution of the eigenvalue problem (12) with $Re(\sigma) > 0$. Moreover, it is not hard to see that if $N_1 \neq N_2$, then the solutions of the eigenvalue problem (12) corresponding to $\sigma(N_1)$ and $\sigma(N_2)$ are different.

Therefore, for a given M , in order to count the number of unstable directions of the stationary solution (10), we need to count the number of N s, such that (i) $N < M$; (ii) $N = 1 \pmod 3$; (iii) N satisfies (20) with the eigenvalue σ , for which $Re(\sigma) > 0$.

Let us fix $N > 0$, satisfying $N = 1 \pmod 3$. The condition (20) becomes

$$\frac{k_{N+1}u_{N+1}}{vk_N^2 + \sigma} \cdot \frac{(\varepsilon - 1)k_{N+1}u_{N+1}}{vk_{N+2}^2 + \sigma} = 1.$$

We get the quadratic equation in σ

$$\sigma^2 + (vk_N^2 + vk_{N+2}^2)\sigma + v^2k_N^2k_{N+2}^2 - (\varepsilon - 1)k_{N+1}^2u_{N+1}^2 = 0.$$

This equation has a real positive solution, provided

$$v^2 k_N^2 k_{N+2}^2 - (\varepsilon - 1) k_{N+1}^2 u_{N+1}^2 < 0. \tag{22}$$

Substituting (10) we obtain the equivalent condition to (22)

$$(\varepsilon - 1) v^{-2} k_{N+1}^{2(\alpha-2)} > v^2 k_{N+1}^2.$$

Rearranging terms, the following condition guarantees the existence of a positive real eigenvalue for (12)

$$(\varepsilon - 1)^{1/2} v^{-2} > \lambda^{(3-\alpha)(N+1)}.$$

Now, we substitute the value of α from (8) to obtain

$$(\varepsilon - 1)^{1/2} v^{-2} > \lambda^{\frac{4-\log_\lambda \frac{\varepsilon-1}{\varepsilon}}{3}(N+1)}.$$

Finally, we get the estimate

$$\begin{aligned} N + 1 &< \frac{3}{4 - \log_\lambda \frac{\varepsilon-1}{\varepsilon}} \log_\lambda ((\varepsilon - 1)^{1/2} v^{-2}) \\ &= \frac{6}{4 - \log_\lambda \frac{\varepsilon-1}{\varepsilon}} \log_\lambda v^{-1} + \frac{3}{8 - 2 \log_\lambda \frac{\varepsilon-1}{\varepsilon}} \log_\lambda (\varepsilon - 1). \end{aligned} \tag{23}$$

Therefore, we showed that if the M that we have chosen at the beginning of the proof, is larger than the right-hand side relation (23), then for such a choice of the forcing term, the number of unstable directions of the stationary solution (10) is bounded from below by

$$\frac{2}{4 - \log_\lambda \frac{\varepsilon-1}{\varepsilon}} \log_\lambda v^{-1} + \frac{1}{8 - 2 \log_\lambda \frac{\varepsilon-1}{\varepsilon}} \log_\lambda (\varepsilon - 1),$$

and the statement of the theorem follows. □

In Ref. 6 we showed that the dimension of the global attractor of the Sabra shell model of turbulence is proportional to the $\log_\lambda v^{-1}$ for small enough viscosity ν . Therefore, our result proves that this bound is tight.

4. LOWER BOUNDS FOR THE DIMENSION OF THE GLOBAL ATTRACTOR—THE “THREE-DIMENSIONAL” PARAMETERS REGIME

The result obtained in the previous section did not give an answer for the case

$$0 < \varepsilon < 1, \tag{24}$$

which is also known as the “three-dimensional” range of parameters. Therefore, we will need to apply a different strategy. First, we will consider the linear stability

of a stationary solution, corresponding to the force acting on a single mode number N , for some $N > 0$. We will show that for every choice of N , and for every value of the parameter $\varepsilon \in (0, 2]$, $\varepsilon \neq 1$, such a stationary solution becomes linearly unstable for sufficiently small viscosity ν . The stability of a single-mode stationary solution was numerically studied previously in Ref. 12, where it was found that such a solution becomes stable around $\varepsilon = 1$. Our rigorous proof contradicts this numerical observation.

Next, we will construct a special type of equilibrium solution, for which we will be able to count the number of unstable directions. The drawback of this method is that we are not able to obtain the exact dependence of the bounds on ε and λ , the parameters of the problem, as we succeeded to do in the “two-dimensional” parameters case.

4.1. On the Linear Stability of a “Single-Mode” Flow

Let us fix $N \geq 1$ and consider the forcing term acting on the single mode N of the form

$$\mathbf{f}^N = (0, \dots, 0, \nu k_N^{-1}, 0, \dots), \tag{25}$$

where all the components of \mathbf{f}^N , except the N -th, are zero. Consider one particular choice of an equilibrium solution corresponding to the above forcing term

$$\mathbf{u}^N = (0, \dots, 0, k_N^{-3}, 0, \dots), \tag{26}$$

which is the analog of the Kolmogorov flow for the Navier–Stokes equations.

Linearizing Eq. (1) around the equilibrium solution \mathbf{u}^N and writing Eq. (12) in the component form we get the following set of equations. For every $j \in \mathbb{N}$, satisfying $2 < |j - N|$, or for $j = N$, we have

$$\nu k_j^2 v_j = -\sigma v_j, \tag{27}$$

together with the four equations, coming from the nonlinear interaction with \mathbf{u}^N

$$\begin{aligned} \nu k_{N-2}^2 v_{N-2} - k_{N-1} k_N^{-3} v_{N-1} &= -\sigma v_{N-2}, \\ \nu k_{N-1}^2 v_{N-1} - k_N k_N^{-3} v_{N+1} + \varepsilon k_{N-1} k_N^{-3} v_{N-2} &= -\sigma v_{N-1} \\ \nu k_{N+1}^2 v_{N+1} + (1 - \varepsilon) k_N k_N^{-3} v_{N-1} + \varepsilon k_{N+1} k_N^{-3} v_{N+2} &= -\sigma v_{N+1}, \\ \nu k_{N+2}^2 v_{N+2} + (1 - \varepsilon) k_{N+1} k_N^{-3} v_{N+1} &= -\sigma v_{N+2}. \end{aligned}$$

Therefore, the eigenvalues of the linear operator $\mathbf{L}_{\mathbf{u}^N}$ (see (11)) are $-\sigma = \nu k_j^2$, for $2 < |j - N|$, or for $j = N$, corresponding to the eigenvectors $\mathbf{v} = (0, 0, \dots, 1, 0, \dots)$, with 1 at the j -th place. Clearly, those eigenvalues are positive, corresponding to $Re(\sigma) < 0$, therefore they do not contribute to the number of linearly unstable directions of the equilibria \mathbf{u}^N .

Other eigenvalues of the linear operator $\mathbf{L}_{\mathbf{u}^N}$ are the eigenvalues of the following matrix

$$J_N = \begin{pmatrix} vk_{N-2}^2 & -k_N^{-2}\lambda^{-1} & 0 & 0 \\ \varepsilon k_N^{-2}\lambda^{-1} & vk_{N-1}^2 & -k_N^{-2} & 0 \\ 0 & (1-\varepsilon)k_N^{-2} & vk_{N+1}^2 & \varepsilon k_N^{-2}\lambda \\ 0 & 0 & (1-\varepsilon)k_N^{-2}\lambda & vk_{N+2}^2 \end{pmatrix}, \tag{28}$$

which will correspond to the eigenvectors $\mathbf{v} = (v_1, v_2, v_3, \dots)$ of the linear operator $\mathbf{L}_{\mathbf{u}^N}$ with the only non-zero components being $v_j, 0 < |j - N| < 2$.

Our goal is to find the condition on the parameters $N, \varepsilon,$ and $v,$ such that the matrix J_N has eigenvalues with a negative real part, which will correspond to σ satisfying $Re(\sigma) > 0$. Let us rewrite expression (28) in the following way

$$J_N = k_N^{-2} \cdot \begin{pmatrix} \lambda^{-4}\beta & -\lambda^{-1} & 0 & 0 \\ \varepsilon\lambda^{-1} & \lambda^{-2}\beta & -1 & 0 \\ 0 & (1-\varepsilon) & \lambda^2\beta & \varepsilon\lambda \\ 0 & 0 & (1-\varepsilon)\lambda & \lambda^4\beta \end{pmatrix}, \tag{29}$$

where we denoted for simplicity

$$\beta = vk_N^4.$$

First, by substituting $\varepsilon = 1,$ we find that for this value of ε the eigenvalues of the matrix J_N always has a positive real part. Therefore, we conclude that in the case $\varepsilon = 1$ the solution \mathbf{u}^N is stable for every N and any $v.$

For other values of the parameters, we substitute $\lambda = 2,$ and write the characteristic polynomial of the matrix J_N

$$\begin{aligned} x^4 - \frac{325}{16}\beta x^3 + \left(4\varepsilon^2 - \frac{19}{4}\varepsilon + 1 + \frac{4497}{64}\beta^2\right)x^2 \\ - \left(\frac{325}{16}\beta^2 + \frac{5}{4}\varepsilon^2 + \frac{257}{16} - \frac{197}{16}\varepsilon\right)\beta x \\ + (\varepsilon^3 - \varepsilon^2) + \left(1 + \frac{239}{16}\varepsilon + \frac{1}{16}\varepsilon^2 + \beta^2\right)\beta^2 = 0. \end{aligned} \tag{30}$$

Next, by fixing an $\varepsilon,$ we find the largest β when the real part of the roots of the polynomial (30) changes its sign. The result of this calculation is shown in Fig. 1.

For every ε in the “three-dimensional” subrange of parameters $0.05 < \varepsilon < 0.97,$ there are two bifurcation points. First, there exists a value of β for which one of the real eigenvalues of J_N crosses 0 and becomes negative. Decreasing the parameter β further, we observe another bifurcation for which the real part of a




Fig. 1. The bifurcation diagram β vs. ε . The dashed line indicates the appearance of one real negative eigenvalue of the matrix J_N , which happens only in the regime $0 < \varepsilon < 1$ (a). The solid line shows the point at which the real part of two conjugate complex eigenvalues of J_N become negative. This bifurcation disappears in the “three-dimensional” parameter regime (a) at $0 \leq \varepsilon \leq 0.05$ and $0.97 \leq \varepsilon \leq 1$. Observe that for positive viscosity ν the solution \mathbf{u}^N is linearly stable for every N at $\varepsilon = 1$, which can be shown rigorously.

pair of complex conjugate eigenvalues becomes negative. Therefore, for $0 < \varepsilon < 1$ there exists a function $m(\varepsilon)$, such that the matrix J_N has 3 negative eigenvalues for

$$0 < \beta = \nu k_N^4 \leq m(\varepsilon), \tag{31}$$

or in other words, for N satisfying

$$4N \leq \log_\lambda \nu^{-1} + \log_\lambda m(\varepsilon). \tag{32}$$

Note, that $m(\varepsilon) > 0$ at $\varepsilon \in (0.05, 0.97)$, and $m(\varepsilon) = 0$ otherwise.

For the range of parameters $0 < \varepsilon \leq 0.05$ and $0.97 \leq \varepsilon < 1$ we observe only one bifurcation point at which one of the real eigenvalues becomes negative.

Finally, for the “two-dimensional” range of parameters $1 < \varepsilon < 2$ the scenario is a little different, as we observe only one bifurcation point for which the real part of a pair of complex conjugate eigenvalues becomes negative. Namely, for $1 < \varepsilon < 2$ there exists a function $d(\varepsilon)$, such that the matrix J_N has two negative eigenvalues for

$$0 < \beta = \nu k_N^4 \leq d(\varepsilon), \tag{33}$$

or equivalently, for N satisfying

$$4N \leq \log_\lambda \nu^{-1} + \log_\lambda d(\varepsilon). \tag{34}$$

In this case, we also have, $d(\varepsilon) > 0$ for all $\varepsilon \in (1, 2]$, and $d(1) = 0$.

4.2. Calculating the Lower Bound of the Dimension of the Attractor

When studying the lower-bounds for the dimension of the global attractors of the 2D NSE and related equations one uses the force and the corresponding stationary solution concentrated on a single mode, the so-called Kolmogorov flow (see, e.g., Refs. 1, 14, 17). In the case of the shell model, the contiguous character of the interactions prevents the unstable manifold of a single mode solution from becoming large. In the previous section we show that the Sabra shell model has at most three unstable directions for the “single-mode” forcing term. In this section we construct a special kind of a lacunary power-law forcing term, and a corresponding stationary solution, with a large unstable manifold. Let us introduce

$$\mathbf{g} = \sum_{j=1}^{\infty} \mathbf{f}^{5j}, \quad |\mathbf{g}| = \nu \frac{1}{\lambda^{5} \sqrt{1 - \lambda^{-10}}}, \tag{35}$$

where \mathbf{f}^{5j} is defined in (25). Then the stationary solution corresponding to that forcing term is

$$\mathbf{u}^{\mathbf{g}} = \sum_{j=1}^{\infty} \mathbf{u}^{5j}, \tag{36}$$

where \mathbf{u}^{5j} is defined in (26). Using the results of the previous section on the stability of the single-mode stationary solution we conclude that for $0 < \varepsilon < 1$, the number of the unstable directions of the solution $\mathbf{u}^{\mathbf{g}}$ equals to $3N/5$, where N satisfies the relation (32). On the other hand, the number of the unstable directions of the solution $\mathbf{u}^{\mathbf{g}}$ for $1 < \varepsilon < 2$ equals to $2N/5$, where N satisfies relation (34).

Recall the definition of the generalized Grashoff number (4), which in our case satisfies

$$G = \frac{|\mathbf{g}|}{\nu^2 \lambda^3} = \frac{1}{\nu \lambda^8 \sqrt{1 - \lambda^{-10}}}.$$

Therefore, we can rewrite the bounds (32) and (32) in terms of the generalized Grashoff number to obtain

$$4N = \log_{\lambda} \nu^{-1} + \log_{\lambda} f(\varepsilon) \leq \log_{\lambda} G + \log_{\lambda} f(\varepsilon), \tag{37}$$

where $f(\varepsilon)$ denotes $m(\varepsilon)$ or $d(\varepsilon)$. Therefore, we proved the following.

Theorem 2. *The Hausdorff and fractal dimensions of the global attractor \mathcal{A} of the Sabra shell model of turbulence with $\nu > 0$ and the forcing term defined in (35) satisfy*

$$\dim_F(\mathcal{A}) \geq \dim_H(\mathcal{A}) \geq K \log_{\lambda} G + \log_{\lambda} f(\varepsilon), \tag{38}$$

for the positive constant K depending on ε which satisfies

$$K(\varepsilon) = \begin{cases} \frac{3}{20}, & \text{for } 0.05 < \varepsilon < 0.97 \\ \frac{1}{10}, & \text{for } 1 < \varepsilon \leq 2 \\ \frac{1}{20}, & \text{for } 0 < \varepsilon \leq 0.05, \quad \text{and } 0.97 \leq \varepsilon < 1, \end{cases} \quad (39)$$

and for some positive real function $f(\varepsilon)$, which is 0 only for $\varepsilon = 1$.

The lower bounds for the global attractor, given by the last theorem do not match exactly the upper bounds, that were obtained previously in Ref. 6, exactly, namely

$$\dim_H(\mathcal{A}) \leq \dim_F(\mathcal{A}) \leq \frac{1}{2} \log_\lambda G - C(\varepsilon), \quad (40)$$

where the function $C(\varepsilon)$ stays positive and bounded for every $\varepsilon \in (0, 2)$. Moreover, the constant K in front of the $\log_\lambda G$ term, although it can be slightly improved, cannot be brought much closer to $\frac{1}{2}$ to match the upper bound of (40).

5. EXISTENCE OF A TRIVIAL GLOBAL ATTRACTOR FOR ANY VALUE OF ν

It is well known that the attractor for the 2-dimensional space-periodic Navier–Stokes equation with a particular form of the forcing term can consist of only one function. This well-known example is due to Yudovich⁽²⁴⁾ and independently by Marchioro⁽¹⁶⁾ (for the proof see also Ref. 8). The same is true for the Sabra shell model for $1 < \varepsilon < 2$, therefore, we need to stress that the bounds that we obtained for the dimension of the global attractor are valid only for the particular type of forcing term that we used in our calculations.

We mentioned in the introduction that, for the 2-dimensional parameters regime, the inviscid Sabra shell model without forcing term conserves the following quantity

$$|A^\gamma u|^2 = \sum_{n=1}^{\infty} k_n^{4\gamma} |u_n|^2,$$

for $4\gamma = -\log_\lambda(\varepsilon - 1)$. This is because we have, in this case, the following identity

$$\text{Re}(\mathbf{B}(\mathbf{u}, \mathbf{u}), \mathbf{A}^{2\gamma} \mathbf{u}) = 0.$$

For $m > 0$ let us denote by \mathbf{P}_m the projection onto the first m coordinates of the sequence \mathbf{u} , and by $\mathbf{Q}_m = \mathbf{I} - \mathbf{P}_m$.

Theorem 3. *Suppose that the forcing term f acts only on the N -th shell for some $N \geq 1$. Let $\mathbf{u}(t)$ be the solution of Eq. (6a) in the “two-dimensional” regime of parameters $1 < \varepsilon < 2$, and let $\gamma = -\frac{1}{4} \log_\lambda(\varepsilon - 1)$. Then we have*

$$\limsup_{t \rightarrow \infty} |\mathbf{Q}_m \mathbf{u}(t)|^2 \leq C \frac{1}{k_{m+1}^{4\gamma}} \liminf_{t \rightarrow \infty} |\mathbf{P}_m \mathbf{u}(t)|^2, \tag{41}$$

for $C = \frac{k_N^{4\gamma} - k_1^{4\gamma}}{1 - \lambda^{-4\gamma}}$ and $m \geq N$.

Proof: Taking the scalar product of Eq. (6a) with \mathbf{u} and with $\mathbf{A}^{2\gamma} \mathbf{u}$ we get two equations

$$\frac{1}{2} \frac{d}{dt} |\mathbf{u}|^2 + \nu(\mathbf{A}\mathbf{u}, \mathbf{u}) = \text{Re}(f_N u_N^*),$$

and

$$\frac{1}{2} \frac{d}{dt} |\mathbf{A}^\gamma \mathbf{u}|^2 + \nu(\mathbf{A}\mathbf{u}, \mathbf{A}^{2\gamma} \mathbf{u}) = \text{Re}(k_N^{4\gamma} f_N u_N^*).$$

Multiplying the energy equation by $k_N^{4\gamma}$ and subtracting it from the last equation we get

$$\frac{1}{2} \frac{d}{dt} (|\mathbf{A}^\gamma \mathbf{u}|^2 - k_N^{4\gamma} |\mathbf{u}|^2) + \nu(|\mathbf{A}^{\gamma+1/2} \mathbf{u}|^2 - k_N^{4\gamma} \|\mathbf{u}\|^2) = 0. \tag{42}$$

On the other hand,

$$\begin{aligned} |\mathbf{A}^{\gamma+1/2} \mathbf{u}|^2 - k_N^{4\gamma} \|\mathbf{u}\|^2 &= \sum_{n=1}^{\infty} k_n^2 (k_n^{4\gamma} - k_N^{4\gamma}) |u_n|^2 \\ &\geq k_N^2 \sum_{n=1}^{\infty} (k_n^{4\gamma} - k_N^{4\gamma}) |u_n|^2 = k_N^2 (|\mathbf{A}^\gamma \mathbf{u}|^2 - k_N^{4\gamma} |\mathbf{u}|^2). \end{aligned}$$

Plugging the last expression into (42) yields

$$\frac{d}{dt} (|\mathbf{A}^\gamma \mathbf{u}|^2 - k_N^{4\gamma} |\mathbf{u}|^2) \leq -2\nu k_N^2 (|\mathbf{A}^\gamma \mathbf{u}|^2 - k_N^{4\gamma} |\mathbf{u}|^2),$$

and therefore,

$$\limsup_{t \rightarrow \infty} (|\mathbf{A}^\gamma \mathbf{u}(t)|^2 - k_N^{4\gamma} |\mathbf{u}(t)|^2) = 0. \tag{43}$$

Finally,

$$|\mathbf{Q}_m \mathbf{u}|^2 = \sum_{i=m+1}^{\infty} |u_i|^2 = \sum_{i=m+1}^{\infty} \frac{k_i^{4\gamma} - k_N^{4\gamma}}{k_i^{4\gamma} - k_N^{4\gamma}} |u_i|^2$$

$$\begin{aligned} &\leq \frac{1}{k_{m+1}^{4\gamma} - k_N^{4\gamma}} (|\mathbf{Q}_m \mathbf{A}^\gamma \mathbf{u}|^2 - k_N^{4\gamma} |\mathbf{Q}_m \mathbf{u}|^2) \\ &= \frac{1}{k_{m+1}^{4\gamma} - k_N^{4\gamma}} ((|\mathbf{A}^\gamma \mathbf{u}|^2 - k_N^{4\gamma} |\mathbf{u}|^2) - (|\mathbf{P}_m \mathbf{A}^\gamma \mathbf{u}|^2 - k_N^{4\gamma} |\mathbf{P}_m \mathbf{u}|^2)) \\ &\leq \frac{1}{k_{m+1}^{4\gamma} - k_N^{4\gamma}} ((|\mathbf{A}^\gamma \mathbf{u}|^2 - k_N^{4\gamma} |\mathbf{u}|^2) + (k_N^{4\gamma} - k_1^{4\gamma}) |\mathbf{P}_m \mathbf{u}|^2), \end{aligned}$$

and the result follows from (43). □

Corollary 1. *The global attractor of the Sabra shell model of turbulence in the “two-dimensional” regime of parameters $1 < \varepsilon < 2$ with the force applied only to the first shell*

$$\mathbf{f}^1 = (f, 0, 0, \dots), \tag{44}$$

is reduced to a single stationary solution

$$\mathbf{u}^1 = \left(\frac{f}{\nu k_1^2}, 0, 0, \dots \right).$$

Proof: Let $\mathbf{u} = (u_1, u_2, \dots)$ be a solution of the Sabra shell model with the forcing term \mathbf{f} defined by (44). Then it immediately follows from Theorem 3 that

$$\limsup_{t \rightarrow \infty} |\mathbf{Q}_1 \mathbf{u}|^2 = 0,$$

which means that $\limsup_{t \rightarrow \infty} |u_n| = 0$, for every $n \geq 2$.

Define $\mathbf{v} = (v_1, v_2, \dots)$ as $\mathbf{v} = \mathbf{u} - \mathbf{u}^1$, which satisfies the equation

$$\frac{d\mathbf{v}}{dt} + \nu \mathbf{A}\mathbf{v} + \mathbf{B}(\mathbf{u}, \mathbf{u}) = 0,$$

where we used the fact that $\mathbf{B}(\mathbf{u}^1, \mathbf{u}^1) = 0$. Taking the inner product of the equation with the vector $\mathbf{P}_1 \mathbf{v} = (v_1, 0, 0, \dots)$ we get that $|v_1(t)|^2$ satisfies

$$\frac{1}{2} \frac{d}{dt} |v_1(t)|^2 + \nu k_1^2 |v_1(t)|^2 + v_1(t) u_2(t) u_3(t) = 0.$$

Using the fact that $u_2(t), u_3(t)$ tend to 0 as $t \rightarrow \infty$ we conclude that $|v_1(t)|^2 \rightarrow 0$ as $t \rightarrow \infty$. Therefore,

$$\limsup_{t \rightarrow \infty} |\mathbf{v}|^2 = \limsup_{t \rightarrow \infty} |\mathbf{u} - \mathbf{u}^1|^2 = 0.$$

finishing the proof. □

6. CONCLUSION

In this work we obtained lower bounds for the dimension of the global attractor of the Sabra shell model of turbulence for specific choices of the forcing term. Our main result states that for these specific choices of the forcing term the Sabra shell model has a large attractor for all values of the governing parameter $\varepsilon \in (0, 2) \setminus \{1\}$. We also showed the scenario of the transition to chaos in the model, which is slightly different for the two- and three-dimensional parameter regime. In addition, in the three-dimensional parameters regime, $\varepsilon \in (0, 1)$, we found that when the parameter ε becomes sufficiently close to 0 or to 1, where the chaotic behavior in the vicinity of the stationary solution changes dramatically.

Finally, we show that in the “two-dimensional” parameters regime the Sabra shell model has a trivial attractor reduced to a single equilibrium solution for any value of viscosity ν , when the forcing term is applied only to the first shell. This result is true also for the two-dimensional NSE due to Yudovich⁽²⁴⁾ and independently by Marchioro⁽¹⁶⁾ (see also Ref. 8).

ACKNOWLEDGMENTS

The authors would like to thank U. Frisch for a very stimulating discussion, and his valuable comments on the first version of the manuscript. The work of P.C. was supported in part by the NSF grant No. DMS-0504213. The work of E.S.T. was also supported in part by the NSF grant No. DMS-0504619, the MAOF Fellowship of the Israeli Council of Higher Education, the ISF grant No. 120/06, and by the BSF grant No. 2004271.

REFERENCES

1. A. V. Babin and M. I. Vishik, Attractors of partial differential equations and estimates of their dimension. *Uspekhi Mat. Nauk* **38**:133–187 (1983) (in Russian); *Russian Math. Surveys* **38**: 151–213 (in English).
2. L. Biferale, Shell models of energy cascade in turbulence. *Annual Rev. Fluid Mech.* **35**:441–468 (2003).
3. L. Biferale, A. Lambert, R. Lima and G. Paladin, Transition to chaos in a shell model of turbulence. *Physica D* **80**:105–119 (1995).
4. T. Bohr, M. H. Jensen, G. Paladin and A. Vulpiani, *Dynamical Systems Approach to Turbulence* (Cambridge University Press, 1998.)
5. P. Constantin, C. Foias, and R. Temam, On the dimension of the attractors in two-dimensional turbulence. *Physica D* **30**:284–296 (1988).
6. P. Constantin, B. Levant and E. S. Titi, Analytic study of the shell model of turbulence. *Physica D* **219**:120–141 (2006).
7. P. Constantin, B. Levant and E. S. Titi, Regularity of inviscid shell models of turbulence. *Phys. Rev. E* **75**(1):016304 (2007).
8. C. Foias, O. Manley, R. Rosa and R. Temam, *Navier–Stokes Equations and Turbulence* (Cambridge University Press, 2001).

9. U. Frisch, *Turbulence: The Legacy of A. N. Kolmogorov* (Cambridge University Press, 1995).
10. E. B. Gledzer, System of hydrodynamic type admitting two quadratic integrals of motion. *Sov. Phys. Dokl.* **18**:216–217 (1973).
11. L. Kadanoff, D. Lohse and N. Schröghofer, Scaling and linear response in the GOY turbulence model. *Physica D* **100**:165–186 (1997).
12. J. Kockelkoren, F. Okkels and M. H. Jensen, Chaotic behavior in shell models and shell maps. *J. Stat. Phys.* **93**:833 (1998).
13. L. D. Landau and E. M. Lifschitz, *Fluid Mechanics* (Pergamon, Oxford, 1977).
14. V. X. Liu, A sharp lower bound for the Hausdorff dimension of the global attractors of the 2D Navier–Stokes equations. *Commun. Math. Phys.* **158**:327–339 (1993).
15. V. S. L'vov, E. Podivilov, A. Pomyalov, I. Procaccia and D. Vandembroucq, Improved shell model of turbulence. *Physical Review E*. **58**(2):1811–1822 (1998).
16. C. Marchioro, An example of absence of turbulence for any Reynolds number. *Comm. Math. Phys.* **105**:99–106 (1986).
17. L. D. Meshalkin and Y. G. Sinai, Investigation of the stability of a stationary solution of a system of equations for the plane movement of an incompressible viscous liquid. *J. Appl. Math. Mech.* **25**:1700–1705 (1961).
18. D. Mitra, J. Bec, R. Pandit and U. Frisch, Is multiscaling an artifact in the stochastically forced burgers equation? *Phys. Rev. Lett.* **95**:194501 (2005).
19. K. Okhitani, M. Yamada Temporal intermittency in the energy cascade process and local Lyapunov analysis in fully developed model of turbulence. *Prog. Theor. Phys.* **89**:329–341 (1989).
20. R. Temam, *Infinite-Dimensional Dynamical Systems in Mechanics and Physics* (Springer-Verlag, New-York, 1988).
21. M. Yamada, and K. Okhitani, Lyapunov spectrum of a chaotic model of three-dimensional turbulence. *J. Phys. Soc. Jpn.* **56**:4210–4213 (1987).
22. M. Yamada and K. Okhitani, Lyapunov spectrum of a model of two-dimensional turbulence. *Phys. Rev. Lett.* **60**(11):983–986 (1988).
23. M. Yamada and K. Okhitani, Asymptotic formulas for the Lyapunov spectrum of fully developed shell model turbulence. *Phys. Rev. E* **57**(6):57–60 (1998).
24. V. I. Yudovich, Example of the generation of a secondary stationary or periodic flow when there is loss of stability of the laminar flow of a viscous incompressible fluid. *J. Appl. Math. Mech.* **29**:527–544 (1965).