

Decay Properties of the Connectivity for Mixed Long Range Percolation Models on \mathbb{Z}^d

Gastão A. Braga · Leandro M. Ciolleti · Rémy Sanchis

Received: 20 July 2006 / Accepted: 10 May 2007 /
Published online: 27 June 2007
© Springer Science+Business Media, LLC 2007

Abstract In this short note we consider mixed short-long range independent bond percolation models on \mathbb{Z}^{k+d} . Let p_{uv} be the probability that the edge (u, v) will be open. Allowing a x, y -dependent length scale and using a multi-scale analysis due to Aizenman and Newman, we show that the long distance behavior of the connectivity τ_{xy} is governed by the probability p_{xy} . The result holds up to the critical point.

1 Introduction

In this short note we consider a long range percolation model on $\mathbb{L} = (\mathbb{Z}^{k+d}, \mathbb{B})$, where $u \in \mathbb{Z}^{k+d}$ is of the form $u = (\vec{u}_0, \vec{u}_1)$, with $\vec{u}_0 \in \mathbb{Z}^k$ and $\vec{u}_1 \in \mathbb{Z}^d$ and \mathbb{B} is the set of edges (unordered pairs) (u, v) , $u \neq v \in \mathbb{Z}^{k+d}$. To each edge (u, v) we associate a Bernoulli random variable ω_b which is open ($\omega_b = 1$) with probability

$$p_{uv} = p_{uv}(\beta) \equiv \beta J_{uv}, \quad u, v \in \mathbb{Z}^{k+d} \quad (1)$$

where $\beta \in [0, 1]$ and, for $\epsilon > 0$, J_{uv} is

$$J_{uv} = \begin{cases} 2(1 + \|\vec{u}_1 - \vec{v}_1\|^{d+\epsilon})^{-1} & \text{if } \vec{u}_0 = \vec{v}_0 \text{ and } \vec{u}_1 \neq \vec{v}_1; \\ 1 & \text{if } \vec{u}_1 = \vec{v}_1 \text{ and } \|\vec{u}_0 - \vec{v}_0\| = 1; \\ 0 & \text{otherwise.} \end{cases} \quad (2)$$

We denote the event $\{\omega \in \Omega : \text{there is an open path connecting } x \text{ to } y\}$ by $\{x \leftrightarrow y\}$ and define the connectivity function by $\tau_{xy} \equiv P\{x \leftrightarrow y\}$. Let $\|x\| = |x_1| + \dots + |x_d|$ be the L^1 norm on \mathbb{Z}^d and $\beta_c = \sup\{\beta \in [0, 1] : \chi(\beta) < \infty\}$. Our aim is to show the following result

Theorem 1 *Suppose $\beta < \beta_c$ and consider the long range percolation model with p_{uv} given by (1) and J_{uv} given by (2). Then there exist positive constants $C = C(\beta)$ and $m = m(\beta)$*

G.A. Braga (✉) · L.M. Ciolleti · R. Sanchis
Departamento de Matemática, UFMG, Caixa Postal 1621, 30161-970 Belo Horizonte, MG, Brazil
e-mail: gbraga@mat.ufmg.br

such that

$$\tau_{xy} \leq \frac{C e^{-m\|\vec{x}_0 - \vec{y}_0\|}}{1 + \|\vec{x}_1 - \vec{y}_1\|^{d+\varepsilon}} \tag{3}$$

for all $x, y \in \mathbb{Z}^{k+d}$.

The above result says that the probability p_{uv} dictates the long distance behavior of the connectivity function in the subcritical regime (similar lower bounds are easily obtained from the FKG inequality). For the one dimensional \mathbb{Z}^{0+1} percolation model, the above result is known to hold, see [1], the same being true for one dimensional \mathbb{Z}^{0+1} $O(N)$ spin models, $1 \leq N \leq 4$, see [2]. The result is expected to hold in the d -dimensional \mathbb{Z}^{0+d} lattice but it is not clear how to prove it if $\beta < \beta_c$, although one can see it holds if $\beta \approx 0$. Our upper bound (3) holds if $\beta < \beta_c$ and for $(k + d)$ -dimensional lattices, $k \geq 0$ and $d \geq 1$. For lattice spin models, the upper bound (3) is known at the high temperature regime, see Ref. [3] for bounded spin models and Ref. [4] for unbounded (and discrete) ones. Reference [5] extends some of the results of [3, 4] to a general class of continuous spin systems, with J_{uv} given by (1) and $u, v \in \mathbb{Z}^{0+d}$, while [6] considers the more general mixed decay model. In both cases, the polymer expansion (see [7] and references therein) is used.

The mixed percolation model we study is motivated by equivalent ferromagnetic models that arise as special cases of discrete space-time field theories as in [6] where the discretized Laplacian is replaced by a nearest neighbor ferromagnetic interaction. The model is interesting by itself and it has already appeared in the literature [8].

The Hammersley–Simon–Lieb inequality [9–11] is a key ingredient in [1] and [2] and here we also adopt this “correlation inequality” point of view. For completeness, we state this inequality in the form we will use, see [12]. For each set $S \subset \mathbb{Z}^d$, let $\tau_{xy}^S \equiv P\{x \leftrightarrow y \text{ inside } S\}$. Then

Hammersley–Simon–Lieb Inequality (HSL) *Given $x, y \in \mathbb{Z}^{k+d}$, if $S \subset \mathbb{Z}^{k+d}$ is such that $x \in S$ and $y \in S^c$, then*

$$\tau_{xy} \leq \sum_{\{u \in S, v \in S^c\}} \tau_{xu}^S p_{uv} \tau_{vy}$$

We now recall some known facts about the long range percolation model defined by (1) and (2). Let $\theta(\beta, \varepsilon) = P_{\beta, \varepsilon}\{0 \leftrightarrow \infty\}$ be the probability that the origin will be connected to infinity. If $k + d \geq 2$ is the space dimension then, by comparing with the nearest neighbor model and for any positive ε , there exists $\beta_c = \beta_c(d, \varepsilon)$ such that $\theta(\beta, \varepsilon) = 0$ if $\beta < \beta_c$ and $\theta(\beta, \varepsilon) > 0$ if $\beta > \beta_c$. For $k = 0$ and $d = 1$, it is known that the existence of β_c depends upon ε , if $\varepsilon > 1$ then there is no phase transition [13] while it shows up if $0 \leq \varepsilon \leq 1$, see [14]. A phase transition can also be measured in terms of χ , the mean cluster size, given by $\chi = \sum_x \tau_{0x}$. Let $\pi_c(d, \varepsilon) = \sup\{\beta : \chi(\beta, \varepsilon) < \infty\}$. Then, it comes from the FKG Inequality [15] that $\pi_c(d, \varepsilon) \leq \beta_c(d, \varepsilon)$. The equality $\pi_c(d, \varepsilon) = \beta_c(d, \varepsilon)$ holds for the class of models we are dealing with and it was proved independently by Aizenman and Barsky in [16] and Menshikov in [17]. We will use the condition $\chi < \infty$ to characterize the subcritical region.

The remaining of this note is divided as follows: in the next section we prove Theorem 1 and in Sect. 3 we make some concluding remarks regarding the validity of our results to ferromagnetic spin models.

2 Proof of the Theorem

Let $x = (\vec{x}_0, \vec{x}_1) \in \mathbb{Z}^{k+d}$. We first observe that

$$\chi = \sum_{n \geq 0} \sum_{\|\vec{x}_0\|=n} \sum_{\vec{x}_1 \in \mathbb{Z}^d} \tau_{0x}.$$

Since $\chi < \infty$, given $\lambda \in (0, 1)$, there exists $n_0 \in \mathbb{N}$ such that, for all $n \geq n_0$, we have

$$\sum_{\|\vec{x}_0\|=n} \sum_{\vec{x}_1 \in \mathbb{Z}^d} \tau_{0x} < \lambda.$$

Consider now $x = (\vec{x}_0, \vec{x}_1)$ with $\|\vec{x}_0\| > n_0$. Using the translation invariance of the model and applying iteratively the HSL Inequality with $S = \{x \in \mathbb{Z}^{k+d}; \|\vec{x}_0\| \leq n_0\}$, we obtain

$$\sum_{\vec{x}_1 \in \mathbb{Z}^d} \tau_{0x} \leq \lambda^{\lfloor \|\vec{x}_0\|/n_0 \rfloor} \leq C_1 \exp(-(m + \delta)\|\vec{x}_0\|),$$

where $\lfloor r \rfloor$ denotes the integer part of r , $\delta > 0$ is given and m is defined by $e^{-(m+\delta)} = \lambda^{1/n_0}$. Next we show that a HSL type inequality holds for the modified connectivity function $T_m(x, y) \equiv e^{m\|\vec{x}_0 - \vec{y}_0\|} \tau_{xy}$ and for the set $S = C_r(x) \equiv \{z \in \mathbb{Z}^{k+d}; \|\vec{x}_1 - \vec{z}_1\| \leq r\}$. Applying the HSL Inequality with the above specified S , we have

$$\tau_{xy} \leq \sum_{\substack{u \in C_r(x) \\ v \in C_r^c(x)}} \tau_{xu} p_{uv} \tau_{vy}.$$

Then, for $y \in C_L^c(x)$ for some $L > 1$, we obtain

$$\begin{aligned} T_m(x, y) &= e^{m\|\vec{x}_0 - \vec{y}_0\|} \tau_{xy} \leq e^{m\|\vec{x}_0 - \vec{y}_0\|} \sum_{\substack{u \in C_L(x) \\ v \in C_L^c(x)}} \tau_{xu} p_{uv} \tau_{vy} \\ &\leq \sum_{\substack{u \in C_L(x) \\ v \in C_L^c(x)}} e^{m\|\vec{x}_0 - \vec{u}_0\|} \tau_{xu} p_{uv} e^{m\|\vec{y}_0 - \vec{v}_0\|} \tau_{vy} \end{aligned}$$

since we necessarily have that $\vec{u}_0 = \vec{v}_0$. It then follows that

$$T_m(x, y) \leq \sum_{\substack{u \in C_L(x) \\ v \in C_L^c(x)}} T_m(x, u) p_{uv} T_m(v, y).$$

We remark that $\chi_m \equiv \sum_{x \in \mathbb{Z}^{k+d}} T_m(0, x) < \infty$ is finite if $\beta < \beta_c$ since

$$\begin{aligned} \sum_{x \in \mathbb{Z}^{k+d}} T_m(0, x) &= \sum_{x \in \mathbb{Z}^{k+d}} e^{m\|\vec{x}_0\|} \tau_{0x} \leq \sum_{k \geq 0} \sum_{\|\vec{x}_0\|=k} e^{m\|\vec{x}_0\|} \sum_{\vec{x}_1 \in \mathbb{Z}^d} \tau_{0x} \\ &\leq \sum_{k \geq 0} 2dk^{d-1} e^{-\delta k} < \infty. \end{aligned}$$

From now on we closely follow Sect. 3 of [1] and prove the polynomial decay of T_m up to the critical point. For fixed $x, y \in \mathbb{Z}^{k+d}$ and $L \equiv \|\vec{x}_1 - \vec{y}_1\|/4$, we know that

$$\begin{aligned}
 T_m(x, y) &\leq \sum_{\substack{u \in C_L(x) \\ v \in C_L^c(x)}} T_m(x, u) p_{uv} T_m(v, y) \\
 &\leq \sum_{\substack{u \in C_L(x) \\ v \in C_L^c(x) \cap C_{3L}(x)}} T_m(u, x) p_{uv} T_m(v, y) + \sum_{\substack{u \in C_L(x) \\ v \in C_L^c(x) \cap C_{3L}^c(x)}} T_m(x, u) p_{uv} T_m(v, y). \tag{4}
 \end{aligned}$$

Let

$$\mathbb{T}_m(L) \equiv \sup\{T_m(0, u); u \in C_L^c(0)\} \quad \text{and} \quad \gamma_L \equiv \sum_{\substack{u \in C_L(x) \\ v \in C_L^c(x)}} T_m(x, u) p_{uv}.$$

Then, the first term on the r.h.s. of (4) is bounded above by $\mathbb{T}_m(L/2)\gamma_L$ while the second one is bounded by

$$\frac{2^{d+\varepsilon} 2\beta \chi_m^2}{1 + \|\vec{x}_1 - \vec{y}_1\|^{d+\varepsilon}},$$

leading to

$$T_m(x, y) \leq \frac{2^{d+\varepsilon} 2\beta \chi_m^2}{1 + \|\vec{x}_1 - \vec{y}_1\|^{d+\varepsilon}} + \gamma_L \mathbb{T}_m\left(\frac{L}{2}\right).$$

Now, since $\chi_m < \infty$ for $\beta < \beta_c$ and since $\sum_u p_{0u} < \infty$, we have that $\gamma_L \rightarrow 0$ as $L \rightarrow \infty$. For $\alpha \in (0, 2^{-(d+\varepsilon)})$, there exists $L_0 > 0$ such that $\gamma_L < \alpha$ for all $L \geq L_0$. Considering $L > L_0$, it follows that

$$\mathbb{T}_m(L) \leq \frac{2^{d+\varepsilon} 2\beta \chi_m^2}{1 + \|\vec{x}_1 - \vec{y}_1\|^{d+\varepsilon}} + \alpha \mathbb{T}_m\left(\frac{L}{2}\right). \tag{5}$$

Iterating (5) n times, with n the smallest integer for which $L2^{-n} \leq L_0$, we have for all $L > L_0$

$$\mathbb{T}_m(L) \leq \frac{2\beta \chi_m^2 \sum_{j=0}^{n-1} (\alpha 2^{d+\varepsilon})^j}{1 + \|\vec{x}_1 - \vec{y}_1\|^{d+\varepsilon}} + \alpha^n \mathbb{T}_m\left(\frac{L}{2^n}\right).$$

Noting that $T_m(x, y) \leq \mathbb{T}_m(L)$, that $\mathbb{T}_m(L) \leq 1$ for any $L > 0$ and that

$$\begin{aligned}
 \alpha^n &\leq 2^{-(d+\varepsilon)n} = \frac{1}{(1 + L^{d+\varepsilon})} \frac{(1 + L^{d+\varepsilon})}{2^{(d+\varepsilon)n}} \leq \frac{2 \cdot 2^{d+\varepsilon}}{1 + \|\vec{x}_1 - \vec{y}_1\|^{d+\varepsilon}} \left(\frac{L}{2^n}\right)^{d+\varepsilon} \\
 &\leq \frac{2(2L_0)^{d+\varepsilon}}{1 + \|\vec{x}_1 - \vec{y}_1\|^{d+\varepsilon}},
 \end{aligned}$$

we can conclude that, for $\beta < \beta_c$,

$$\tau_{xy} \leq \frac{C e^{-m\|\vec{x}_0 - \vec{y}_0\|}}{1 + \|\vec{x}_1 - \vec{y}_1\|^{d+\varepsilon}}$$

and the bound (3) holds.

3 Concluding Remarks

The strategy used to prove Theorem 1 can also be applied to \mathbb{Z}^d ferromagnetic models with free boundary conditions and pair interaction J_{uv} given by (2). For this class of models the Griffiths inequalities [18, 19] are valid, guaranteeing the positivity of spin-spin correlations $\langle \sigma_x \sigma_y \rangle$. Simon–Lieb Inequality also holds in this case (see [7] and references therein), with p_{uv} replaced by βJ_{uv} , where β is the inverse of the temperature, and with τ_{xy} replaced by $\langle \sigma_x \sigma_y \rangle$. Finally, since the uniqueness of the critical point is guaranteed in [20], the results of Sect. 2 are also valid for these models.

Acknowledgements G.B. was partially supported by CNPq; L.C. acknowledges CNPq for a graduate scholarship; R.S. was partially supported by Pró-Reitoria de Pesquisa—UFMG under grant 10023.

References

1. Aizenman, M., Newman, C.M.: Discontinuity of the percolation density in one-dimensional $1/|x - y|^2$ percolation models. *Commun. Math. Phys.* **107**(4), 611–647 (1986)
2. Spohn, H., Zwirger, W.: Decay of the two-point function in one-dimensional $O(N)$ spin models with long-range interactions. *J. Stat. Phys.* **94**, 1037–1043 (1999)
3. Gross, L.: Decay of correlations in classical lattice models at high temperature. *Commun. Math. Phys.* **68**, 9–27 (1979)
4. Israel, R., Nappi, C.: Exponential clustering for long-range integer-spin systems. *Commun. Math. Phys.* **68**, 29–37 (1979)
5. Procacci, A., Scoppola, B.: On decay of correlations for unbounded spin systems with arbitrary boundary conditions. *J. Stat. Phys.* **105**, 453–482 (2001)
6. Thebaldi, R.S., Pereira, E., Procacci, A.: A cluster expansion for the decay of correlations of light-mass quantum crystals and some stochastic models under intense noise. *J. Math. Phys.* **46**, 053303 (2005)
7. Glimm, J., Jaffe, A.: *Quantum Physics: A Functional Integral Point of View*, 2nd edn. Springer, New York (1987)
8. Sidoravicius, V., Surgailis, D., Vares, M.E.: On the truncated anisotropic long-range percolation on \mathbb{Z}^2 . *Stoch. Process. Appl.* **88**, 337–349 (1999)
9. Hammersley, J.M.: Percolation processes. Lower bounds for the critical probability. *Ann. Math. Stat.* **28**, 790–795 (1957)
10. Simon, B.: Correlation inequalities and the decay of correlations in ferromagnets. *Commun. Math. Phys.* **77**, 111–126 (1980)
11. Lieb, E.H.: A refinement of Simon’s correlation inequality. *Commun. Math. Phys.* **77**, 127–135 (1980)
12. Aizenman, M., Newman, C.M.: Tree graph inequalities and critical behaviour in percolation models. *J. Stat. Phys.* **36**, 107–143 (1984)
13. Schulman, L.S.: Long range percolation in one dimension. *J. Phys. Lett. A* **16**, L639 (1983)
14. Newman, C.M., Schulman, L.S.: One-dimensional $1/|j - i|^s$ percolation models: the existence of a transition for $s \leq 2$. *Commun. Math. Phys.* **104**, 547–571 (1986)
15. Grimmett, G.: *Percolation*, 2nd edn. Springer, Berlin (1999)
16. Aizenman, M., Barsky, D.J.: Sharpness of the phase transition in percolation models. *Commun. Math. Phys.* **108**(3), 489–526 (1987)
17. Menshikov, M.V.: Coincidence of critical points in percolation problems. *Sov. Math. Dokl.* **33**, 856–859 (1986)
18. Griffiths, R.B.: Correlations in Ising ferromagnets, I. *J. Math. Phys.* **8**, 478–483 (1967)
19. Griffiths, R.B.: Correlations in Ising ferromagnets, II: external magnetic fields. *J. Math. Phys.* **8**, 484–489 (1967)
20. Aizenman, M., Barsky, D.J., Fernández, R.: The phase transition in a general class of Ising-type models is sharp. *J. Stat. Phys.* **47**(3–4), 343–374 (1987)