

SLE on Doubly-Connected Domains and the Winding of Loop-Erased Random Walks

Christian Hagendorf · Pierre Le Doussal

Received: 25 March 2008 / Accepted: 13 August 2008 / Published online: 10 September 2008
© Springer Science+Business Media, LLC 2008

Abstract Two-dimensional loop-erased random walks (LERWs) are random planar curves whose scaling limit is known to be a Schramm-Loewner evolution SLE_κ with parameter $\kappa = 2$. In this note, some properties of an SLE_κ trace on doubly-connected domains are studied and a connection to passive scalar diffusion in a Burgers flow is emphasised. In particular, the endpoint probability distribution and winding probabilities for SLE_2 on a cylinder, starting from one boundary component and stopped when hitting the other, are found. A relation of the result to conditioned one-dimensional Brownian motion is pointed out. Moreover, this result permits to study the statistics of the winding number for SLE_2 with fixed endpoints. A solution for the endpoint distribution of SLE_4 on the cylinder is obtained and a relation to reflected Brownian motion pointed out.

Keywords Loop-erased random walks · Stochastic Loewner evolution · Stochastic processes · Burgers equation

1 Introduction

The Schramm-Loewner Evolution SLE_κ is a one-parameter family of conformally invariant measures on non-self-crossing planar curves. In many cases, interfaces in two-dimensional models of statistical mechanics at criticality are conjectured to be described by SLE. For example, the scaling limit of the planar loop-erased random walk (LERW) is known to be SLE_2 , interfaces in the 2D Ising model SLE_3 and critical percolation hulls to be described by SLE_6 . While these systems can be also studied using traditional methods of theoretical physics like (boundary) conformal field theory and Coulomb gas methods, Schramm has given a novel and rigorous approach via conformally invariant stochastic growth processes.

LERWs were the starting point and motivation for the development of SLE in Schramm's seminal paper [30]. Today their scaling limits and relations to SLE are well understood in

C. Hagendorf (✉) · P. Le Doussal
CNRS-Laboratoire de Physique Théorique de l'Ecole Normale Supérieure, 24, rue Lhomond, 75231
Paris cedex 05, France
e-mail: hagendorf@lpt.ens.fr

simply-connected domains, where conformal invariance was shown by Lawler, Schramm and Werner [27]. Properties in multiply-connected domains are more involved but much recent progress was obtained, e.g. in a series of works by Dapeng Zhan [32–34]. In this paper, we mainly study properties of planar LERWs/SLE₂ in doubly-connected domains, emerging from one boundary component and conditioned to finish when hitting the second boundary component. In contrast to the simply-connected case, curves in doubly-connected domains may wind non-trivially around various boundary components. Here we concentrate on these winding properties of loop-erased random walks on cylinders in two cases: (1) with one endpoint fixed on one boundary component, and the other endpoint free on the other boundary component, (2) with both endpoints fixed on different boundary components. We establish these results by solving a differential equation arising from SLE_κ with $\kappa = 2$ on the cylinder. Another motivation to study the cylinder is that numerical simulations of critical systems are often performed in a cylinder geometry, see e.g. [12].

The paper is organised as follows. In Sect. 2 we give a brief description of SLE. In particular, we outline two variants of SLE in simply-connected domains which prove to be limits (in some sense) of the doubly-connected case. After this we discuss SLE on cylinders, being prototypes of doubly-connected domains. Section 3 provides a detailed discussion on endpoint probabilities and winding properties of SLE_κ on cylinders. Inspired from analogies with diffusion-advection in $(1 + 1)$ -dimensional Burgers flows, we present analytical results for the case $\kappa = 2$, study various limits and point out a relation between the winding of LERWs and conditioned one-dimensional Brownian motion. Finally, conditioning walks to exit via a given point, we derive the winding statistics for LERW with fixed endpoints on the cylinder. In Sect. 4, somewhat aside the main topic, we provide the endpoint distribution in the case $\kappa = 4$ and point out a relation to reflected two-dimensional Brownian motion. We conclude in Sect. 5. The appendix contains computations of exit probabilities for Brownian motion and random walks on the lattice, in relation to distributions of exit points for SLE₂ and their finite-size corrections. It also contains a partial (periodised) result about bulk left-passage probabilities with respect to a point in the covering space of the cylinder, as well as a path-integral version of the endpoint distribution problem.

2 Basic Notions of SLE

In this Section we give an informal description of SLE. For detailed accounts we refer the reader to the many existing excellent review articles on this subject (see e.g. [6, 13, 26]).

Stochastic Loewner evolutions (SLE) describe the growth of non-intersecting random planar curves or hulls emerging from the boundary of planar domains \mathbb{D} by means of stochastic differential equations. A crucial feature of SLE is *conformal invariance*: if two domains \mathbb{D} and \mathbb{D}' are related by a conformal mapping $g(z)$, then the image of SLE in \mathbb{D} is SLE in \mathbb{D}' . Another key property is the so-called *domain Markov property* of SLE. Conditioning on the curve/hull γ_t grown up to time t in \mathbb{D} , the remainder of the curve is just SLE in the slit domain $\mathbb{D}_t = \mathbb{D} \setminus \gamma_t$. In fact, these two properties (almost) imply that the growing hulls are characterised by a single positive parameter κ .

For simply-connected domains, conformal invariance and the Riemann mapping theorem naturally lead to a description of the curves in terms of conformal mappings $g_t(z)$ which uniformise \mathbb{D}_t to \mathbb{D} , and map the tip of γ_t to some point on the boundary of \mathbb{D} . $g_t(z)$ is solution of a differential equation in t with initial condition $g_{t=0}(z) = z$, called the *Loewner equation*. SLE in simply-connected domains comes in different flavours, according to which the Loewner differential equation varies. In Sect. 2.1, we shall concentrate on so-called radial and dipolar SLEs.

Compared to simply-connected domains, SLE in multiply-connected domains proves to be more involved because no equivalent of the Riemann mapping theorem is at hand [5, 10, 32, 33]. In this article, we concentrate on doubly-connected domains \mathbb{D} with the topology of an annulus. Their boundary is given by two disjoint simple curves $\partial_1\mathbb{D}$ and $\partial_2\mathbb{D}$. A classical theorem of complex analysis states that each such a domain may be mapped onto the annulus $\mathbb{A}_p = \{z \in \mathbb{C} \mid e^{-p} < |z| < 1\}$ with some $p > 0$, such that for example $\partial_1\mathbb{D}$ is mapped to $|z| = 1$ and $\partial_2\mathbb{D}$ to $|z| = e^{-p}$. Conformal invariance of SLE legitimates the choice of \mathbb{A}_p as a reference domain. However, we shall rather work with a cylinder $\mathbb{T}_p = \{z = x + iy \in \mathbb{C} \mid -\pi < x \leq \pi, 0 < y < p\}$, identifying $x = -\pi$ and $x = \pi$. The cylinder \mathbb{T}_p may always be mapped back onto \mathbb{A}_p via the conformal mapping $w = e^{iz}$. The height of the cylinder $p > 0$ is called *the modulus*, and the boundary components are given by $\partial_1\mathbb{T}_p = [-\pi, \pi]$ and $\partial_2\mathbb{T}_p = [-\pi, \pi] + ip$ which we shall refer to as “lower” and “upper boundary” respectively. We shall study hulls γ_t growing from $z = 0 \in \partial_1\mathbb{T}_p$ to $\partial_2\mathbb{T}_p$ as shown on Fig. 1a and b. In Sect. 2.2, we analyse the Loewner equation for $g_t(z)$ mapping $\mathbb{T}_p \setminus \gamma_t$ to some cylinder \mathbb{T}_q with (in general) different modulus $q \neq p$. A mapping back to \mathbb{T}_p does not exist since one may show that there is no conformal mapping sending a cylinder of modulus p to a cylinder of modulus $q \neq p$ [29]. Moreover, we show how the doubly-connected version of SLE on \mathbb{T}_p interpolates between radial and dipolar SLE.

2.1 SLE in Simply-Connected Domains

Before embarking into considerations on general values for the modulus p we shall consider limiting cases of semi-infinite cylinders $p \rightarrow +\infty$ and very thin cylinders $p \rightarrow 0^+$ in order to establish a connection with well-known variants of SLE in simply-connected domains.

$p \rightarrow +\infty$ and radial SLE. Radial SLE starts from a boundary point $x_0 \in \partial\mathbb{D}$ to a bulk point $x_\infty \in \mathbb{D}$ of some simply-connected planar domain \mathbb{D} . Its Loewner equation is most conveniently written for the unit disc $\mathbb{D} = \mathbb{U}$ from $x_0 = 1$ to $x_\infty = 0$ where

$$\frac{dg_t(z)}{dt} = -g_t(z) \frac{g_t(z) + e^{i\xi_t}}{g_t(z) - e^{i\xi_t}}, \tag{1}$$

where ξ_t is some real-valued function of $t \geq 0$ and $g_t(0) = 0$ (the point x_∞ remains unchanged during time evolution). The time parametrisation has been chosen so that $g'_t(0) = e^t$. The real-valued function ξ_t is related to the image of the tip τ_t of the growing trace γ_t by $g_t(\tau_t) = e^{i\xi_t}$. In fact, conformal invariance and the domain Markov property imply that $\xi_t = \sqrt{\kappa}B_t + \alpha t$ is simple Brownian motion with a diffusion constant κ and a linear drift αt . If we ask for reflection symmetry (or left-right-symmetry) as it occurs in physical systems (and throughout this paper), the drift vanishes $\alpha = 0$. For later considerations, we outline another useful version of radial SLE. In fact, the unit disc \mathbb{U} may be sent to the semi-infinite cylinder \mathbb{T}_∞ via the conformal mapping $w = -i \log z$ (here the principal branch of the logarithm has been chosen). In this geometry, the Loewner equation for the new uniformising map $\tilde{g}_t(z)$ is given by

$$\frac{d\tilde{g}_t(z)}{dt} = \cot\left(\frac{\tilde{g}_t(z) - \xi_t}{2}\right), \quad \xi_t = \sqrt{\kappa}B_t. \tag{2}$$

obtained from (1) using $g_t(z) = \exp[i\tilde{g}_t(w(z))]$.

Since we aim at studying winding probabilities, let us recall how to give an intuitive (yet non-rigorous) argument in order to estimate the distribution of the winding angle θ_t

for large t [13]. For the unit disc \mathbb{U} as $t \gg 1$, the tip of the radial SLE trace τ_t approaches $z = 0$. We may therefore write $g_t(\tau_t) = e^{i\xi_t} \approx e^t \tau_t$, by using the Taylor series expansion $g_t(z) = e^t z + \dots$ near $z = 0$, from what follows $\tau_t \approx e^{-t+i\xi_t}$. Hence the winding angle $\theta_t = \arg \tau_t \approx \xi_t$ turns out to be a Gaussian random variable with zero mean and variance κt . For the infinite cylinder \mathbb{T}_∞ , the winding angle corresponds to the real part of the tip allowed to vary continuously on \mathbb{R} , and its probability distribution is approximately given by

$$\lambda_r(t, x) \approx \frac{1}{\sqrt{2\pi\kappa t}} \exp\left(-\frac{x^2}{2\kappa t}\right), \quad \text{for large } t. \tag{3}$$

The intuitive argument should be taken with caution since the power series expansion of $g_t(z)$ has a radius of convergence less than $|\tau_t|$. However, Schramm has shown by rigorous methods that indeed $\theta_t/\sqrt{\kappa t}$ converges to a Gaussian with unit variance in the limit $t \rightarrow \infty$ [30] as suggested by (3). Within later considerations on doubly-connected domains we bypass these difficulties by providing exact solutions (at least for $\kappa = 2$) of the winding angle problem, without making reference to approximate series expansions.

p $\rightarrow 0^+$ and dipolar SLE. Another version of SLE is obtained by considering planar curves in a simply-connected domain \mathbb{D} from a boundary point $x_0 \in \partial\mathbb{D}$ to a side arc $S \subset \partial\mathbb{D}$ [7] not containing x_0 . Let us choose the geometry of an infinite strip $\mathbb{S}_p = \{z \in \mathbb{C} \mid 0 < \text{Im} z < p\}$ of height p , $x_0 = 0$ and $S = \mathbb{R} + ip$. Then Loewner’s equation is given by

$$\frac{dg_t(z)}{dt} = \frac{\pi}{p} \coth\left[\frac{\pi(g_t(z) - \xi_t)}{2p}\right]. \tag{4}$$

$g_t(z)$ maps the tip of the trace γ_t to ξ_t . Again, because of conformal invariance, the domain Markov property and reflection symmetry we have $\xi_t = \sqrt{\kappa} B_t$. The trace γ_t hits the boundary arc $\mathbb{R} + ip$ at some (random) point $x + ip$ as $t \rightarrow +\infty$. This version of SLE is related to the cylinder problem on \mathbb{T}_p in the limit $p \rightarrow 0^+$. Loosely spoken, if p is very small the SLE trace will not feel the periodicity of the cylinder but only hit the upper boundary close to the point ip on the upper boundary (this statement can be made more precise, see below). The equivalent to the winding angle distribution for radial SLE is the probability distribution of this exit point for dipolar SLE, which we shall denote $\lambda_d(p, x)$. For any κ it is given by [7]

$$\lambda_d(p, x) = \frac{\sqrt{\pi} \Gamma(1/2 + 2/\kappa)}{2p \Gamma(2/\kappa)} \frac{1}{[\cosh(\pi x/2p)]^{4/\kappa}}. \tag{5}$$

2.2 Stochastic Loewner Evolution on Doubly-Connected Domains

Let us now turn to the case of general modulus p . The Loewner equation for the uniformising map $g_t(z)$ of SLE_κ on \mathbb{T}_p is given by [33]

$$\frac{dg_t(z)}{dt} = H(p - t, g_t(z) - \sqrt{\kappa} B_t) \tag{6}$$

with standard Brownian motion B_t and the function

$$H(p, z) = \lim_{n \rightarrow \infty} \sum_{k=-n}^n \cot\left(\frac{z}{2} - ikp\right) = \cot \frac{z}{2} + \sum_{k=1}^\infty \frac{2 \sin z}{\cosh 2kp - \cos z}. \tag{7}$$

The process is defined between $t = 0$ ($g_0(z) = z$) where the trace starts from point $z = 0$ on the lower boundary up to $t = p$ where the SLE trace hits the upper boundary at some random point γ_p with $\text{Im } \gamma_p = p$. The time parametrisation of the curves is chosen in such a way that $g_t(z)$ maps $\mathbb{T}_p \setminus \gamma_t$ to \mathbb{T}_q with $q = p - t$. One possible way to derive (6) and (7) is by considering an infinitesimal hull, and using invariance properties of \mathbb{T}_p , namely translation invariance $z \mapsto z + a$, $a \in \mathbb{R}$ and reflection invariance $z \mapsto ip - z$ (see [5] where this has been explicitly carried out for an annulus geometry).

Let us describe the properties of the function defined in (7). $H(p, z)$ is an odd function in z with period 2π . Furthermore it is quasi-periodic in the imaginary direction: $H(p, z + 2ip) = H(p, z) - 2i$. Moreover, notice the interesting property $H(p, z) = (i\pi H(\pi^2/p, i\pi z/p) - z)/p$. It may be used in order to derive an alternative series expansion

$$H(p, z) = \frac{\pi}{p} \lim_{n \rightarrow \infty} \sum_{k=-n}^n \coth \left(\frac{\pi(z + 2\pi k)}{2p} \right) - \frac{z}{p}. \tag{8}$$

For large positive p , we have $H(p - t, z) = \cot(z/2) + \dots$ as it can be seen from (7). Therefore, as we let $p \rightarrow +\infty$, the Loewner equation (6) reduces to the one for radial SLE, defined in (2). The dipolar limit $p \rightarrow 0^+$ is more involved. Consider the function $h_t(z) = g_t(z) - \sqrt{\kappa} B_t$ mapping $\mathbb{T}_p \setminus \gamma_t$ to \mathbb{T}_{p-t} and the tip τ_t of the curve to 0. Rescaling $k_s(z) = \pi h_t(z)/(p - t)$ maps the problem to a cylinder of fixed height π , and width $2\pi^2/(p - t)$. The new mapping $k_s(z)$ depends on a time $s = \pi^2/(p - t)$, $0 \leq t < p$. Performing the time change leads to a stochastic differential equation $dk_s(z) = iH(s, i\pi k_s(z))ds - \sqrt{\kappa} d\tilde{B}_s$ where \tilde{B}_s is another standard Brownian motion, obtained from the time-change formula. Now, as $p \rightarrow 0^+$ we have $s \rightarrow +\infty$ so that we may use $H(s, z) \sim \cot z/2 + \dots$. Finally for $K_s(z) = k_s(z) + \sqrt{\kappa} \tilde{B}_s$, we find the dipolar Loewner equation

$$\frac{dK_s(z)}{ds} = \coth \left(\frac{K_s(z) - \sqrt{\kappa} \tilde{B}_s}{2} \right) \tag{9}$$

in the limit $s \rightarrow +\infty$.

The covering space. In the sequel it shall be convenient to lift the processes to the covering space of the cylinder \mathbb{T}_p . In fact, this is simply obtained by allowing $\text{Re } \gamma_t$ to vary continuously on \mathbb{R} . An illustration is given on Fig. 1b. It is easy to see that the difference between the endpoints of a given trace on the cylinder and its analogue on the covering space is always given by $2\pi N$ with an integer N . We shall call N the winding number of the SLE trace. For the covering space, the Loewner equation (6) still holds. However, probabilities and probability densities computed on the covering space have different boundary conditions when compared to the periodic case of cylinders [33].

3 Endpoint Probabilities for SLE on the Cylinder

In this section, we characterise the endpoint distributions for SLE on cylinders of height p which we denote $\Lambda(p, x)$, $-\pi \leq x \leq \pi$. To be precise, $\Lambda(p, x)dx$ corresponds to the probability that the trace hits the upper boundary within $\text{Re } \gamma_p \in [x, x + dx]$ for some infinitesimal dx . In order to study the winding of the traces around the cylinder, we introduce an




Fig. 1 (a) Illustration for SLE on an annulus \mathbb{A}_p from l to the target circle $|z| = e^{-p}$. (b) SLE on the cylinder and the covering space: the shaded region represents the cylinder, with sides $x = -\pi$ and $x = \pi$ identified. We obtain the covering space by periodic extension along the real axis. The endpoints on the cylinder and the covering space differ by $2\pi N$ where N denotes the winding number of the trace. The interfaces are samples of loop-erased random walks

analogous object $\lambda(p, x)$, $x \in \mathbb{R}$ for the covering space. Results on the cylinder are obtained upon identification of points differing by $2\pi n$:

$$\Lambda(p, x) = \sum_{n=-\infty}^{\infty} \lambda(p, x + 2\pi n). \tag{10}$$

3.1 Endpoint Distributions for General κ

In order to compute $\Lambda(p, x)$ consider the SLE trace γ_{dt} evolving for an infinitesimal time dt on the cylinder \mathbb{T}_p . The uniformising map $g_{dt}(z)$ maps the remainder of the curve $\gamma_p \setminus \gamma_{dt}$ onto SLE in a cylinder \mathbb{T}_{p-dt} starting at $\sqrt{\kappa} dB_0$. Under this mapping $g_{dt}(z)$ the point $x + ip$ on the upper boundary of \mathbb{T}_p is sent to some point $x' + i(p - dt)$ on the upper boundary of \mathbb{T}_{p-dt} , with $x' = x + (H(p, x + ip) + i)dt$, see (6). Using the Markov property and conformal invariance of the SLE measure we obtain, upon averaging over the infinitesimal Brownian increment dB_0 , the equation

$$\Lambda(p, x)dx = \mathbb{E}_{dB_0}[\Lambda(p - dt, x' - \sqrt{\kappa} dB_0)dx']. \tag{11}$$

An expansion to first order in dt , using $\mathbb{E}_{dB_0}[dB_0] = 0$ and $\mathbb{E}_{dB_0}[dB_0^2] = dt$, leads to the Fokker-Planck equation

$$\frac{\partial \Lambda(p, x)}{\partial p} = \frac{\partial}{\partial x} (v(p, x)\Lambda(p, x)) + \frac{\kappa}{2} \frac{\partial^2 \Lambda(p, x)}{\partial x^2}, \tag{12}$$

where we have introduced the drift function (compare with (7))

$$v(p, x) = H(p, x + ip) + i = \frac{\pi}{p} \lim_{N \rightarrow \infty} \sum_{n=-N}^N \tanh \frac{\pi(x + 2\pi n)}{2p} - \frac{x}{p}. \tag{13}$$

Like $H(p, z)$ the drift function $v(p, z)$ is an odd and periodic function with period 2π . Furthermore, it is quasiperiodic in imaginary direction: $v(p, z + 2ip) = v(p, z) - 2i$. Notice that $\lambda(p, x)$ is a solution of (12) as well, however with vanishing boundary conditions

for $x \rightarrow \pm\infty$. Leaving aside the identification of the points x and $x + 2\pi$, we ask for the probability

$$\omega(p, x) = \int_{-\infty}^x dy \lambda(p, y) \tag{14}$$

that the SLE traces hits the upper boundary to the left of a given point $x + ip$ on the covering space. It is solution of the convection-diffusion equation

$$\frac{\partial \omega(p, x)}{\partial p} = v(p, x) \frac{\partial \omega(p, x)}{\partial x} + \frac{\kappa}{2} \frac{\partial^2 \omega(p, x)}{\partial x^2}, \tag{15}$$

with boundary conditions $\lim_{p \rightarrow 0^+} \omega(p, x) = \Theta(x)$, $\lim_{x \rightarrow +\infty} \omega(p, x) = 1$ and $\lim_{x \rightarrow -\infty} \omega(p, x) = 0$. Here $\Theta(x)$ denotes the Heaviside function which is 1 for $x > 0$, and 0 otherwise. It is a remarkable property of the function $v(p, x)$ to be a solution of the $(1 + 1)$ -dimensional Burgers equation

$$\frac{\partial v(p, x)}{\partial p} = v(p, x) \frac{\partial v(p, x)}{\partial x} + \frac{\partial^2 v(p, x)}{\partial x^2}, \tag{16}$$

where p plays the role of time and viscosity is set to unity [33]. In fact, this equation even holds for complex x . Furthermore it can be shown that $H(p, z)$ obeys the same equation by using the relationship between $H(p, z)$ and $v(p, z)$. Hence we interpret (15) as the evolution equation of a passive scalar convected and diffusing in the Burgers flow $-v(p, x)$, also described by a Langevin equation:

$$dx_p = -v(p, x_p)dp + \sqrt{\kappa}dB_p. \tag{17}$$

The problem of finding SLE probabilities for arbitrary κ can thus be reformulated as finding the density of a passive scalar convected by a (particular) one-dimensional Burgers flow (along x). In its general formulation it is a classic problem [4, 11, 17, 19, 20, 28] characterised by the dimensionless Prandtl number P [21], with $P = 2/\kappa$ for our problem.

By means of the well-known Cole-Hopf transformation [15, 23] we rewrite

$$v(p, x) = 2 \frac{\partial}{\partial x} \ln \epsilon(p, x), \tag{18}$$

where $\epsilon(p, x)$ satisfies the diffusion equation $\partial \epsilon(p, x) / \partial p = \partial^2 \epsilon(p, x) / \partial x^2$ with solution

$$\epsilon(p, x) = \sum_{n=-\infty}^{+\infty} (-1)^n \exp(inx - n^2 p) = \sqrt{\frac{\pi}{p}} \sum_{n=-\infty}^{+\infty} \exp\left(-\frac{(x - \pi - 2\pi n)^2}{4p}\right) \tag{19}$$

corresponding to an initial condition $\epsilon(0, x) = 2\pi \sum_{n=-\infty}^{\infty} \delta(x - \pi - 2\pi n)$. In the language of Burgers equation this corresponds to an initial condition at $p = 0^+$ with a periodic set of shocks at positions $x_n^s = 2\pi n$. As the ‘‘time’’ p increases the shocks are broadened by the viscosity and acquire a width proportional to p , leading to the smooth periodic function (13). Finding the solution of (12) for general κ is a complicated time-dependent problem, and only results in special cases are obtained below (see Appendix 4 for a path-integral formulation).

In the limit of large modulus $p \gg 1$ the drift term in (12) vanishes like $v(p, x) \sim 4e^{-p} \sin x$. Therefore we recover a simple diffusion equation for the endpoint distribution $\lambda(p, x)$ which, in terms of the scaling variable x/\sqrt{p} , converges to the Gaussian result

for radial SLE (3). However if one does not rescale x , there are non-trivial corrections and one expects instead that the cumulants converge at large p to κ dependent constants, $\langle x^{2n} \rangle_c - \kappa p \delta_{n,2} \rightarrow X_{2n,\kappa}$, as shown explicitly below for $\kappa = 2$. Going from \mathbb{T}_p to an annulus with radii $R_1 > R_2 = R_1 e^{-p}$ via the conformal mapping $w = R_1 e^{iz}$ (see Fig. 1a for an illustration) we obtain the winding angle distribution $\theta = x$ for SLE starting from the outer boundary and stopped when first hitting the inner boundary (or the reverse if we admit reversibility of annulus SLE). We expect its cumulants of the winding angle to have the form

$$\langle \theta^{2n} \rangle_c = \kappa \ln(R_1/R_2) \delta_{n,2} + X_{2n,\kappa} \tag{20}$$

in the limit of large R_1/R_2 . This is more precise than the standard result obtained from Coulomb gas methods [13, 18], or the argument given for radial SLE, namely $\langle \theta^2 \rangle = (4/g) \ln(R_1/R_2)$ with $4/g = \kappa$ (see (3)), since the universal constants $X_{2n,\kappa}$ take explicitly into account the presence of the two boundaries. One can compare to the corresponding cumulative distribution of the winding angle for the standard two-dimensional Brownian motion with absorbing conditions on the boundary of the annulus (it can be obtained from dipolar SLE with $\kappa = 2$, the direct derivation being recalled in Appendix A.1.1)

$$\omega_b(\theta) = \frac{1}{1 + e^{-\pi\theta/\ln(R_1/R_2)}} \tag{21}$$

which yields much larger cumulants $\langle \theta^{2n} \rangle_c \sim (\ln(R_1/R_2))^{2n}$ with prefactors given below in the context of dipolar SLE.

As $p \rightarrow 0^+$ a simple argument allows to recover the dipolar SLE hitting probability (5) from the form of Burgers shocks, which are well separated in that limit. Let us suppose that we may restrict (13) to a single shock, $v_s(p, x) = -x/p + \pi/p \tanh(\pi x/2p)$, i.e. take an initial condition $\epsilon(0, x) = 2\pi(\delta(x + \pi) + \delta(x - \pi))$. This gives a potential $u(p, x) = 2 \log \epsilon(p, x) = -x^2/2p + 2 \log \cosh(\pi x/2p) + \text{const.}$ which in the scaling region $x \sim p$ strongly confines the diffusing particle described by (12) near the origin. One easily checks that the (adiabatic) Gibbs measure $\lambda_{\text{adiab}}(x, p) \sim e^{-(2/\kappa)u(p,x)} = C(p)\epsilon(x, p)^{-4/\kappa} = \lambda_d(x, p)$ satisfies (12) in the region $x \sim p$ (the term $\partial \lambda(p, x)/\partial p$ on the l.h.s. being subdominant). One can be more precise and rescale $\xi = x/p$. The new probability $\tilde{\omega}(p, \xi) = \omega(p, x)$ is solution of the differential equation

$$p^2 \frac{\partial \tilde{\omega}(p, \xi)}{\partial p} = \tilde{v}(p, \xi) \frac{\partial \tilde{\omega}(p, \xi)}{\partial \xi} + \frac{\kappa}{2} \frac{\partial^2 \tilde{\omega}(p, \xi)}{\partial \xi^2}, \tag{22}$$

where $\tilde{v}(p, \xi) = p(v(p, x) + \xi)$. Inserting the single shock profile one finds $\tilde{v}_s(p, \xi) = \pi \tanh(\pi \xi/2)$, i.e. a stationary velocity field. Hence we may convince ourselves that the stationary solution of (22) is $\partial \omega_{\text{stat}}(p, x)/\partial x = \lambda_d(p, x)$, i.e. the convergence to the dipolar result (5) includes also (non exponential) subdominant terms in the scaling region $x \sim p$. Interesting exact solutions of (22) for all p are easily obtained from a Laplace transform in $\tau = 1/p$ or using methods as in [21], but do not obviously relate to SLE.

3.2 Results for $\kappa = 2$ and the Winding of Loop-Erased Random Walks

The Cole-Hopf transformation (18) will lead to interesting simplifications allowing to treat explicitly $\kappa = 2$. This case corresponds to the scaling limit of the LERW on the cylinder. First we recall its definition on the lattice [26]. Consider a rectangular lattice domain of lattice mesh δ embedded into the cylinder domain \mathbb{T}_p . Start a simple random walk \mathcal{W} from




Fig. 2 Illustration of loop-erasure on the cylinder for a square lattice. **(a)** A Brownian excursion from $z = 0$ to the upper boundary $y = p$. **(b)** Loop-erasure of the same walk. Loops like $(14, 15, \dots, 27)$, winding around the cylinder for several times, may be erased

0 to $\text{Re } z = y = p$ with absorbing boundary conditions for $y = 0$ (see Fig. 2a). Chronological loop erasure from W yields a simple path γ on the lattice domain which is the LERW (see Fig. 2b). Notice that the endpoints of W and γ always coincide. However, the random walk may wind several times around the cylinder whereas its loop erasure does not. This makes it a non-trivial exercise to compute the winding properties of the latter (i.e. its endpoint distribution on the covering space). As the lattice mesh δ goes to 0 , the paths W converge to two-dimensional Brownian excursions on \mathbb{T}_p whereas the paths γ converge to SLE_2 on \mathbb{T}_p [34].

Since the endpoints of W and γ coincide on the cylinder \mathbb{T}_p , we may find the endpoint distribution for SLE_2 by a simple computation on Brownian excursion on the cylinder, starting from $z = 0$ and stopped as soon as it hits altitude $y = p$ (see Appendix A.1.1). The result is

$$\Lambda_2(p, x) = \frac{\pi}{4p} \sum_{n=-\infty}^{\infty} \frac{1}{(\cosh(\frac{\pi(x+2\pi n)}{2p}))^2}. \tag{23}$$

An equivalent result for random walks on a finite cylindrical lattice domain is presented in Appendix 2. Indeed, $\Lambda_2(p, x)$ solves (12) for $\kappa = 2$ with periodic boundary conditions and may be related to the drift function $v(p, x)$ via $\Lambda_2(p, x) = (p \partial v(p, x) / \partial x + 1) / 2\pi$ [33]. Moreover, it corresponds to the periodised endpoint distribution (5) for dipolar SLE with $\kappa = 2$.

In order to compute the winding probabilities and to find the distribution on the covering space $\lambda(p, x)$, we solve (15) with boundary conditions as indicated above. The idea is to write $\omega(p, x) = \alpha(p, x)\psi(p, x)$ and impose that terms proportional to $\partial \psi(p, x) / \partial x$ cancel, which, for arbitrary κ , yields $\alpha(p, x) \propto \epsilon(p, x)^{-2/\kappa}$ [21]. It turns out that the function

$\psi(p, x)$ is solution to

$$\frac{\partial \psi(p, x)}{\partial p} = \left(\frac{2 - \kappa}{2\kappa} \right) \frac{\partial v(p, x)}{\partial x} \psi(p, x) + \frac{\kappa}{2} \frac{\partial^2 \psi(p, x)}{\partial x^2}. \tag{24}$$

Hence for $\kappa = 2$ we have a simple diffusion equation $\partial \psi(p, x) / \partial p = \partial^2 \psi(p, x) / \partial x^2$ with initial condition

$$\psi(0, x) = \epsilon(0, x) \omega(0, x) = 2\pi \sum_{n=-\infty}^{\infty} \delta(x - \pi - 2\pi n) \Theta(x). \tag{25}$$

In this case the solution therefore is immediate:

$$\psi(p, x) = \int_{-\infty}^{\infty} \frac{dy}{\sqrt{4\pi p}} e^{-(x-y)^2/4p} \psi(0, y) = \sqrt{\frac{\pi}{p}} \sum_{n=0}^{\infty} e^{-(x-\pi(2n+1))^2/4p}. \tag{26}$$

Thus, for $\kappa = 2$ the endpoint probability reads


$$\begin{aligned} \omega_2(p, x) &= \frac{\sum_{n=0}^{\infty} \exp(-(x - \pi(2n + 1))^2/4p)}{\sum_{n=-\infty}^{\infty} \exp(-(x - \pi(2n + 1))^2/4p)} \\ &= \frac{\sum_{n=0}^{\infty} \exp(\pi n(x - \pi(n + 1))/p)}{\sum_{n=-\infty}^{\infty} \exp(\pi n(x - \pi(n + 1))/p)}. \end{aligned} \tag{27}$$

Note that $\omega(p, x) = \psi(p, x) / (\psi(p, x) + \psi(p, -x))$. The endpoint distribution is given by

$$\lambda_2(p, x) = \frac{\sum_{n,m=0}^{\infty} (m+n+1) \exp(-\frac{(x-\pi-2\pi n)^2+(x+\pi+2\pi m)^2}{4p})}{(p \epsilon(p, x) / \pi)^2}. \tag{28}$$

This result is exact and allows to analyse for $\kappa = 2$ how the SLE trace wraps around the cylinder. An illustration of the distributions is given in Fig. 3. Some remarks are at order. First of all, we have checked through tedious calculation that (28) inserted in (10) correctly reproduces (23). Second, the very particular form of (27), given by a ratio of propagators for one-dimensional Brownian motion, suggests that it might be obtained as a conditional probability for a simple one-dimensional process. Indeed, such a relation can be established (see Sect. 3.3). Moreover, notice an interesting relation to (boundary) conformal field theory. Setting $q = \exp(-2\pi^2/p)$, we see that the different terms in (27) contain $q^{h_{1,n+1}}$ where

Fig. 3 The endpoint distributions $\Lambda_2(p, x)$ and $\lambda_2(p, x)$ for SLE₂ in the case $p = \pi$



$h_{1,n} = \dim \psi_{1,n} = n(n - 1)/2$ are the scaling dimensions of the operators $\psi_{1,n}$, creating $n - 1$ curves at the boundary, for the $O(N = -2)$ model [14]. Finally, one may hope that the simplifications in the case $\kappa = 2$ for the endpoint probabilities may be extended to bulk probabilities of the cylinder and the covering space, a case which we treat (partially) in Appendix 3.

The closed forms (27) and (28) converge well in the dipolar limit $p \rightarrow 0^+$, however are not suitable for computations in the radial limit $p \rightarrow +\infty$. Therefore, we study the moment generating function $\langle e^{-\mu x} \rangle$ here below, and shall give an exact expression adapted to both limits. To this end, it will be useful to consider $\omega_2(p, z) = \psi(p, z)/\epsilon(p, z)$ for complex z . From (27) we see that it is periodic with period $2ip$. Moreover, since the denominator can be written as an infinite product [1]

$$\epsilon(p, z) \propto \prod_{k=1}^{\infty} (1 - e^{iz-(2k-1)p}) (1 - e^{-iz-(2k-1)p}) \tag{29}$$

(up to a p -dependent prefactor), we conclude that the function $\omega_2(p, z)$ has simple poles in the complex plane for $z_{k,\ell} = (2k - 1)ip + 2\pi\ell$, $k, \ell \in \mathbb{Z}$. In order to compute the generating function, we write

$$\langle e^{-\mu x} \rangle = \mu \int_{-\infty}^{\infty} dx e^{-\mu x} \omega_2(p, x), \quad \mu > 0. \tag{30}$$

Deformation of the integration contour in the complex plane from \mathbb{R} to $\mathbb{R} + 2ip$ leads to two distinct contributions. First, the integration along $\mathbb{R} + 2ip$ simply yields $e^{-2ip\mu} \langle e^{-\mu x} \rangle$ because of the periodicity of $\omega_2(p, z)$. Second, we must take into account the simple poles at $z_\ell = z_{1,\ell} = ip + 2\pi\ell$, $\ell \in \mathbb{Z}$ and compute the corresponding residues. Solving for $\langle e^{-\mu x} \rangle$, we find

$$\begin{aligned} \langle e^{-\mu x} \rangle &= \frac{2\pi i \mu}{1 - e^{-2ip\mu}} \sum_{\ell \in \mathbb{Z}} \operatorname{res}_{z=z_\ell} e^{-\mu z} \omega_2(p, z) \\ &= \frac{\pi \mu}{\sin p\mu} \sum_{\ell \in \mathbb{Z}} e^{-2\pi\mu\ell} \psi(p, ip + 2\pi\ell) \left(\frac{\partial \epsilon(p, ip + 2\pi\ell)}{\partial z} \right)^{-1}. \end{aligned} \tag{31}$$

The sum can explicitly be computed because the derivative of $\epsilon(p, z)$ at $z = z_\ell$ does not depend on ℓ by periodicity. After a little algebra we obtain

$$\langle e^{-\mu x} \rangle = \frac{p\mu}{\sin p\mu} \frac{1}{\sinh \pi\mu} \frac{\sum_{\ell=0}^{\infty} (-1)^\ell \sinh((2\ell + 1)\pi\mu) e^{-\pi^2\ell(\ell+1)/p}}{\sum_{\ell=0}^{\infty} (-1)^\ell (2\ell + 1) e^{-\pi^2\ell(\ell+1)/p}}. \tag{32}$$

For large p it is convenient to transform this expression into a dual series

$$\langle e^{-\mu x} \rangle = \frac{e^{p\mu^2} \pi \mu}{\sinh \pi \mu} \frac{1}{\sin p\mu} \frac{\sum_{\ell=0}^{\infty} (-1)^\ell \sin((2\ell + 1)p\mu) e^{-p\ell(\ell+1)}}{\sum_{\ell=0}^{\infty} (-1)^\ell (2\ell + 1) e^{-p\ell(\ell+1)}} \tag{33}$$

by using identities for Jacobi theta functions [1] or Poisson’s summation formula. Let us now study the radial and dipolar limit of our result.

Limit $p \rightarrow +\infty$. In the radial limit, we may write (33) as

$$\langle e^{-\mu x} \rangle = \frac{\pi \mu e^{p\mu^2}}{\sinh \pi \mu} (1 + 2e^{-2p} (1 - \cos 2p\mu)) + O(e^{-4p}). \tag{34}$$

In fact, it is possible to recover this result by approximation of $\epsilon(p, x)$ with its series expansion (19) and Gaussian integration. The result turns out to be finite result for all μ since the probability decays as a Gaussian at large x , i.e. $1 - \omega_2(p, x) = \omega_2(p, -x) \sim \exp(-(x + \pi)^2/4p)$ as $x \rightarrow +\infty$, up to a prefactor periodic in x . We take the logarithm and find a cumulant expansion

$$\begin{aligned} \log\langle e^{-\mu x} \rangle &= \sum_{k=1}^{\infty} (-1)^k \langle x^k \rangle_c \frac{\mu^k}{k!} \\ &= p\mu^2 + \log\left(\frac{\pi\mu}{\sinh \pi\mu}\right) + 2e^{-2p}(1 - \cos 2p\mu) + O(e^{-4p}). \end{aligned} \tag{35}$$

The result shows that, as announced above, we do not recover a Gaussian distribution in the limit $p \rightarrow +\infty$, in contrast to the simple argument from radial SLE. It rather provides all the constants $X_{2n,\kappa=2}$ from the Taylor series expansion of the function $\log[\pi\mu/(\sinh \pi\mu)]$ with respect to μ . For instance, we find

$$\langle x^2 \rangle_c = 2p - \pi^2/3 + 8p^2e^{-2p} + O(e^{-4p}), \tag{36}$$

$$\langle x^4 \rangle_c = 2\pi^4/15 - 32p^4e^{-2p} + O(e^{-4p}) \tag{37}$$

up to corrections of the form $p^n e^{-4p}$.

Limit $p \rightarrow 0^+$. In the dipolar limit, we expand the cumulant generating function, using (32), as

$$\log\langle e^{-\mu x} \rangle = \log\left(\frac{p\mu}{\sin p\mu}\right) - 4\sinh^2(\pi\mu)e^{-2\pi^2/p}. \tag{38}$$

Hence the first two non-vanishing cumulants are given by

$$\langle x^2 \rangle_c = p^2/3 - 8\pi^2e^{-2\pi^2/p} + O(e^{-4\pi^2/p}), \tag{39}$$

$$\langle x^4 \rangle_c = 2p^4/15 - 32\pi^4e^{-2\pi^2/p} + O(e^{-4\pi^2/p}). \tag{40}$$

Since the first corrections to the dipolar result at small p originate from erasure of a single loop wrapping around the cylinder, it seems plausible that the probability for erasing such a loop behaves like $\exp(-2\pi^2/p)$ in the limit $p \rightarrow 0^+$. This factor has indeed the same magnitude as the probability that planar Brownian motion on \mathbb{T}_p winds by at least 2π around the cylinder (we recall the winding distribution for Brownian motion in Appendix A.1.1).

Notice that the dipolar limit presents a subtlety, which does not allow to conclude immediately from expansion of the probability $\omega_2(p, x)$. For any fixed x such that $-\pi < x < \pi$ we find estimate as $p \rightarrow 0^+$ (a neighbourhood of size $\sim p$ of the points $\pm\pi$ must be excluded):

$$\omega_2(p, x) = \frac{1}{1 + e^{-\pi x/p}} + \tanh\left(\frac{\pi x}{2p}\right)e^{-2\pi^2/p} + o(e^{-2\pi^2/p}). \tag{41}$$

Up to exponential corrections, and uniformly in the interval $]-\pi, \pi[$, we recover the dipolar result (5) for $\kappa = 2$. However, the approximation is not uniform beyond this interval what leads to complications when computing moments by using this approximation.

3.3 Probabilistic Argument for $\kappa = 2$

The simplifications for SLE_2 seem surprising at first sight. Here we shall try to understand the underlying mechanism by giving a probabilistic argument for the simple solution of (15) by (27) in the case $\kappa = 2$.

Let us consider the stochastic process $X_t = g_t(x + ip) - i(p - t) - \sqrt{\kappa}B_t$ with initial condition $X_0 = x$. It describes the motion of a given point $x + ip$ on the upper boundary of the cylinder under the flow $g_t(z)$ and is solution of the stochastic differential equation

$$dX_t = v(p - t, X_t)dt + \sqrt{\kappa}dB_t. \tag{42}$$

In particular, it has been shown that $\lim_{t \rightarrow p^-} X_t = (2k + 1)\pi$ with $k \in \mathbb{Z}$ (almost surely) [33]. We may relate the probability that the SLE trace hits the upper boundary of the covering space within $(-\infty, x]$ to X_t via

$$\begin{aligned} \omega(p, x) &= \mathbb{P}[\text{Re } \gamma_p \in (-\infty, x]] = \mathbb{P}[\text{Re } g_t(\gamma_p) \in (-\infty, \text{Re } g_t(x + ip))] \\ &\stackrel{t \rightarrow p}{=} \mathbb{P}[\sqrt{\kappa} B_p \in (-\infty, \text{Re } g_p(x + ip))] = \mathbb{P}_x[X_p \geq 0], \end{aligned} \tag{43}$$

where \mathbb{P}_x denotes ‘‘probability’’, taking into account that the process starts from x . Along these lines we have used the conformal invariance of the SLE measure, leading to invariance of probabilities under conformal transport.

We now show that the right hand side of (43) may be computed easily by establishing a relation to conditioned Brownian motion. Consider the process $x_t = \sqrt{\kappa}B_t$ of simple Brownian motion with diffusion constant κ on the interval $I = [-\pi, \pi]$, identifying its endpoints, which starts from $x_0 = x \in I$. We introduce the propagator via $P(y, t; x)dy = \mathbb{P}_x[x_t \in [y, y + dy]]$. It is solution of the diffusion equation $\partial P(y, t; x)/\partial t = (\kappa/2)\partial^2 P(y, t; x)/\partial x^2$, and explicitly given by

$$P(y, t; x) = \frac{1}{\sqrt{2\pi\kappa t}} \sum_{n=-\infty}^{\infty} \exp\left(-\frac{(y - x - 2\pi n)^2}{2\kappa t}\right). \tag{44}$$

Let us condition x_t to arrive at $x_p = \pi \pmod{2\pi}$ for some given time $p > 0$. Using elementary facts about conditional probabilities we see that the new process has a propagator $Q(y, t; x)$ defined via

$$\begin{aligned} Q(y, t; x)dy &= \mathbb{P}_x[x_t \in [y, y + dy] | x_p = \pi \pmod{2\pi}] \\ &= \frac{P(\pi, p - t; y)}{P(\pi, p; x)} P(y, t; x)dy, \quad \text{for } 0 < t < p. \end{aligned} \tag{45}$$

The conditioning therefore leads to a drift [16] that can be read off from the diffusion equation for $Q(y, t; x)$:

$$\frac{\partial Q(y, t; x)}{\partial t} = -\kappa \frac{\partial}{\partial y} \left(\left(\frac{\partial}{\partial y} \ln P(\pi, p - t; y) \right) Q(y, t; x) \right) + \frac{\kappa}{2} \frac{\partial^2 Q(y, t; x)}{\partial y^2} \tag{46}$$

Specialising to $\kappa = 2$, we conclude that the conditioned process is solution of the stochastic differential equation

$$\begin{aligned} dx_t &= 2 \left(\frac{\partial}{\partial x} \ln P(\pi, p - t; x_t) \right) dt + \sqrt{2} dB_t \\ &= v(p - t, x_t)dt + \sqrt{2} dB_t. \end{aligned} \tag{47}$$

Hence we see that in the case $\kappa = 2$ the motion X_t induced by the Loewner mapping $g_t(z)$ on the upper boundary of the cylinder is the same as Brownian motion x_t on a circle, starting from x and conditioned to visit $\pi \pmod{2\pi}$ at time $t = p$. However, it is a simple exercise to compute the statistics of the winding number of the latter. Again, we shall say that x_t has winding number $N = n$ if its equivalent on the covering space \mathbb{R} of I arrives at $x + 2\pi n$ at time t . From simple conditioning, we have

$$P_x[N = n \mid x_p = \pi \pmod{2\pi}] = \frac{\exp(-(\pi - x + 2\pi n)^2/2\kappa p)}{\sum_{n=-\infty}^{\infty} \exp(-(\pi - x + 2\pi n)^2/2\kappa p)}. \tag{48}$$

Summation over n from 0 to ∞ yields the announced result for $\kappa = 2$

$$\omega(p, x) = P_x[X_p \geq 0] = \frac{\sum_{n=0}^{\infty} \exp(-(x - \pi(2n + 1))^2/4p)}{\sum_{n=-\infty}^{\infty} \exp(-(x - \pi(2n + 1))^2/4p)}. \tag{49}$$

Thus we have shown that we may indeed reinterpret the winding of SLE_2 in terms of the winding of a conditioned one-dimensional Brownian motion. In fact, this is the profound reason explaining why the preceding transformation leads to simplifications for $\kappa = 2$.

3.4 The Winding of Loop-Erased Random Walks with Fixed Endpoints

In this Section we consider the case of a LERW with fixed endpoints from 0 to $x + ip$, $-\pi \leq x \leq \pi$ in the cylinder geometry. On the covering space, it is thus allowed to exit at $x + 2\pi N + ip$ with arbitrary winding number $N \in \mathbb{Z}$. We shall be interested in the law of N which may be obtained by conditioning LERWs to exit at the given boundary points.

Let us ask for the probability that the LERW has made $N = n$ windings if we condition its trace to exit on a subinterval $[a, b] + ip$, $-\pi \leq a, b \leq \pi$ on the upper boundary. It is given by

$$\begin{aligned} P[N = n \mid \text{Re } \gamma_p \in [a, b]] &= \frac{P[N = n \cap \text{Re } \gamma_p \in [a, b]]}{P[\text{Re } \gamma_p \in [a, b]]} \\ &= \frac{\int_a^b dx \lambda_2(p, x + 2\pi n)}{\int_a^b dx \Lambda_2(p, x)}, \end{aligned} \tag{50}$$

where $\Lambda_2(p, x)$ and $\lambda_2(p, x)$ are the distributions defined in (23,28). Taking the limit $a, b \rightarrow x$ amounts to forcing the LERW to exit at $x + ip$. It has been noted previously that (at least for simply-connected domains) this procedure of conditioning leads to chordal SLE_2 from 0 to $x + ip$ [3, 9, 25]. Supposing that this property holds for doubly-connected domains as well, we obtain the statistics of the winding number of SLE_2 with fixed endpoints 0 and $x + ip$ on the boundary of \mathbb{T}_p :

$$P_{0 \rightarrow x+ip}[N = n] = \frac{\lambda_2(p, x + 2\pi n)}{\Lambda_2(p, x)} = \frac{\lambda_2(p, x + 2\pi n)}{\sum_{k=-\infty}^{\infty} \lambda_2(p, x + 2\pi k)}, \tag{51}$$

where $-\pi < x < \pi$. This result is non-trivial (at least for us), since there does not seem to exist an obvious way to recover this probability law from considerations of underlying random walks/Brownian motions.

As an application of (51) we compute the generating function for the moments of N . For $\mu > 0$ let us write

$$\begin{aligned} \langle e^{-\mu N} \rangle &= \frac{1}{\Lambda_2(p, x)} \sum_{n=-\infty}^{\infty} e^{-\mu n} \lambda_2(p, x + 2\pi n) \\ &= \frac{1}{\Lambda_2(p, x)} \frac{\partial}{\partial x} \left(\frac{1}{\epsilon(p, x)} \sum_{n=-\infty}^{\infty} e^{-\mu n} \psi(p, x + 2\pi n) \right) \\ &= \frac{1}{\Lambda_2(p, x)} \frac{e^{p\mu^2/4\pi^2}}{2 \sinh(\mu/2)} \frac{\partial}{\partial x} \left(\frac{e^{\mu x/2\pi} \epsilon(p, x + p\mu/\pi)}{\epsilon(p, x)} \right). \end{aligned} \tag{52}$$

Strictly speaking, the calculation is valid for $\mu > 0$ but the result extends to negative μ , too. The result may be used in order to study the winding behaviour in the radial and dipolar limit.

Radial limit. For $p \rightarrow +\infty$, we find a generating function for the cumulants given by

$$\log \langle e^{-\mu N} \rangle = \frac{\mu x}{2\pi} + \frac{p\mu^2}{4\pi^2} + \log \left(\frac{\mu/2}{\sinh(\mu/2)} \right) + O(e^{-p}), \tag{53}$$

where we have used in that limit $\Lambda_2(p, x) \rightarrow 1/(2\pi)$. As in Sect. 3.2 the cumulants $\langle N^n \rangle_c$ with $n > 2$ converge to constants which can be obtained from the Taylor series expansion of the function $\log[\mu/(2 \sinh(\mu/2))]$ with respect to μ . Corrections to this limit are exponentially small $\propto e^{-p}$.

Dipolar limit. Having in mind the discussion in Sect. 3.2, we expect the probability distribution of N to be concentrated at $N = 0$ as $p \rightarrow 0^+$ (independently of the value taken by $|x| < \pi$), and the probabilities for events occurring with $N \neq 0$ to be exponentially small $\propto e^{-2\pi^2/p}$. By means of explicit computation we find

$$\langle e^{-\mu N} \rangle = 1 + 4 \sinh \left(\frac{\mu}{2} \right) \left(2 \sinh \left(\frac{2\pi x + \mu p}{2p} \right) + \sinh \left(\frac{4\pi x + \mu p}{2p} \right) \right) e^{-2\pi^2/p} \tag{54}$$

up to corrections of the order of $O(e^{-4\pi^2/p})$, what confirms the preceding discussion.

4 Endpoint Probabilities for $\kappa = 4$ on the Cylinder

It is surprising that we obtain the endpoint distribution on the cylinder $\kappa = 2$ by periodisation of the equivalent dipolar result (5). For $\kappa = 4$ we have applied the same idea to look for a solution in the form of a 2π -periodised result found in dipolar SLE_4 . These two values seem to be the only ones for which the periodisation yields exact results. The case $\kappa = 4$ is related to the massless free field theory with central charge $c = 1$. In fact, SLE_4 curves arise as zero lines of the field $\varphi(z, \bar{z})$ emerging from discontinuities in Dirichlet boundary conditions. Moreover, SLE_4 arises in the scaling limit of several lattice models such as the harmonic explorer and domino tilings [24, 31]. A variant of the harmonic explorer leading SLE_4 in doubly-connected domains as discussed in this article was suggested in [33].

In this section, we shall show that the periodic endpoint distribution

$$\Lambda_4(p, x) = \frac{1}{2p} \sum_{n=-\infty}^{\infty} \frac{1}{\cosh[\frac{\pi(x+2\pi n)}{2p}]} \tag{55}$$

is solution of (12) with $\kappa = 4$. Let us notice, that (55) corresponds to the endpoint distribution of two-dimensional Brownian motion on the cylinder with reflecting boundary conditions for $y = 0$, stopped for the first time when it reaches altitude $y = p$ (see Appendix A.1.2).

We analytically continue $\Lambda_4(p, z)$ to complex $z \in \mathbb{C}$ and introduce the function

$$g(p, z) = \frac{\partial \Lambda_4(p, z)}{\partial p} - \frac{\partial}{\partial z}(v(p, z)\Lambda_4(p, z)) - 2\frac{\partial^2 \Lambda_4(p, z)}{\partial z^2} \tag{56}$$

with $v(z, p)$ as defined previously. If $\Lambda_4(p, z)$ is a solution of (12) with $\kappa = 4$ then $g(p, z)$ must vanish for $z \in \mathbb{R}$. We shall even show that $g(p, z)$ vanishes for $\Lambda_4(p, z)$ defined in (55) for all $z \in \mathbb{C}$.

$\Lambda_4(p, z)$ is elliptic with periods $\omega_1 = 2\pi$ and $\omega_2 = 4pi$ and antiperiodic with respect to the half-period $\omega_2/2 = 2pi$. Thus, $g(p, z)$ is elliptic for all p with the same periods ω_1 and ω_2 (as $z \rightarrow z + 4pi$, the additional term arising from the quasi-periodicity of $v(p, z)$ is compensated by the derivative of Λ_4 with respect to p). We may therefore restrict our analysis to the rectangle defined via $-\pi < \text{Re } z < \pi$ and $0 < \text{Im } z < 4p$. The poles of $\Lambda_4(p, z)$, $v(p, z)$ and their derivatives are located at $z = ip$ and $z = 3ip$. The principal parts of the Laurent series expansions at $z = ip$ for the different terms in (56) read

$$\begin{aligned} \frac{\partial \Lambda_4(p, z)}{\partial p} &= \frac{1}{\pi(z - ip)^2} + O(1), & \frac{\partial^2 \Lambda_4(p, z)}{\partial z^2} &= -\frac{2i}{\pi(z - ip)^3} + O(1), \\ \frac{\partial(v(p, z)\Lambda_4(p, z))}{\partial z} &= \frac{1}{\pi(z - ip)^2} + \frac{4i}{\pi(z - ip)^3} + O(1). \end{aligned} \tag{57}$$

Insertion into (56) therefore shows that all divergent terms cancel out so that $g(p, z)$ has a removable singularity at $z = ip$. The Laurent expansion around $z = 3ip$ leads to the same result. In fact, we have

$$\begin{aligned} \frac{\partial \Lambda_4(p, z)}{\partial p} &= -\frac{3}{\pi(z - 3ip)^2} + O(1), & \frac{\partial^2 \Lambda_4(p, z)}{\partial z^2} &= \frac{2i}{\pi(z - 3ip)^3} + O(1), \\ \frac{\partial(v(p, z)\Lambda_4(p, z))}{\partial z} &= -\frac{3}{\pi(z - 3ip)^2} - \frac{4i}{\pi(z - 3ip)^3} + O(1). \end{aligned} \tag{58}$$

Therefore $g(p, z)$ has a removable singularity at $z = 3pi$, too. Hence it must be a constant $g(p, z) \equiv A(p)$ what follows from Liouville’s theorem [2]. However, because of $g(p, z) = -g(p, z + 2pi)$ we find $A(p) = 0$ for all p , so that $g(p, z) \equiv 0$ which is equivalent to say that for $\kappa = 4$ the given form for $\Lambda_4(p, x)$ is the correct probability distribution function.

Let us note that (55) may be written in terms of the Burgers potential $\epsilon(p, x)$ as

$$\Lambda_4(p, x) = \mathcal{N}(p) \frac{\epsilon(p, x - \pi)}{\epsilon(p, x)}. \tag{59}$$

The normalisation factor $\mathcal{N}(p)$ may be found in a similar way as (32) and (33), and is given by

$$\mathcal{N}(p) = \frac{1}{\pi \epsilon(p, ip - \pi)} \frac{\partial \epsilon(p, ip)}{\partial z} = \frac{1}{\pi} \frac{\sum_{n \in \mathbb{Z}} (-1)^n n e^{-n(n+1)p}}{\sum_{n \in \mathbb{Z}} e^{-n(n+1)p}}. \quad (60)$$

However, despite this very suggestive form of $\Lambda_4(p, x)$ as a ratio of two solutions to the simple diffusion equation we have not found algebraic simplifications in order to solve the partial differential equations for $\kappa = 4$ with general boundary conditions.

5 Conclusion

In this paper, we have studied winding properties of loop-erased random walks around a finite cylinder by means of stochastic Loewner evolutions. Relating the computation of the endpoint distribution $\lambda(p, x)$ of SLE_κ on a cylinder to a problem of diffusion-advection of a passive scalar in a Burgers flow, we were able to explicitly determine $\lambda_2(p, x)$ in the case $\kappa = 2$. The behaviour in the limit of very thin and very large cylinders was studied and we pointed out a non-Gaussian behaviour of the winding properties for long cylinders. Furthermore, conditioning the loop-erased random walks to exit via a given boundary point, we were able to compute the probability distribution of the winding number for walks with fixed endpoints. We have shown that the somewhat surprising simplifications for $\kappa = 2$ may be related to the winding of conditioned one-dimensional Brownian motion on a circle. Moreover, we have determined cylinder endpoint distribution in the case $\kappa = 4$. A relation to reflected Brownian motion was pointed out. As for $\kappa = 2$, it is related to the periodisation of the dipolar endpoint distribution. However, these seem to be the only values for κ for which this property holds. It remains to see whether closed results can also be obtained for other values of κ . We hope that our results can be useful to test recent conjectures [12] about SLE properties of interfaces in numerical studies in a cylinder geometry.

Acknowledgements We would like to thank Michel Bauer, Denis Bernard and Alan Middleton for very useful discussions. Moreover, we would like to thank the anonymous referees for useful comments. PLD acknowledges support from ARN BLAN05-0099-01.

Appendix 1: Endpoint Distributions for Two-Dimensional Brownian Motion

In this appendix we compute the endpoint distribution for planar Brownian motion $z_t = x_t + iy_t$ on the cylinder, starting from $z = 0$ and stopped as soon as it reaches the altitude $y = p$. For simplicity, we first consider the geometry of an infinite strip $\{z \in \mathbb{C} \mid 0 < \text{Im} z < p\}$ and then periodise along the real axis in order to obtain the results on the cylinder \mathbb{T}_p . The diffusion in x -direction is unconstrained and has a propagator $\mathbb{P}_{x_0=0}[x_t \in [x, x + dx]] = \exp(-x^2/2t) dx / \sqrt{2\pi t}$. For the motion in y -direction, we consider absorbing and mixed boundary conditions.

A.1.1 Absorbing Boundary Conditions

Consider the motion in y -direction, starting from $y_0 = \epsilon > 0$. We shall impose absorbing boundary conditions at $y = 0$ and $y = p$, and condition the process to exit at $y = p$. The propagator $P_A(y, t)$ defined via $P_A(y, t) dy = \mathbb{P}_{y_0=\epsilon}[y_t \in [y, y + dy]]$ is solution of the dif-

fusion equation $\partial P_A(y, t)/\partial t = 1/2 \partial^2 P_A(y, t)/\partial y^2$ and given by

$$P_A(y, t) = \frac{2}{p} \sum_{n=1}^{\infty} \sin\left(\frac{n\pi y}{p}\right) \sin\left(\frac{n\pi \epsilon}{p}\right) \exp\left(-\frac{n^2 \pi^2 t}{2p^2}\right). \tag{61}$$

We obtain the exit-time distribution $f_A(t)$ at $y = p$ from the probability current at this point. However, if we condition the diffusion to exit at $y = p$, we furthermore must divide this current by the exit probability $\mathbb{P}_{y_0=\epsilon}[\text{exit at } y = p] = \epsilon/p$. Hence

$$\begin{aligned} f_A(t) &= -\frac{p}{2\epsilon} \frac{\partial P(y, t; \epsilon)}{\partial y} \Big|_{y=p} = \frac{\pi}{p\epsilon} \sum_{n=1}^{\infty} (-1)^{n+1} n \sin\left(\frac{n\pi \epsilon}{p}\right) \exp\left(-\frac{\pi^2 n^2 t}{2p^2}\right) \\ &= \frac{\pi^2}{p^2} \sum_{n=1}^{\infty} (-1)^{n+1} n^2 \exp\left(-\frac{\pi^2 n^2 t}{2p^2}\right), \quad \text{as } \epsilon \rightarrow 0^+. \end{aligned} \tag{62}$$

We obtain the distribution $\mu_A(p, x)$ of the endpoint by integration of the free propagator in x -direction weighted by $f_A(t)$ with respect to time t :

$$\begin{aligned} \mu_A(p, x) &= \int_0^{\infty} \frac{dt}{\sqrt{2\pi t}} \exp\left(-\frac{x^2}{2t}\right) f_A(t) \\ &= \frac{\pi}{p} \sum_{k=1}^{\infty} (-1)^{k+1} k e^{-\pi k|x|/p} = \frac{\pi}{4p} \frac{1}{(\cosh(\pi x/2p))^2}. \end{aligned} \tag{63}$$

Periodisation in x with period 2π yields the endpoint distribution on the cylinder $\Lambda_2(p, x)$ (23).

A.1.2 Mixed Boundary Conditions

Consider the same problem as above, but impose reflecting boundary conditions at $y = 0$ and absorbing boundary conditions at $y = p$. Let $P_M(y, t)$ the propagator in y -direction, defined via $P_M(y, t)dy = \mathbb{P}_{y_0=0}[y_t \in [y, y + dy]]$. It is solution of the diffusion equation $\partial P_M(y, t)/\partial t = 1/2 \partial^2 P_M(y, t)/\partial y^2$ and given by

$$P_M(t, y) = \frac{2}{p} \sum_{n=0}^{\infty} \cos\left(\frac{(2n+1)\pi y}{2p}\right) \exp\left(-\frac{(2n+1)^2 \pi^2 t}{8p^2}\right). \tag{64}$$

As previously, we compute the exit-time distribution from the probability current at $y = p$:

$$f_M(t) = \frac{\pi}{2p^2} \sum_{n=0}^{\infty} (-1)^n (2n+1) \exp\left(-\frac{(2n+1)^2 \pi^2 t}{8p^2}\right). \tag{65}$$

Consequently the distribution of the exit point at the upper boundary of the strip reads

$$\begin{aligned} \mu_M(p, x) &= \int_0^{\infty} \frac{dt}{\sqrt{2\pi t}} \exp\left(-\frac{x^2}{2t}\right) f_M(t) \\ &= \frac{1}{p} \sum_{n=0}^{\infty} (-1)^n \exp\left(-\frac{(2n+1)\pi|x|}{2p}\right) = \frac{1}{2p} \frac{1}{\cosh(\pi x/2p)}. \end{aligned} \tag{66}$$

Finally, periodisation with respect to x with period 2π leads to $\Lambda_4(p, x)$ for the cylinder \mathbb{T}_p (55).

Appendix 2: Endpoint Distribution for Hexagonal Lattices

The purpose of this appendix is to give some idea about finite-size corrections to the scaling limit on the cylinder. Here we evaluate the discrete equivalent to (23) and compute the leading correction due to lattice effects. In principle, we dispose of a great variety for the choice of the lattice. Finite-size corrections not only turn out to be dependent on the choice of geometry, but on the lattice type as well. We shall content ourselves with the honeycomb lattice—the most common choice.

We consider a finite lattice tube of $m \times l$ honeycombs, as shown in Fig. 4a. We decompose the honeycomb lattice into two triangular sub-lattices whose sites are coloured by \bullet and \circ . Nearest neighbours of a \bullet -site thus belong to the \circ -lattice and vice versa. We study lattice walks starting from a \bullet -site at the lower boundary to some any \circ -site at the upper boundary and shall impose absorbing boundary conditions on \circ -sites at either boundary. In fact, it shall be convenient to map the problem to a “brick wall” lattice as shown on Fig. 4b, and introduce a suitable coordinate system. Then it is straightforward to write the master equations for the probability $P_{\bullet/\circ}(x, y; t)$ that the walker starting from $(0, 0)$ can be found at (x, y) at time t :

$$P_o(x, y; t + 1) = \frac{1}{3}(P_\bullet(x + 1, y; t) + P_\bullet(x - 1, y; t) + P_\bullet(x, y - 1; t)), \tag{67}$$

$$P_\bullet(x, y; t + 1) = \frac{1}{3}(P_o(x + 1, y; t) + P_o(x - 1, y; t) + P_o(x, y + 1; t)) \tag{68}$$

with boundary conditions $P_o(x, y = 0; t) = P_o(x, y = l; t) = 0$ and $P_\bullet(x, y; t = 0) = \delta_{x,0}\delta_{y,0}$ where $\delta_{i,j}$ denotes the Kronecker symbol which takes the value 1 if $i = j$, and 0 otherwise. Moreover, identification of x and $x + 2m$ leads to periodicity $P_o(x + 2m, y; t) = P_o(x, y; t)$, $P_\bullet(x + 2m, y; t) = P_\bullet(x, y; t)$. We shall need the local occupation times $G_{\bullet/\circ}(x, y) = \sum_{t=0}^\infty P_{\bullet/\circ}(x, y; t)$. The lattice walker is stopped if it walks from some \bullet -site at $y_e = 1$ or $y_e = l - 1$ to a \circ -site at $y_e = 0$ or $y_e = l$ respectively, what happens with probability $1/3$. Therefore, the exit probability at the upper boundary is given by

$$P[x_e = x, y_e = l] = \frac{1}{3}G_\bullet(x, l - 1), \tag{69}$$

where x_e denotes the (random) horizontal position of the exiting walker. Notice that x_e only takes odd/even integer values for even/odd l . Combining (67, 68), it is possible to eliminate




Fig. 4 (a) Cylindrical lattice domain of 5×3 honeycombs with walk from lower to upper boundary. The highlighted curve represent some walk from the lower to the upper boundary. (b) Coordinate system for equivalent “brick wall” lattice

$G_\bullet(x, y)$ and find the second-order difference equation

$$G_\bullet(x, y) = \frac{1}{6}(G_\bullet(x + 2, y) + G_\bullet(x + 1, y + 1) + G_\bullet(x + 1, y - 1) + G_\bullet(x - 2, y) + G_\bullet(x - 1, y + 1) + G_\bullet(x - 1, y - 1)), \tag{70}$$

for $0 < y < l - 1$. Using the boundary conditions, we furthermore find the equations

$$G_\bullet(x, 0) = \frac{1}{8}(G_\bullet(x + 1, 1) + G_\bullet(x - 1, 1)) + \frac{9}{8} \delta_{x,0} \tag{71}$$

$$G_\bullet(x, l - 1) = \frac{1}{7}(G_\bullet(x + 1, l - 2) + G_\bullet(x + 2, l - 1) + G_\bullet(x - 1, l - 2) + G_\bullet(x - 2, l - 1)). \tag{72}$$

Solving (70), (71) and (72) all together leads to a rather lengthy and tedious calculation. The reader may find a detailed account of the strategy with applications to other lattice types in [22]. Here we only sketch the solution.

First, we solve the bulk equation (70) by a separation ansatz $G_\bullet(x, y) = P(x)Q(y)$. Forming linear combinations of solutions found from this ansatz, we obtain the general bulk solution

$$G_\bullet(x, y) = A_0 + C_0y + (A_m + C_my)(-1)^{x+y} + \sum_{k \in I_1} (A_k e^{i\alpha_k x} + B_k e^{-i\alpha_k x})(e^{\gamma_k y} + C_k e^{-\gamma_k y}) + \sum_{k \in I_2} (-1)^y (A_k e^{i\alpha_k x} + B_k e^{-i\alpha_k x})(e^{\gamma_k y} + C_k e^{-\gamma_k y}), \tag{73}$$

where A_i, B_i and C_i are constants, $I_1 = \{k \in \mathbb{Z} \mid 0 < k < m/2 \vee 3m/2 < k < 2m\}$ and $I_2 = \{k \in \mathbb{Z} \mid m/2 < k < 3m/2, k \neq m/2\}$ index sets, and

$$\alpha_k = \frac{\pi k}{m}, \quad \cosh \gamma_k = \frac{3 - \cos 2\alpha_k}{2|\cos \alpha_k|} \quad \text{and} \quad k = 0, 1, \dots, 2m - 1 \tag{74}$$

However, we exclude $k = m$. Moreover, for some technical reason the formula only is valid as long as m is odd. The remaining constants are determined by matching (73) to (71) and (72) what yields

$$A_k = B_k = \frac{1}{2m|\cos \alpha_k| \sinh(\gamma_k l)}, \quad C_k = -e^{2\gamma_k(l-1)} \frac{1 + 2e^{\gamma_k} |\cos \alpha_k|}{1 + 2e^{-\gamma_k} |\cos \alpha_k|},$$

and $A_0 = A_m = (3l - 1)/(4ml)$, $C_0 = C_m = -3/(4ml)$. Putting all pieces together, we find the solution for (69). However, if we condition the walk to exit at the upper boundary, we still have to divide by the probability $P[y_e = l]$ what amounts to a normalisation. After some algebra we obtain the final result (for odd m)

$$P[x_e = x \mid y_e = l] = \frac{1}{m} + \sum_{k=1}^{(m-1)/2} \frac{2l \cos(\alpha_k x) \sinh \gamma_k}{m \sinh \gamma_k l}. \tag{75}$$

Having this lattice result, it is interesting to compute its scaling limit and study corrections. In the case where m is odd, x takes even values from $-(m - 1)$ to $m - 1$. Introducing a lattice scale a such that $x = \xi/a$, $\xi \in \mathbb{R}$, we would like to study the scaling limit $a \rightarrow 0^+$, $m, l \rightarrow +\infty$ in such a way that the geometrical height $p = \sqrt{3}(l - 1/3)a$ and width $2\pi = 2am$ remain finite. Therefore $\sqrt{3}l/m = p/\pi + a/(\sqrt{3}\pi)$.

$$\begin{aligned} \Lambda_2(p, \xi, a) &= \frac{1}{2a} \mathbb{P}[x_e = x | y_e = l] = \frac{\sqrt{3}l}{\pi m} \sum_{k=-\infty}^{\infty} \frac{k \cos k\xi}{\sinh(\sqrt{3}(l/m)k)} \\ &= \frac{1}{4\sqrt{3}(l/m)} \sum_{n=-\infty}^{\infty} \frac{1}{\left[\cosh\left(\frac{(x+2\pi n)}{2\sqrt{3}(l/m)}\right) \right]^2} \\ &= \left(1 + \frac{a}{\sqrt{3}} \frac{\partial}{\partial p} \right) \Lambda_2(p, \xi) + O(a^2). \end{aligned} \tag{76}$$

We see that the first correction to the scaling limit is completely naturally proportional to the lattice scale a .

Appendix 3: Analysis in the Bulk

This appendix collects some facts about bulk properties for SLE on the cylinder. Let us denote the probability, that the trace γ passes to the left of some given point z on the covering space by $\omega(p, z, \bar{z})$. It is solution of the diffusion equation

$$\frac{\partial \omega(p, z, \bar{z})}{\partial p} = H(p, z) \frac{\partial \omega(p, z, \bar{z})}{\partial z} + H(p, \bar{z}) \frac{\partial \omega(p, z, \bar{z})}{\partial \bar{z}} + \frac{\kappa}{2} \left(\frac{\partial}{\partial z} + \frac{\partial}{\partial \bar{z}} \right)^2 \omega(p, z, \bar{z}) \tag{77}$$

with boundary conditions $\omega(p, z, \bar{z}) \rightarrow 0$ as $\text{Re } z \rightarrow -\infty$ and $\omega(p, z, \bar{z}) \rightarrow 1$ as $\text{Re } z \rightarrow +\infty$. The derivation of (77) is similar to (11) and (12), using an infinitesimal argument. Using the relationship $H(p, z) = v(p, z - ip) - i$ to the velocity field defined above and (16), it is not difficult to show that $H(p, z)$ is a Burgers flow, too. We have

$$\frac{\partial H(p, z)}{\partial p} = H(p, z) \frac{\partial H(p, z)}{\partial z} + \frac{\partial^2 H(p, z)}{\partial z^2}. \tag{78}$$

Using the Cole-Hopf transformation (18), we may write $H(p, z) = \partial U(p, z)/\partial z$ with $U(p, z) = 2 \log \epsilon(p, z - ip) - iz - p/2$.

For $\kappa = 2$ as $\text{Im } z \rightarrow p^-$ we require that $\omega(p, z, \bar{z})$ tends to $\lambda_2(p, x)$ (with $x = \text{Re } z$), and $\omega(p, z, \bar{z}) \rightarrow \Theta(x)$ as $\text{Im } z \rightarrow 0^+$. Although we have not found the solution with these boundary conditions, we have determined a special solution $\tilde{\omega}(p, z, \bar{z})$ with a periodised version of these boundary conditions which has interesting relations to dipolar SLE₂. We ask that for $\text{Im } z \rightarrow p^-$ it becomes $\tilde{\omega}(p, z, \bar{z}) = \int_0^x dx \Lambda_2(p, x) = (p v(p, x) + x)/2\pi$, and that for $\text{Im } z \rightarrow 0^+$ it reproduces the quasi-periodic step function $\tilde{\omega}(p, z, \bar{z}) = \lim_{n \rightarrow \infty} \sum_{k=-n}^n (\text{sign}(x - 2\pi k))/2$.

Obviously, both $H(p, z)$ and $H(p, \bar{z})$ are solutions to (77), however with a simple pole at $z = 0$. Inspired from the boundary condition at $\text{Im } z = p$, we shall consider the function

$f(p, z, \bar{z}) = (\text{Im } z)(H(p, z) + H(p, \bar{z}))$, which has no pole. However, using the differential operator \mathcal{A} defined as

$$\mathcal{A} = \left[\frac{\partial}{\partial p} - H(p, z) \frac{\partial}{\partial z} - H(p, \bar{z}) \frac{\partial}{\partial \bar{z}} - \left(\frac{\partial}{\partial z} + \frac{\partial}{\partial \bar{z}} \right)^2 \right] \tag{79}$$

we see that $\mathcal{A}f(p, z, \bar{z}) = -(H(z)^2 - H(\bar{z})^2)/2i$. Hence we should find some solution to $\mathcal{A}g(p, z) = iH(z)^2/2$ without poles (since this would again lead to a singular solution). Hence we have to search for a suitable $g(p, z)$ with at most a logarithmic singularity at $z = 0$. This can be done by observing that $U(p, z) = 2 \log \epsilon(p, z - ip) - iz - p/2$ is solution to the equation

$$\begin{aligned} \frac{\partial U(p, z)}{\partial p} &= \frac{1}{2} \left(\frac{\partial U(p, z)}{\partial z} \right)^2 + \frac{\partial^2 U(p, z)}{\partial z^2} \\ &= H(p, z) \frac{\partial U(p, z)}{\partial z} + \frac{\partial^2 U(p, z)}{\partial z^2} - \frac{1}{2} H(p, z)^2 \end{aligned} \tag{80}$$

as can be seen from Burgers equation for $H(p, z)$ (78). Hence we have found a solution $\tilde{\omega}(p, z, \bar{z}) = C_1 + C_2(f(p, z, \bar{z}) - i(U(p, z) - U(p, \bar{z})))$, where C_1, C_2 are constants. It is a solution of (77) without poles for $0 \leq \text{Im } z \leq p$. The choice $C_1 = 1/2$ and $C_2 = 1/4\pi$ leads to the desired boundary conditions for $\tilde{\omega}(p, z, \bar{z})$:

$$\tilde{\omega}(p, z, \bar{z}) = \frac{1}{2} + \frac{\text{Im } z}{4\pi} (H(p, z) + H(p, \bar{z})) + \frac{i}{4\pi} (U(p, z) - U(p, \bar{z})) \tag{81}$$

$\tilde{\omega}(p, z, \bar{z})$ is related to $\omega(p, z, \bar{z})$ by periodisation

$$\tilde{\omega}(p, z, \bar{z}) = \lim_{n \rightarrow \infty} \sum_{k=-n}^n \left(\omega(p, z + 2\pi k, \bar{z} + 2\pi k) - \frac{1}{2} \right) \tag{82}$$

i.e. we have only found a (quasi-)periodic version of $\omega(p, z, \bar{z})$.

The dipolar limit $p \rightarrow 0^+$ turns out to be quite interesting. In that limit the quasi-periodicity is irrelevant and one obtains the result for dipolar SLE₂. One can replace in that limit $H(p, z) \rightarrow H_{ss}(p, z) = \frac{\pi}{p} \coth \pi z/2p - z/p$, the single shock expression, and the corresponding expression $2 \log \epsilon(p, z) \sim -z^2/(2p) + 2 \log \cosh \pi z/2p$, up to a p -dependent unimportant constant. One then easily obtains from (81):

$$\tilde{\omega}(p, z, \bar{z}) + \frac{1}{2} \sim \omega_d(p, z, \bar{z}) = 1 - \frac{1}{\pi} \text{Im} \log \sinh \frac{\pi z}{2p} + \frac{\text{Im } z}{4p} \left(\coth \frac{\pi z}{2p} + \coth \frac{\pi \bar{z}}{2p} \right). \tag{83}$$

This is nothing but the probability that dipolar SLE₂ passes to the left of a given point z within a strip of height p . It satisfies the general equation:

$$\left[\frac{\pi}{p} \coth \frac{\pi z}{2p} \frac{\partial}{\partial z} + \frac{\pi}{p} \coth \frac{\pi \bar{z}}{2p} \frac{\partial}{\partial \bar{z}} + \frac{\kappa}{2} \left(\frac{\partial}{\partial z} + \frac{\partial}{\partial \bar{z}} \right)^2 \right] \omega_d(p, z, \bar{z}) = 0 \tag{84}$$

studied in [7] where solutions could be found only for $\kappa \geq 4$ (in the form of simple harmonic functions). Here we recover, from a limit of a more general object defined on \mathbb{T}_p , a result for $\kappa = 2$ obtained only very recently by a rather different method [8]. The fact that (83) satisfies both (84) (an equation with no ∂_p terms) and the dipolar (i.e. single shock) limit

of (77) (an evolution equation as a function of p) is easily understood by noting that it also satisfies $(p\partial_p + z\partial_z + \bar{z}\partial_{\bar{z}})\omega_d(p, z, \bar{z}) = 0$. This equation expresses nothing but dilatation invariance of the dipolar SLE, explicitly broken by the periodicity of cylinder geometries.

One may wonder whether there is a way to extend the known harmonic solutions [7] of the dipolar limit, to the full cylinder. Here we only present the case $\kappa = 4$ which can be constructed by integration of the result given in Sect. 4. In fact, this amounts to periodise the result from dipolar SLE for a strip of height p

$$\omega_d(p, z, \bar{z}) = 1 - \frac{1}{\pi} \operatorname{Im} \log \tanh \frac{\pi z}{4p} \tag{85}$$

in an appropriate way. Symmetric periodisation and leads to

$$\tilde{\omega}(p, z, \bar{z}) = \lim_{n \rightarrow \infty} \sum_{k=-n}^n \left(\frac{1}{2} - \frac{1}{\pi} \operatorname{Im} \log \tanh \left(\frac{\pi(z + 2\pi k)}{4p} \right) \right) \tag{86}$$

which is a harmonic solution to (77) with required boundary conditions at $z = x + ip$

$$\tilde{\omega}(p, x + ip, x - ip) = \int_0^x dx \Lambda_4(p, x). \tag{87}$$

Appendix 4: Path Integral for General κ

Let us indicate how one can take advantage of (24) to express the solution of (12) as a path integral for general κ . We may write

$$\psi(p, x) = \int_{-\infty}^{\infty} dy K(p, x; y) \psi(0, y), \tag{88}$$

where $K(p, x; y)$ is the Euclidean propagator for (24). It given by a path-integral representation

$$K(p, x; y) = \int_{x(0)=y}^{x(p)=x} [dx(t)] e^{-S[x(t)]}, \tag{89}$$

where the action is defined as

$$S[x(t)] = \int_0^p dt \left(\frac{\dot{x}(t)^2}{2\kappa} + \frac{\kappa - 2}{2\kappa} \frac{\partial v(t, x(t))}{\partial x} \right). \tag{90}$$

Hence, we find a nice path-integral representation [19, 20] for the general solution $\omega(p, x)$ of (15):

$$\omega(p, x) = \epsilon(p, x)^{-2/\kappa} \int_{-\infty}^{\infty} dy \epsilon(0, y)^{2/\kappa} \omega(0, y) \int_{x(0)=y}^{x(p)=x} [dx(t)] e^{-S[x(t)]} \tag{91}$$

which reproduces the exact solution (27) in the case $\kappa = 2$. It may be useful for perturbation theory around $\kappa = 2$, using the Fourier series decomposition:

$$v(p, x) = \sum_{n=1}^{\infty} \frac{4e^{-np}}{1 - e^{-2np}} \sin(nx) \tag{92}$$

if one treats the apparent singular behaviour of $\epsilon(0, y)^{2/\kappa}$ for $\kappa \neq 2$ (e.g. shifting the integration from 0 to small p_0 and using the known dipolar result for $\omega(p_0, x)$).

References

1. Abramowitz, M., Stegun, I.: Handbook of Mathematical Functions. Dover, New York (1970)
2. Ahlfors, L.: Complex Analysis. McGraw-Hill, New York (1979)
3. Bauer, M., Bernard, D.: private communication
4. Bauer, M., Bernard, D.: Sailing the deep blue sea of burgers turbulence. *J. Phys. A: Math. Gen.* **32**, 5179–5199 (1999)
5. Bauer, M., Bernard, D.: SLE, CFT and zig-zag probabilities. In: Conformal Invariance and Random Spatial Processes. NATO Advanced Study Institute (2003)
6. Bauer, M., Bernard, D.: 2D growth processes: SLE and Loewner chains. *Phys. Rep.* **432**, 115–221 (2006)
7. Bauer, M., Bernard, D., Houdayer, J.: Dipolar SLEs. *J. Stat. Mech.* **0503**, P001 (2005)
8. Bauer, M., Bernard, D., Kennedy, T.: Conditioning SLEs and loop erased random walks. [arXiv:0806.2246](https://arxiv.org/abs/0806.2246) (2008)
9. Bauer, M., Bernard, D., Kytölä, K.: LERW as an example of off-critical SLEs. [arXiv:0712.1952](https://arxiv.org/abs/0712.1952) (2007)
10. Bauer, R.O., Friedrich, R.: On Chordal and Bilateral SLE in multiply connected domains. [arXiv:math.PR/0503178](https://arxiv.org/abs/math.PR/0503178) (2005)
11. Bec, J., Khanin, K.: Burgers turbulence. *Phys. Rep.* **447**, 1–66 (2007)
12. Bernard, D., LeDoussal, P., Middleton, A.A.: Are domain walls in spin glasses described by stochastic loewner evolutions? *Phys. Rev. B* **76**, 020403(R) (2007)
13. Cardy, J.: SLE for theoretical physicists. *Ann. Phys.* **318**, 81–115 (2005)
14. Cardy, J.: The $O(n)$ model on the annulus. *J. Stat. Phys.* **125**, 1–21 (2006)
15. Cole, J.: On a quasilinear parabolic equation occurring in aerodynamics. *Quart. Appl. Math.* **9**, 225 (1951)
16. Doob, J.L.: Classical Potential Theory and Its Probabilistic Counterpart. Springer, New York (1984)
17. Drossel, B., Kardar, M.: Passive sliders on growing surfaces and advection in burgers flows. *Phys. Rev. B* **66**, 195414 (2002)
18. Duplantier, B., Saleur, H.: Winding-angle distributions of two-dimensional self-avoiding walks from conformal invariance. *Phys. Rev. Lett.* **60**, 2343–2346 (1988)
19. Ginanneschi, P.M.: Models of passive and reactive tracer motion: an application of Ito calculus. *J. Phys. A: Math. Gen.* **30**, L519–L523 (1997)
20. Ginanneschi, P.M.: On the mass transport by a Burgers velocity field. *Physica D* **115**, 341–352 (1998)
21. Gorodtsov, V.: Convective heat conduction and diffusion in one-dimensional hydrodynamics. *JETP* **89**(5), 872 (1999)
22. Henri, B., Batchelor, M.: Random walks on finite lattice tubes. *Phys. Rev. E* **68**, 016112 (2003)
23. Hopf, E.: The partial differential equation $u_t + uu_x = u_{xx}$. *Commun. Pure Appl. Math.* **3**, 210 (1950)
24. Kenyon, R.: Conformal invariance of domino tiling. *Ann. Probab.* **28**, 759–795 (2000)
25. Lawler, G.F.: private communication
26. Lawler, G.F.: Conformally Invariant Processes in the Plane. American Mathematical Society, Providence (2005)
27. Lawler, G.F., Schramm, O., Werner, W.: Conformal invariance of planar loop-erased random walks and uniform spanning trees. *Ann. Probab.* **32**, 939–995 (2004)
28. Nagar, A., Majumdar, S.N., Barma, M.: Strong clustering of noninteracting, sliding passive scalars driven by fluctuating surfaces. *Phys. Rev. E* **74**, 021124 (2006)
29. Nehari, Z.: Conformal Mapping. Dover, New York (1982)
30. Schramm, O.: Scaling limits of loop-erased random walks and uniform spanning trees. *Israel J. Math.* **118**, 221–288 (2000)
31. Schramm, O., Sheffield, S.: Contour lines of the two-dimensional discrete Gaussian free field. [math.PR/0605337](https://arxiv.org/abs/math.PR/0605337) (2006)
32. Zhan, D.: Stochastic Loewner evolutions in doubly connected domains. *Prob. Theory Relat. Fields* **129**, 340–380 (2004)
33. Zhan, D.: Some properties of annulus SLE. *Electronic J. Probab.* **11**(41), 1069–1093 (2006)
34. Zhan, D.: The scaling limits of planar LERW in finitely connected domains. [arXiv:math.PR/06100304](https://arxiv.org/abs/math.PR/06100304) (2006)