

Quantum Dynamics with Mean Field Interactions: a New Approach

László Erdős · Benjamin Schlein

Received: 23 April 2008 / Accepted: 13 May 2008 / Published online: 5 March 2009
© Springer Science+Business Media, LLC 2008

Abstract We propose a new approach for the study of the time evolution of a factorized N -particle bosonic wave function with respect to a mean-field dynamics with a bounded interaction potential. The new technique, which is based on the control of the growth of the correlations among the particles, leads to quantitative bounds on the difference between the many-particle Schrödinger dynamics and the one-particle nonlinear Hartree dynamics. In particular the one-particle density matrix associated with the solution to the N -particle Schrödinger equation is shown to converge to the projection onto the one-dimensional subspace spanned by the solution to the Hartree equation with a speed of convergence of order $1/N$ for all fixed times.

Keywords Many-body quantum dynamics · Mean field limit · Nonlinear Hartree equation

1 Introduction

We consider a system of N interacting bosons in ν dimensions described on the Hilbert space $\mathcal{H}_N = L^2_s(\mathbb{R}^{N\nu}, dx_1 \dots dx_N)$, the subspace of $L^2(\mathbb{R}^{N\nu}, dx_1 \dots dx_N)$ consisting of permutation symmetric wave functions. Here the variables $x_1, \dots, x_N \in \mathbb{R}^\nu$ refer to the positions of the N particles. The Hamiltonian is given by

$$H_N = \sum_{j=1}^N (-\Delta_{x_j} + U(x_j)) + \frac{1}{N} \sum_{i < j}^N V(x_i - x_j) \quad (1.1)$$

Dedicated to Jürg Fröhlich on the occasion of his 60th birthday, with admiration and gratitude.

L. Erdős is partially supported by SFB/TR12 Project from DFG.

B. Schlein is on leave from Cambridge University; his research is supported by a Kovalevskaia Award from Me Humboldt Foundation.

L. Erdős · B. Schlein (✉)

Institute of Mathematics, University of Munich, Theresienstr. 39, 80333 Munich, Germany
e-mail: schlein@math.lmu.de

and, under suitable conditions on the potentials U and V , acts as a self-adjoint operator on \mathcal{H}_N . The coupling constant $1/N$ in front of the interaction characterizes mean field models; it guarantees that the kinetic and the potential part of the Hamiltonian are typically of the same order.

The dynamics of the system is governed by the N -particle Schrödinger equation

$$i \partial_t \psi_{N,t} = H_N \psi_{N,t} \Rightarrow \psi_{N,t} = e^{-iH_N t} \psi_N. \tag{1.2}$$

In particular we will be interested in the time evolution of factorized initial data $\psi_N(x_1, \dots, x_N) = \prod_{j=1}^N \varphi(x_j)$. It turns out that, because of the mean field nature of the interaction, for large N , factorization is approximately preserved by the evolution (1.2). In a sense to be made precise, we have $\psi_{N,t} \simeq \varphi_t^{\otimes N}$, for a suitable $\varphi_t \in L^2(\mathbb{R}^v)$. If factorization were indeed preserved, it is simple to check that φ_t must be determined by the solution to the self-consistent nonlinear Hartree equation

$$i \partial_t \varphi_t = (-\Delta + U)\varphi_t + (V * |\varphi_t|^2)\varphi_t, \tag{1.3}$$

with initial data $\varphi_{t=0} = \varphi$.

A more precise analysis shows that the solution $\psi_{N,t}$ can be approximated, in the limit $N \rightarrow \infty$, by products of solutions to the nonlinear Hartree equation (1.3) only if we are interested in quantities depending non-trivially on a finite number of particles. In other words, we cannot expect the L^2 -difference between $\psi_{N,t}$ and $\varphi_t^{\otimes N}$ to converge to zero. What can be expected and, for a large class of potentials, has also been rigorously proven, is that for an arbitrary fixed k -particle observable $J^{(k)}$ (for example, a compact operator on $L^2(\mathbb{R}^{kv})$), and for every fixed time $t \in \mathbb{R}$,

$$\langle \psi_{N,t}, (J^{(k)} \otimes 1^{(N-k)}) \psi_{N,t} \rangle \rightarrow \langle \varphi_t^{\otimes N}, (J^{(k)} \otimes 1^{(N-k)}) \varphi_t^{\otimes N} \rangle = \langle \varphi_t^{\otimes k}, J^{(k)} \varphi_t^{\otimes k} \rangle \tag{1.4}$$

as $N \rightarrow \infty$, where φ_t is the solution to (1.3) with initial data $\varphi_{t=0} = \varphi$. The convergence (1.4) can be interpreted as the convergence of the marginal densities. Recall that, for a given N -particle wave-function ψ_N , we define the density matrix $\gamma_N = |\psi_N\rangle\langle\psi_N|$ as the orthogonal projection onto ψ_N , and, for $k = 1, \dots, N$, we define the k -particle marginal density $\gamma_N^{(k)}$ associated with ψ_N by taking the partial trace of γ_N over the last $N - k$ particles. In other words, $\gamma_N^{(k)}$ is defined for $k = 1, \dots, N$, as a non-negative trace class operator on $L^2(\mathbb{R}^{vk})$ with kernel given by

$$\begin{aligned} \gamma_N^{(k)}(\mathbf{x}_k; \mathbf{x}'_k) &= \int d\mathbf{x}_{N-k} \gamma_N(\mathbf{x}_k, \mathbf{x}_{N-k}; \mathbf{x}'_k, \mathbf{x}_{N-k}) \\ &= \int d\mathbf{x}_{N-k} \psi_N(\mathbf{x}_k, \mathbf{x}_{N-k}) \overline{\psi_N(\mathbf{x}'_k, \mathbf{x}_{N-k})}. \end{aligned} \tag{1.5}$$

Here and henceforth we use the notation $\mathbf{x}_k = (x_1, \dots, x_k)$, $\mathbf{x}'_k = (x'_1, \dots, x'_k)$, $\mathbf{x}_{N-k} = (x_{k+1}, \dots, x_N)$. We will also use the shorthand notation $\mathbf{x} \equiv \mathbf{x}_N = (x_1, \dots, x_N)$. Note that we choose the convention $\text{Tr} \gamma_N^{(k)} = 1$ for all $N \geq 1$, and $1 \leq k \leq N$.

For $k = 1, \dots, N$ denote by $\gamma_{N,t}^{(k)}$ the k -particle marginal density associated with the solution to the N -particle Schrödinger equation (1.2) with factorized initial data $\psi_N = \varphi^{\otimes N}$. In terms of marginal densities, (1.4) is equivalent to the convergence

$$\gamma_{N,t}^{(k)} \rightarrow |\varphi_t\rangle\langle\varphi_t|^{\otimes k} \text{ as } N \rightarrow \infty \tag{1.6}$$

for all fixed $t \in \mathbb{R}$ and $k \geq 1$, with respect to the weak* topology defined on the space of trace class operators. It is simple to check that the weak* convergence in this case is equivalent to the convergence in the trace norm topology, because the limit is a rank one projection.

The first rigorous results of the form (1.4) or (1.6) were obtained by Hepp in [16], and then extended by Ginibre and Velo in [14, 15]. These works are based on the analysis of the time evolution of coherent states (in a second quantized representation of the system). The use of coherent states (characterized by a non-fixed number of particles) makes it possible to isolate the main part of the evolution (described by the Hartree equation) and to study the fluctuations around it. With this method, it was recently proved in [21] that (1.6) can be improved (for example for the case $k = 1$) to the quantitative estimate

$$\text{Tr} |\gamma_{N,t}^{(1)} - |\varphi_t\rangle\langle\varphi_t|| \leq \text{const} \cdot \frac{e^{Ct}}{N^{1/2}} \tag{1.7}$$

for factorized initial data (while for coherent initial data, the results of [21] establish a better bound, proportional to $1/N$) and for interaction potentials V such that $V^2(x) \leq C(1 - \Delta_x)$.

A different approach was proposed by Spohn in [22]. The main idea of the technique introduced in [22] consists in characterizing the limit of the marginal densities $\gamma_{N,t}^{(k)}$ through the evolution equations that they satisfy. For finite N , the dynamics of the marginal densities $\{\gamma_{N,t}^{(k)}\}_{k=1}^N$ is governed by the so-called BBGKY hierarchy

$$\begin{aligned} i \partial_t \gamma_{N,t}^{(k)} &= \sum_{j=1}^k [-\Delta_{x_j} + U(x_j), \gamma_{N,t}^{(k)}] + \frac{1}{N} \sum_{i < j}^k [V(x_i - x_j), \gamma_{N,t}^{(k)}] \\ &\quad + \frac{N - k}{N} \sum_{j=1}^k \text{Tr}_{k+1} [V(x_j - x_{k+1}), \gamma_{N,t}^{(k+1)}], \end{aligned} \tag{1.8}$$

where Tr_{k+1} denotes the partial trace over the $(k + 1)$ -th particle. Using $\text{Tr} \gamma_{N,t}^{(k)} = 1$, an abstract argument shows that the sequence $\{\gamma_{N,t}^{(k)}\}_{k=1}^N$ is compact with respect to a suitable weak topology. The first main step towards a proof of (1.6) is then to show that an arbitrary limit point $\{\gamma_{\infty,t}^{(k)}\}_{k \geq 1}$ of the sequence of marginals $\{\gamma_{N,t}^{(k)}\}_{k=1}^N$ is a solution to the infinite hierarchy

$$i \partial_t \gamma_{\infty,t}^{(k)} = \sum_{j=1}^k [-\Delta_{x_j} + U(x_j), \gamma_{\infty,t}^{(k)}] + \sum_{j=1}^k \text{Tr}_{k+1} [V(x_j - x_{k+1}), \gamma_{\infty,t}^{(k+1)}] \tag{1.9}$$

which is obtained from (1.8) by formally letting $N \rightarrow \infty$. The second main step to show (1.6) consists in proving the uniqueness of the solution to (1.9). Since it is simple to verify that the ansatz $\gamma_t^{(k)} = |\varphi_t\rangle\langle\varphi_t|^{\otimes k}$ is indeed a solution to (1.9) if φ_t satisfies the Hartree equation (1.3), the compactness of the sequence $\{\gamma_{N,t}^{(k)}\}_{k=1}^N$ leads to (1.6). This method, which was first used in [22] to prove (1.6) for bounded interaction, was recently extended to singular potential (see [1–11]; in some of these works, for technical reasons, only the case of a translation invariant Hamiltonian, with $U = 0$, was considered). Note that this argument does not provide an effective estimate for the speed of convergence in (1.6).

Recently, a new proof of (1.6) has been presented by Fröhlich et al. in [12]. This method provides convergence estimates uniformly in Planck’s constant up to an exponentially small remainder, and therefore it can be used to combine the mean field- with the semiclassical limit (note that, in the semiclassical limit the Hartree equation converges to a Vlasov

equation). In [13] it has been shown that the limit (1.6) can be interpreted as a Egorov-type theorem, in the sense that the time evolution of observables commutes with their Wick quantization, up to corrections which vanish in the mean-field limit.

In the present work we propose yet a different approach. With respect to the techniques mentioned above, our approach has the advantage of being very simple and of giving a quantitative bounds of the order $1/N$ for the speed of convergence in (1.6). The estimates we obtain here are therefore better, in their N -dependence, than the bounds (1.7) derived, for factorized initial data, in [21]. However, our analysis is restricted to a smaller class of interaction potential V ; in fact, we need to assume that the operator norm $\|V\|$ is finite (the operator norm of the multiplication operator V equals the L^∞ -norm of the function $V(x)$), and that it has a Fourier transform $\widehat{V} \in L^1(\mathbb{R}^v)$. The main result of this paper is the following theorem.

Theorem 1.1 *Fix an arbitrary dimension $v \geq 1$. Assume that the one-particle potential U is such that $-\Delta + U$ is a self-adjoint operator on $L^2(\mathbb{R}^v)$. Suppose that $V \in L^\infty(\mathbb{R}^v)$ is such that $\|\widehat{V}\|_1 < \infty$. Denote by $\gamma_{N,t}^{(k)}$ the k -particle density associated with the solution $\psi_{N,t} = e^{-iH_N t} \psi_N$ of the Schrödinger equation (1.2) with Hamiltonian H_N given by (1.1) and with initial data $\psi_N = \varphi^{\otimes N}$, for some $\varphi \in H^1(\mathbb{R}^v)$ with $\|\varphi\|_2 = 1$. Then we have*

$$\text{Tr} |\gamma_{N,t}^{(k)} - |\varphi_t\rangle\langle\varphi_t|^{\otimes k}| \leq \frac{k^2 \lambda_V}{N} e^{2k\|V\|} (e^{8\|V\|t} - 1)$$

for every $t \in \mathbb{R}$, $N \geq 1$ and $1 \leq k \leq N$. Here we defined $\lambda_V = 1 + (\|\widehat{V}\|_1 / \|V\|)$ and φ_t is the solution to the nonlinear Hartree equation

$$i \partial_t \varphi_t = (-\Delta + U)\varphi_t + (V * |\varphi_t|^2)\varphi_t,$$

with $\varphi_{t=0} = \varphi$.

In order to prove Theorem 1.1 we control the growth of the correlations among the particles (at time $t = 0$, the wave function is factorized, and therefore there are no correlations). Our proof has been inspired by the techniques used in [20] to prove Lieb-Robinson inequalities for anharmonic lattice systems (see also the proof of Lieb-Robinson bounds for quantum spin systems given in [17–19]). To prove Lieb-Robinson bounds, one has to control the growth of the correlations between spatially separated observables. To prove Theorem 1.1, on the other hand, we have to control the growth of correlations between observables acting on different particles.

In Sect. 2, we prove that if A and B are two operators acting on different particles, then the commutator $[A, e^{iH_N t} B e^{-iH_N t}]$ remains of order $1/N$ for all fixed times (the commutator is zero at time $t = 0$); see Theorem 2.1. We apply this result to prove that, if $\gamma_{N,t}^{(k)}$ denote the marginal density associated with the evolution $\psi_{N,t} = e^{-iH_N t} \psi_N$, for a factorized initial data $\psi_N = \varphi^{\otimes N}$, then the difference $\gamma_{N,t}^{(m+n)} - \gamma_{N,t}^{(m)} \otimes \gamma_{N,t}^{(n)}$ remains of order $1/N$ for all fixed times (it is zero at time $t = 0$), when tested against product observables; see Proposition 2.2. Note that both Theorem 2.1 and Proposition 2.2 only require bounded potential to hold (the condition $\|\widehat{V}\|_1 < \infty$ is not required here). Finally, in Sect. 3, we apply the result of Proposition 2.2 to show Theorem 1.1.

Notation. For $k \geq 1$, denote by \mathcal{B}_k the algebra of bounded operators over $L^2(\mathbb{R}^{kv})$. For an operator $A \in \mathcal{B}_k$, and for integers $1 \leq i_1 < i_2 < \dots < i_k \leq N$, we denote by $A^{[i_1, \dots, i_k]}$ the operator on $L^2(\mathbb{R}^{Nv})$ acting as A over the particles i_1, \dots, i_k and as the identity over all other particles. We will use the notation $\|A\|$ to indicate the operator norm of A (where A

is meant as an operator from an L^2 space to itself). In particular $\|V\|$ denotes the operator norm of the potential (interpreted as a multiplication operator) which of course equals the L^∞ -norm of the function $V(x)$.

2 Growth of Correlations

Theorem 2.1 *Suppose that the external potential U is such that $-\Delta + U$ is self-adjoint, and assume that $V \in L^\infty(\mathbb{R}^v)$. Then, for every bounded m -particle observable $A \in \mathcal{B}_m$, and every bounded n -particle observable $B \in \mathcal{B}_n$, we have*

$$\| [A^{[i_1, \dots, i_m]}, e^{iH_N t} B^{[j_1, \dots, j_n]} e^{-iH_N t}] \| \leq \frac{mn \|A\| \|B\|}{N} (e^{4\|V\||t|} - 1) \tag{2.10}$$

for all integers $1 \leq i_1 < \dots < i_m \leq N$, $1 \leq j_1 < \dots < j_n \leq N$, with $\{i_1, \dots, i_m\} \cap \{j_1, \dots, j_n\} = \emptyset$.

Proof Without loss of generality, we assume that $t \geq 0$. For $n, m \in \mathbb{N}$, we define the quantity

$$f_{m,n}(t) = \sup_{A \in \mathcal{B}_m, B \in \mathcal{B}_n} \frac{\| [A^{[i_1, \dots, i_m]}, e^{iH_N t} B^{[j_1, \dots, j_n]} e^{-iH_N t}] \|}{\|A\| \|B\|} \tag{2.11}$$

for arbitrary integers $1 \leq i_1 < \dots < i_m \leq N$, and $1 \leq j_1 < \dots < j_n \leq N$ such that $\{i_1, \dots, i_m\} \cap \{j_1, \dots, j_n\} = \emptyset$. Note that, because of the permutation symmetry of H_N , the quantity $f(t)$ is independent of the choice of the indices i_ℓ, j_ℓ and

$$f_{m,n}(t) = \sup_{A \in \mathcal{B}_m, B \in \mathcal{B}_n} \frac{\| [A^{[n+1, \dots, n+m]}, e^{iH_N t} B^{[1, \dots, n]} e^{-iH_N t}] \|}{\|A\| \|B\|}.$$

We define next the modified Hamiltonian

$$H_N^{(n)} = H_N - \frac{1}{N} \sum_{\ell=1}^n \sum_{j>n} V(x_\ell - x_j).$$

With respect to the dynamics generated by $H_N^{(n)}$, particles $1, \dots, n$ are decoupled from the rest of the system. In particular this implies that $e^{-iH_N^{(n)} t} B^{[1, \dots, n]} e^{iH_N^{(n)} t}$ is still an operator acting only on the degrees of freedom of particles $1, \dots, n$, with norm equal to the norm of B . Therefore, we have

$$f_{m,n}(t) = \sup_{A \in \mathcal{B}_m, B \in \mathcal{B}_n} \frac{\| [A^{[n+1, \dots, n+m]}, e^{iH_N t} e^{-iH_N^{(n)} t} B^{[1, \dots, n]} e^{iH_N^{(n)} t} e^{-iH_N t}] \|}{\|A\| \|B\|}. \tag{2.12}$$

For given $A \in \mathcal{B}_m$ and $B \in \mathcal{B}_n$ we define the time-dependent bounded operator acting on $L^2(\mathbb{R}^{vN})$

$$g_{A,B}(t) = [A^{[n+1, \dots, n+m]}, e^{iH_N t} e^{-iH_N^{(n)} t} B^{[1, \dots, n]} e^{iH_N^{(n)} t} e^{-iH_N t}]$$

and we compute its time-derivative

$$\frac{d}{dt} g_{A,B}(t) = [A^{[n+1, \dots, n+m]}, e^{iH_N t} [i(H_N - H_N^{(n)}), e^{-iH_N^{(n)} t} B^{[1, \dots, n]} e^{iH_N^{(n)} t}] e^{-iH_N t}]$$

$$\begin{aligned}
 &= [A^{[n+1, \dots, n+m]}, [ie^{iH_N t} (H_N - H_N^{(n)})e^{-iH_N t}, e^{iH_N t} e^{-iH_N^{(n)} t} B^{([1, \dots, n])} e^{iH_N^{(n)} t} e^{-iH_N t}]] \\
 &= [ie^{iH_N t} (H_N - H_N^{(n)})e^{-iH_N t}, g_{A,B}(t)] \\
 &\quad + [e^{iH_N t} e^{-iH_N^{(n)} t} B^{[1, \dots, n]} e^{iH_N^{(n)} t} e^{-iH_N t}, \\
 &\quad [A^{[n+1, \dots, n+m]}, e^{iH_N t} (H_N - H_N^{(n)})e^{-iH_N t}]]
 \end{aligned}$$

where, in the last step we used the Jacobi identity. Next, we define

$$\mathcal{H}^{(n)}(t) = e^{iH_N t} (H_N - H_N^{(n)})e^{-iH_N t}.$$

It is simple to see that $\mathcal{H}^{(n)}(t)$ generates a two-parameter group of unitary transformations $\mathcal{U}^{(n)}(t, s)$ satisfying

$$i \partial_t \mathcal{U}^{(n)}(t, s) = \mathcal{H}^{(n)}(t) \mathcal{U}^{(n)}(t, s), \quad \text{with } \mathcal{U}^{(n)}(s, s) = 1 \quad \text{for all } s \in \mathbb{R}.$$

Therefore, we obtain

$$\begin{aligned}
 &\frac{d}{dt} \mathcal{U}^{(n)}(0, t) g_{A,B}(t) \mathcal{U}^{(n)}(t, 0) \\
 &= \mathcal{U}^{(n)}(0, t) [e^{iH_N t} e^{-iH_N^{(n)} t} B^{[1, \dots, n]} e^{iH_N^{(n)} t} e^{-iH_N t}, \\
 &\quad [A^{[n+1, \dots, n+m]}, e^{iH_N t} (H_N - H_N^{(n)})e^{-iH_N t}]] \mathcal{U}^{(n)}(t, 0). \tag{2.13}
 \end{aligned}$$

Integrating this identity from time 0 to time t , using that $g_{A,B}(0) = 0$ and the definition of $H_N^{(n)}$, we find that

$$\begin{aligned}
 g_{A,B}(t) &= \frac{1}{N} \sum_{\ell=1}^n \sum_{j>n} \int_0^t ds \mathcal{U}^{(n)}(t, s) \\
 &\quad \times [e^{iH_N s} e^{-iH_N^{(n)} s} B^{[1, \dots, n]} e^{iH_N^{(n)} s} e^{-iH_N s}, \\
 &\quad [A^{[n+1, \dots, n+m]}, e^{iH_N s} V(x_\ell - x_j) e^{-iH_N s}]] \mathcal{U}^{(n)}(s, t). \tag{2.14}
 \end{aligned}$$

Taking norms, we get

$$\begin{aligned}
 \|g_{A,B}(t)\| &\leq \frac{2}{N} \sum_{\ell=1}^n \sum_{j>n} \int_0^t ds \|B\| \| [A^{[n+1, \dots, n+m]}, e^{iH_N s} V(x_\ell - x_j) e^{-iH_N s}] \| \\
 &\leq \frac{4mnt \|A\| \|B\| \|V\|}{N} \\
 &\quad + 2n \|B\| \int_0^t ds \| [A^{[n+1, \dots, n+m]}, e^{iH_N s} V(x_1 - x_{n+m+1}) e^{-iH_N s}] \|
 \end{aligned}$$

where the first term on the r.h.s. corresponds to the terms with $j = n + 1, \dots, n + m$, while in the second term we used the permutation symmetry. From (2.11) and (2.12), it follows that

$$f_{m,n}(t) \leq \frac{4mnt \|V\|}{N} + 2n \|V\| \int_0^t ds f_{m,2}(s). \tag{2.15}$$

Iterating this equation k times, we obtain that

$$\begin{aligned}
 f_{m,n}(t) &\leq \frac{4mn\|V\|}{N}t + \frac{4mn\|V\|}{N} \sum_{r=1}^{k-1} (4\|V\|)^r \int_0^t ds_1 \cdots \int_0^{s_{r-1}} ds_r s_r \\
 &\quad + 2n\|V\|(4\|V\|)^{k-1} \int_0^t \cdots \int_0^{s_{k-1}} ds_k f_{m,2}(s_r).
 \end{aligned}
 \tag{2.16}$$

With the a-priori bound $f_{m,2}(s) \leq 2$, it follows that

$$f_{m,n}(t) \leq \frac{mn}{N} \sum_{r=0}^{k-1} \frac{(4\|V\|t)^{r+1}}{(r+1)!} + n \frac{(4\|V\|t)^k}{k!} \leq \frac{mn}{N} (e^{4\|V\|t} - 1) + n \frac{(4\|V\|t)^k}{k!}$$

for all $k \in \mathbb{N}$. Since the l.h.s. is independent of k , we obtain (2.10). □

The following proposition is a useful consequence of Theorem 2.1.

Proposition 2.2 *Suppose that U is such that $-\Delta + U$ is self-adjoint on $L^2(\mathbb{R}^\nu)$ and assume that $V \in L^\infty(\mathbb{R}^\nu)$. Denote $\psi_{N,t} = e^{-iH_{Nt}}\psi_N$ the solution to the N -particle Schrödinger equation with factorized initial data $\psi_N = \varphi^{\otimes N}$, for some $\varphi \in L^2(\mathbb{R}^\nu)$ with $\|\varphi\|_2 = 1$. Then, for any $A \in \mathcal{B}_m, B \in \mathcal{B}_n$, we have*

$$\begin{aligned}
 &|\langle \psi_{N,t}, A^{[i_1, \dots, i_m]} B^{[j_1, \dots, j_n]} \psi_{N,t} \rangle - \langle \psi_{N,t}, A^{[i_1, \dots, i_m]} \psi_{N,t} \rangle \langle \psi_{N,t}, B^{[j_1, \dots, j_n]} \psi_{N,t} \rangle| \\
 &\leq \frac{mn\|A\|\|B\|}{N} (e^{8\|V\|t} - 1)
 \end{aligned}
 \tag{2.17}$$

for arbitrary integers $1 \leq i_1 < \dots < i_m \leq N, 1 \leq j_1 < \dots < j_n \leq N$ with $\{i_1, \dots, i_m\} \cap \{j_1, \dots, j_n\} = \emptyset$. In particular, if $\gamma_{N,t}^{(k)}$ denotes the k -particle marginal associated with $\psi_{N,t}$, we have

$$|\text{Tr}(A^{[1, \dots, m]} \otimes B^{[m+1, \dots, m+n]})(\gamma_{N,t}^{(m+n)} - \gamma_{N,t}^{(m)} \otimes \gamma_{N,t}^{(n)})| \leq \frac{mn\|A\|\|B\|}{N} (e^{8\|V\|t} - 1).$$

Proof Because of the permutation symmetry we have

$$\begin{aligned}
 &\langle \psi_{N,t}, A^{[i_1, \dots, i_m]} B^{[j_1, \dots, j_n]} \psi_{N,t} \rangle - \langle \psi_{N,t}, A^{[i_1, \dots, i_m]} \psi_{N,t} \rangle \langle \psi_{N,t}, B^{[j_1, \dots, j_n]} \psi_{N,t} \rangle \\
 &= \langle \psi_{N,t}, A^{[1, \dots, m]} B^{[m+1, \dots, m+n]} \psi_{N,t} \rangle \\
 &\quad - \langle \psi_{N,t}, A^{[1, \dots, m]} \psi_{N,t} \rangle \langle \psi_{N,t}, B^{[m+1, \dots, m+n]} \psi_{N,t} \rangle.
 \end{aligned}
 \tag{2.18}$$

We observe that

$$\begin{aligned}
 &\langle \psi_{N,t}, A^{[1, \dots, m]} B^{[m+1, \dots, m+n]} \psi_{N,t} \rangle \\
 &= \langle \psi_N, e^{iH_{Nt}} A^{[1, \dots, m]} e^{-iH_{Nt}} e^{iH_{Nt}} B^{[m+1, \dots, m+n]} e^{-iH_{Nt}} \psi_N \rangle \\
 &= \langle \psi_N, e^{iH_{Nt}} A^{[1, \dots, m]} e^{-iH_{Nt}} |\varphi\rangle \langle \varphi|^{\otimes N} e^{iH_{Nt}} B^{[m+1, \dots, m+n]} e^{-iH_{Nt}} \psi_N \rangle \\
 &\quad + \sum_{j=1}^N \langle \psi_N, e^{iH_{Nt}} A^{[1, \dots, m]} e^{-iH_{Nt}} [|\varphi\rangle \langle \varphi|^{\otimes(j-1)} \otimes (1 - |\varphi\rangle \langle \varphi|) \otimes 1^{\otimes(N-j)}] \rangle
 \end{aligned}$$

$$\times e^{iH_N t} \mathbf{B}^{[m+1, \dots, m+n]} e^{-iH_N t} \psi_N \rangle. \tag{2.19}$$

Since $\psi_N = \varphi^{\otimes N}$, we get

$$\begin{aligned} & \langle \psi_{N,t}, A^{[1, \dots, m]} \mathbf{B}^{[m+1, \dots, m+n]} \psi_{N,t} \rangle \\ &= \langle \psi_{N,t}, A^{[1, \dots, m]} \psi_{N,t} \rangle \langle \psi_{N,t}, \mathbf{B}^{[m+1, \dots, m+n]} \psi_{N,t} \rangle \\ &+ \sum_{j=1}^m \langle \psi_N, e^{iH_N t} A^{[1, \dots, m]} e^{-iH_N t} (|\varphi\rangle\langle\varphi|^{\otimes(j-1)} \otimes \mathbf{1}^{\otimes(N-j+1)}) \\ &\times [(1 - |\varphi\rangle\langle\varphi|)^{[j]}, e^{iH_N t} \mathbf{B}^{[m+1, \dots, m+n]} e^{-iH_N t}] \psi_N \rangle \\ &+ \sum_{j=m+1}^{m+n} \langle \psi_N, [e^{iH_N t} A^{[1, \dots, m]} e^{-iH_N t}, (1 - |\varphi\rangle\langle\varphi|)^{[j]}] \\ &\times (|\varphi\rangle\langle\varphi|^{\otimes(j-1)} \otimes \mathbf{1}^{\otimes(N-j+1)}) e^{iH_N t} \mathbf{B}^{[m+1, \dots, m+n]} e^{-iH_N t} \psi_N \rangle \\ &+ \sum_{j=m+n+1}^N \langle \psi_N, [e^{iH_N t} A^{[1, \dots, m]} e^{-iH_N t}, (1 - |\varphi\rangle\langle\varphi|)^{[j]}] \\ &\times (|\varphi\rangle\langle\varphi|^{\otimes(j-1)} \otimes \mathbf{1}^{\otimes(N-j+1)}) [(1 - |\varphi\rangle\langle\varphi|)^{[j]}, e^{iH_N t} \mathbf{B}^{[m+1, \dots, m+n]} e^{-iH_N t}] \psi_N \rangle, \tag{2.20} \end{aligned}$$

where we used the notation $(1 - |\varphi\rangle\langle\varphi|)^{[j]} = \mathbf{1}^{(j-1)} \otimes (1 - |\varphi\rangle\langle\varphi|) \otimes \mathbf{1}^{(N-j-1)}$ for the operator acting as the projection $(1 - |\varphi\rangle\langle\varphi|)$ over the j -th particle, and as the identity over the other $(N - 1)$ particles. Note that, by definition $((1 - |\varphi\rangle\langle\varphi|)^{[j]})^2 = 1 - |\varphi\rangle\langle\varphi|$. From Theorem 2.1, we obtain

$$\begin{aligned} & |\langle \psi_{N,t}, A^{[1, \dots, m]} \mathbf{B}^{[m+1, \dots, m+n]} \psi_{N,t} \rangle - \langle \psi_{N,t}, A^{[1, \dots, m]} \psi_{N,t} \rangle \langle \psi_{N,t}, \mathbf{B}^{[m+1, \dots, m+n]} \psi_{N,t} \rangle| \\ &\leq \frac{2mn \|A\| \|B\|}{N} (e^{4\|V\|t} - 1) + \frac{mn \|A\| \|B\|}{N} (e^{4\|V\|t} - 1)^2 \\ &= \frac{mn \|A\| \|B\|}{N} (e^{8\|V\|t} - 1) \end{aligned}$$

which completes the proof of the proposition. □

3 Derivation of the Hartree Equation

Using the bounds on the correlations obtained in the previous section, we can now state and prove our main result.

Proof of Theorem 1.1 From the BBGKY hierarchy (1.8), we obtain the integral equation

$$\begin{aligned} \gamma_{N,t}^{(k)} &= \mathcal{U}^{(k)}(t) |\varphi\rangle\langle\varphi|^{\otimes k} - \frac{i}{N} \sum_{i < j}^k \int_0^t ds \mathcal{U}^{(k)}(t-s) [V(x_i - x_j), \gamma_{N,s}^{(k)}] \\ &- i \frac{(N-k)}{N} \sum_{j=1}^k \int_0^t ds \mathcal{U}^{(k)}(t-s) \text{Tr}_{k+1} [V(x_j - x_{k+1}), \gamma_{N,s}^{(k+1)}], \tag{3.21} \end{aligned}$$

where we defined the free evolution

$$\mathcal{U}^{(k)}(t)\gamma^{(k)} = e^{-i\sum_{j=1}^k(-\Delta_j+U(x_j))t} \gamma^{(k)} e^{i\sum_{j=1}^k(-\Delta_j+U(x_j))t},$$

and where Tr_{k+1} denotes the partial trace over the $(k + 1)$ -th particle. On the other hand, it is simple to check that, if φ_t is a solution to the nonlinear Hartree equation

$$i\partial_t\varphi_t = (-\Delta + U)\varphi_t + (V * |\varphi_t|^2)\varphi_t$$

then the orthogonal projection $|\varphi_t\rangle\langle\varphi_t|^{\otimes k}$ satisfies the equation

$$|\varphi_t\rangle\langle\varphi_t|^{\otimes k} = \mathcal{U}^{(k)}(t)|\varphi\rangle\langle\varphi|^{\otimes k} - i\sum_{j=1}^k\int_0^t ds\mathcal{U}^{(k)}(t-s)\text{Tr}_{k+1}[V(x_j-x_{k+1}),|\varphi_t\rangle\langle\varphi_t|^{\otimes(k+1)}].$$

Therefore, for an arbitrary k -particle observable $J^{(k)}$ over $L^2(\mathbb{R}^{kv})$, we obtain

$$\begin{aligned} &\text{Tr}J^{(k)}(\gamma_{N,t}^{(k)} - |\varphi_t\rangle\langle\varphi_t|^{\otimes k}) \\ &= -\frac{i}{N}\sum_{i<j}^k\int_0^t ds\text{Tr}(\mathcal{U}^{(k)}(s-t)J^{(k)})[V(x_i-x_j),\gamma_{N,s}^{(k)}] \\ &\quad + i\frac{k}{N}\sum_{j=1}^k\int_0^t ds\text{Tr}(\mathcal{U}^{(k)}(s-t)J^{(k)})[V(x_j-x_{k+1}),\gamma_{N,s}^{(k+1)}] \\ &\quad - i\sum_{j=1}^k\int_0^t ds\text{Tr}(\mathcal{U}^{(k)}(s-t)J^{(k)})[V(x_j-x_{k+1}),(\gamma_{N,s}^{(k+1)} - \gamma_{N,s}^{(k)} \otimes \gamma_{N,s}^{(1)})] \\ &\quad - i\sum_{j=1}^k\int_0^t ds\text{Tr}(\mathcal{U}^{(k)}(s-t)J^{(k)})[V(x_j-x_{k+1}),(\gamma_{N,s}^{(k)} - |\varphi_s\rangle\langle\varphi_s|^{\otimes k}) \otimes |\varphi_s\rangle\langle\varphi_s|] \\ &\quad - i\sum_{j=1}^k\int_0^t ds\text{Tr}(\mathcal{U}^{(k)}(s-t)J^{(k)})[V(x_j-x_{k+1}),\gamma_{N,s}^{(k)} \otimes (\gamma_{N,s}^{(1)} - |\varphi_s\rangle\langle\varphi_s|)]. \end{aligned} \tag{3.22}$$

We bound the fourth and fifth term using that

$$\begin{aligned} &|\text{Tr}(\mathcal{U}^{(k)}(s-t)J^{(k)})[V(x_j-x_{k+1}),(\gamma_{N,s}^{(k)} - |\varphi_s\rangle\langle\varphi_s|^{\otimes k}) \otimes |\varphi_s\rangle\langle\varphi_s|]| \\ &\leq 2\|V\|\|J^{(k)}\|\text{Tr}|\gamma_{N,s}^{(k)} - |\varphi_s\rangle\langle\varphi_s|^{\otimes k}| \end{aligned} \tag{3.23}$$

and that, analogously,

$$\begin{aligned} &|\text{Tr}(\mathcal{U}^{(k)}(s-t)J^{(k)})[V(x_j-x_{k+1}),\gamma_{N,s}^{(k)} \otimes (\gamma_{N,s}^{(1)} - |\varphi_s\rangle\langle\varphi_s|)]| \\ &\leq 2\|V\|\|J^{(k)}\|\text{Tr}|\gamma_{N,s}^{(1)} - |\varphi_s\rangle\langle\varphi_s|. \end{aligned} \tag{3.24}$$

As for the third term on the r.h.s. of (3.22), we expand the potential in a Fourier integral. We obtain

$$\begin{aligned}
 & |\text{Tr}(\mathcal{U}^{(k)}(s-t)J^{(k)})[V(x_j-x_{k+1}), (\gamma_{N,s}^{(k+1)} - \gamma_{N,s}^{(k)} \otimes \gamma_{N,s}^{(1)})]| \\
 & \leq \int \text{d}q |\widehat{V}(q)| |\text{Tr}(\mathcal{U}^{(k)}(s-t)J^{(k)})[e^{iq \cdot x_j} \otimes e^{-iq \cdot x_{k+1}}, (\gamma_{N,s}^{(k+1)} - \gamma_{N,s}^{(k)} \otimes \gamma_{N,s}^{(1)})]| \\
 & \leq \int \text{d}q |\widehat{V}(q)| |\text{Tr}(\mathcal{U}^{(k)}(s-t)J^{(k)})e^{iq \cdot x_j} \otimes e^{-iq \cdot x_{k+1}} (\gamma_{N,s}^{(k+1)} - \gamma_{N,s}^{(k)} \otimes \gamma_{N,s}^{(1)})| \\
 & \quad + \int \text{d}q |\widehat{V}(q)| |\text{Tr} e^{iq \cdot x_j} (\mathcal{U}^{(k)}(s-t)J^{(k)}) \otimes e^{-iq \cdot x_{k+1}} (\gamma_{N,s}^{(k+1)} - \gamma_{N,s}^{(k)} \otimes \gamma_{N,s}^{(1)})| \\
 & \leq \frac{2k \|\widehat{V}\|_1 \|J^{(k)}\|}{N} (e^{8\|V\|s} - 1)
 \end{aligned} \tag{3.25}$$

where we applied Proposition 2.2. Thus, from (3.22), we find that

$$\begin{aligned}
 & |\text{Tr} J^{(k)}(\gamma_{N,t}^{(k)} - |\varphi_t\rangle\langle\varphi_t|^{\otimes k})| \\
 & \leq 2k\|V\| \|J^{(k)}\| \int_0^t \text{d}s \text{Tr} |\gamma_{N,s}^{(k)} - |\varphi_s\rangle\langle\varphi_s|^{\otimes k}| + 2k\|V\| \|J^{(k)}\| \int_0^t \text{d}s \text{Tr} |\gamma_{N,s}^{(1)} - |\varphi_s\rangle\langle\varphi_s|| \\
 & \quad + \frac{k^2 \|\widehat{V}\|_1 \|J^{(k)}\|}{4\|V\|N} (e^{8\|V\|t} - 1) + \frac{3tk^2\|V\| \|J^{(k)}\|}{N}.
 \end{aligned} \tag{3.26}$$

For a trace class operator A over $L^2(\mathbb{R}^{\nu k})$, let $\|A\|_1 = \text{Tr}|A|$ be the trace norm of A . If we denote by \mathcal{K}_k the algebra of compact operators on $L^2(\mathbb{R}^{\nu k})$, we have the variational characterization of the trace norm

$$\|A\|_1 = \text{Tr}|A| = \sup_{J \in \mathcal{K}_k, \|J\|=1} |\text{Tr} JA|.$$

Therefore we find

$$\begin{aligned}
 \|\gamma_{N,t}^{(k)} - |\varphi_t\rangle\langle\varphi_t|^{\otimes k}\|_1 & \leq 2k\|V\| \int_0^t \text{d}s (\|\gamma_{N,s}^{(k)} - |\varphi_s\rangle\langle\varphi_s|^{\otimes k}\|_1 + \|\gamma_{N,s}^{(1)} - |\varphi_s\rangle\langle\varphi_s|\|_1) \\
 & \quad + \frac{k^2 \lambda_V}{2N} (e^{8\|V\|t} - 1)
 \end{aligned} \tag{3.27}$$

with $\lambda_V = 1 + (\|\widehat{V}\|_1/\|V\|)$. For $k = 1$, we obtain

$$\|\gamma_{N,t}^{(1)} - |\varphi_t\rangle\langle\varphi_t|\|_1 \leq 4\|V\| \int_0^t \text{d}s \|\gamma_{N,s}^{(1)} - |\varphi_s\rangle\langle\varphi_s|\|_1 + \frac{\lambda_V}{2N} (e^{8\|V\|t} - 1).$$

Iterating this inequality n times, it follows that

$$\begin{aligned}
 & \|\gamma_{N,t}^{(1)} - |\varphi_t\rangle\langle\varphi_t|\|_1 \\
 & \leq \frac{\lambda_V}{2N} (e^{8\|V\|t} - 1) + \frac{\lambda_V}{2N} \sum_{m=1}^{n-1} (4\|V\|)^m \int_0^t \text{d}s_1 \int_0^{s_1} \text{d}s_2 \cdots \int_0^{s_{m-1}} \text{d}s_m (e^{8\|V\|s_m} - 1) \\
 & \quad + (4\|V\|)^n \int_0^t \text{d}s_1 \cdots \int_0^{s_{n-1}} \text{d}s_n \|\gamma_{N,s_n}^{(1)} - |\varphi_{s_n}\rangle\langle\varphi_{s_n}|\|_1.
 \end{aligned} \tag{3.28}$$

Since

$$\int_0^t ds_1 \cdots \int_0^{s_{m-1}} ds_m e^{8\|V\|s_m} \leq \frac{(e^{8\|V\|t} - 1)}{(8\|V\|)^m}$$

and $\|\gamma_{N,s_n}^{(1)} - |\varphi_{s_n}\rangle\langle\varphi_{s_n}|\|_1 \leq 2$, we find

$$\|\gamma_{N,t}^{(1)} - |\varphi_t\rangle\langle\varphi_t|\|_1 \leq \frac{\lambda_V}{N}(e^{8\|V\|t} - 1). \tag{3.29}$$

Inserting this bound into the r.h.s. of (3.27), we conclude that

$$\|\gamma_{N,t}^{(k)} - |\varphi_t\rangle\langle\varphi_t|^{\otimes k}\|_1 \leq 2k\|V\| \int_0^t ds \|\gamma_{N,s}^{(k)} - |\varphi_s\rangle\langle\varphi_s|^{\otimes k}\|_1 + \frac{k^2\lambda_V}{N}(e^{8\|V\|t} - 1).$$

Iteration leads to

$$\begin{aligned} \|\gamma_{N,t}^{(k)} - |\varphi_t\rangle\langle\varphi_t|^{\otimes k}\|_1 &\leq \frac{k^2\lambda_V}{N}(e^{8\|V\|t} - 1) \\ &+ \frac{k^2\lambda_V}{N} \sum_{m=1}^{n-1} (2k\|V\|)^m \int_0^t ds_1 \int_0^{s_1} ds_2 \cdots \int_0^{s_{m-1}} ds_m (e^{8\|V\|s_m} - 1) \\ &+ (2k\|V\|)^n \int_0^t ds_1 \cdots \int_0^{s_{n-1}} ds_n \|\gamma_{N,s_n}^{(k)} - |\varphi_{s_n}\rangle\langle\varphi_{s_n}|^{\otimes k}\|_1 \end{aligned} \tag{3.30}$$

which implies that

$$\|\gamma_{N,t}^{(k)} - |\varphi_t\rangle\langle\varphi_t|^{\otimes k}\|_1 \leq \frac{k^2\lambda_V}{N} e^{2k\|V\|} (e^{8\|V\|t} - 1). \quad \square$$

References

1. Adami, R., Bardos, C., Golse, F., Teta, A.: Towards a rigorous derivation of the cubic nonlinear Schrödinger equation in dimension one. *Asymptot. Anal.* **40**(2), 93–108 (2004)
2. Adami, R., Golse, F., Teta, A.: Rigorous derivation of the cubic NLS in dimension one. *J. Stat. Phys.* **127**(6), 1193–1220 (2007)
3. Bardos, C., Golse, F., Mauser, N.: Weak coupling limit of the N -particle Schrödinger equation. *Methods Appl. Anal.* **7**, 275–293 (2000)
4. Elgart, A., Erdős, L., Schlein, B., Yau, H.-T.: Gross–Pitaevskii equation as the mean field limit of weakly coupled bosons. *Arch. Ration. Mech. Anal.* **179**(2), 265–283 (2006)
5. Elgart, A., Schlein, B.: Mean field dynamics of boson stars. *Commun. Pure Appl. Math.* **60**(4), 500–545 (2007)
6. Erdős, L., Schlein, B., Yau, H.-T.: Derivation of the Gross–Pitaevskii hierarchy for the dynamics of Bose–Einstein condensate. *Commun. Pure Appl. Math.* **59**(12), 1659–1741 (2006)
7. Erdős, L., Schlein, B., Yau, H.-T.: Derivation of the cubic non-linear Schrödinger equation from quantum dynamics of many-body systems. *Invent. Math.* **167**, 515–614 (2007)
8. Erdős, L., Schlein, B., Yau, H.-T.: Derivation of the Gross–Pitaevskii equation for the dynamics of Bose–Einstein condensate. *Ann. Math.* (to appear). Preprint [arXiv:math-ph/0606017](https://arxiv.org/abs/math-ph/0606017)
9. Erdős, L., Schlein, B., Yau, H.-T.: Rigorous derivation of the Gross–Pitaevskii equation. *Phys. Rev. Lett.* **98**(4), 040404 (2007)
10. Erdős, L., Schlein, B., Yau, H.-T.: Rigorous derivation of the Gross–Pitaevskii equation: the case of a strong potential. Preprint [arXiv:0802.3877](https://arxiv.org/abs/0802.3877)
11. Erdős, L., Yau, H.-T.: Derivation of the nonlinear Schrödinger equation from a many body Coulomb system. *Adv. Theor. Math. Phys.* **5**(6), 1169–1205 (2001)

12. Fröhlich, J., Graffi, S., Schwarz, S.: Mean-field- and classical limit of many body Schrödinger dynamics for bosons. *Commun. Math. Phys.* **271**(3), 681–697 (2007)
13. Fröhlich, J., Knowles, A., Pizzo, A.: Atomism and quantization. *J. Phys. A* **40**(12), 3033–3045 (2007)
14. Ginibre, J., Velo, G.: The classical field limit of scattering theory for non-relativistic many-boson systems, I. *Commun. Math. Phys.* **66**, 37–76 (1979)
15. Ginibre, J., Velo, G.: The classical field limit of scattering theory for non-relativistic many-boson systems, II. *Commun. Math. Phys.* **68**, 45–68 (1979)
16. Hepp, K.: The classical limit for quantum mechanical correlation functions. *Commun. Math. Phys.* **35**, 265–277 (1974)
17. Lieb, E.H., Robinson, D.W.: The finite group velocity of quantum spin systems. *Commun. Math. Phys.* **28**, 251–257 (1972)
18. Nachtergaele, B., Sims, R.: Lieb-Robinson bounds and the exponential clustering theorem. *Commun. Math. Phys.* **265**, 119–130 (2006)
19. Nachtergaele, B., Ogata, Y., Sims, R.: Propagation of correlations in quantum lattice systems. *J. Stat. Phys.* **124**(1), 1–13 (2006)
20. Nachtergaele, B., Raz, H., Schlein, B., Sims, R.: Lieb-Robinson bounds for harmonic and anharmonic lattice systems. Preprint [arXiv:0712.3820](https://arxiv.org/abs/0712.3820)
21. Rodnianski, I., Schlein, B.: Quantum fluctuations and rate of convergence towards mean field dynamics. Preprint [arXiv:0711.3087](https://arxiv.org/abs/0711.3087)
22. Spohn, H.: Kinetic equations from Hamiltonian dynamics. *Rev. Mod. Phys.* **52**(3), 569–615 (1980)