

From Discrete Velocity Boltzmann Equations to Gas Dynamics Before Shocks

Florent Berthelin · Athanasios E. Tzavaras · Alexis Vasseur

Received: 25 August 2008 / Accepted: 25 February 2009 / Published online: 14 March 2009
© Springer Science+Business Media, LLC 2009

Abstract This article is devoted to the proof of the hydrodynamic limit for a discrete velocity Boltzmann equation before appearance of shocks in the limit system.

Keywords Hydrodynamic limit · Entropy methods · Discrete Boltzmann equations · Gas dynamics

1 Introduction

We consider the system of discrete velocity Boltzmann equations

$$\partial_t f_i + v_i \partial_x f_i = \frac{1}{\varepsilon} Q_i(f, f), \quad \text{for } i = 1, \dots, N, \quad (1.1)$$

Partially supported by the National Science Foundation Grants DMS-0607053, DMS-0555272 and DMS-0757227, and by the program “Pythagoras” of the Greek Secretariat of Research.

F. Berthelin
Laboratoire J.A. Dieudonné, UMR 6621 CNRS, Université de Nice, Parc Valrose, 06108 Nice Cedex 2, France
e-mail: Florent.Berthelin@unice.fr

A.E. Tzavaras
Department of Mathematics, University of Maryland, College Park, 20742-4015, USA
e-mail: tzavaras@math.umd.edu

A.E. Tzavaras
Department of Applied Mathematics and Institute for Applied and Computational Mathematics (FORTH), University of Crete, 71409 Heraklion, Greece

A. Vasseur (✉)
Mathematics Department, University of Texas at Austin, 1 University Station C1200, Austin, TX 78712-0257, USA
e-mail: vasseur@math.utexas.edu

where

$$Q_i(f, f) = \sum_{jkl} S_{ijkl} (f_k f_l - f_i f_j), \quad (1.2)$$

and $N \geq 3$. Such systems have been extensively studied in the literature (see *e.g.* Cabannes, Gatignol and Luo [11] or Platkowski and Illner [24] and references therein) because they offer a simplification and approximation of the Boltzmann equation that shares remarkable similarities to the latter model. Nevertheless, these systems are quite simpler than the Boltzmann equation and for instance their existence theory is relatively well understood in both the cases of one dimension [1, 7, 19] as well as in several space dimensions [8, 15]. Discrete velocity models present certain pathologies in several space dimensions and we will refrain from working with them here.

The parameter ε is called mean free path or Knudsen number and, under certain conditions on the interaction coefficients S_{ijkl} that will be precised later, the system formally converges as $\varepsilon \rightarrow 0$ to equations,

$$\begin{aligned} \partial_t \rho + \partial_x(\rho u) &= 0, \\ \partial_t(\rho u) + \partial_x(\rho E) &= 0, \\ \partial_t(\rho E) + \partial_x(\rho J(u, E)) &= 0. \end{aligned} \quad (1.3)$$

The objective of this article is to establish the hydrodynamic limit from discrete Boltzmann equations (1.1) to the gas dynamics system in the form (1.3) in the regime where the solutions of (1.3) remain smooth.

It is well known that several models of fluid mechanics arise as limits of kinetic equations in a variety of scalings. The best known examples are the compressible Euler equations and the incompressible Navier-Stokes equations that can be formally derived from the Boltzmann equation, see *e.g.* the review article [16]. The mathematical justification of such scaling limits tests the limits of modern theories of weak solutions, with the most successful example being the derivation of incompressible Navier-Stokes equations from the Boltzmann equation, see [17] and references therein. In the hyperbolic scaling, the Boltzmann equation leads to the system of compressible Euler equations consisting of the balance of mass, momentum and energy. Due to difficulties with the theory of the Euler system, the validation of this limit in full generality is currently an open problem, with partial results achieved either when the limiting solution is smooth and the kinetic data are near a Maxwellian [29] or for solutions of the Euler system containing non-interacting shocks [32].

A simpler paradigm for studying the hyperbolic scaling is offered by discrete velocity problems (1.1). The formal limiting system then takes the form (1.3) which has certain similarity with the usual compressible Euler system (but also the nonphysical feature that the pressure and heat flux depend on the velocity). The usual approach for studying the Euler limit proceeds by studying spectral properties of the linearized Boltzmann operator, see the analysis of Calfisch and Papanicolaou [12] justifying the Euler limit from the Broadwell model to an isothermal Euler system consisting of balance of mass and momentum, and also the asymptotic in time convergence of (1.1) to global Maxwellians established in Kawashima [21]. We note that an analysis of the linearized Boltzmann operator lies in the core of the derivation of the Euler system from the Boltzmann equation for smooth solutions and near Maxwellian data [2, 29]. The novelty of this article is that we will use an energy method to establish this limit in the smooth regime, based on the natural entropy of the system (1.1) and outlined below. Especially, we do not need any regularity assumption on the

kinetic functions at ε fixed. Convergence in L^2 of the initial values are sufficient to ensure the convergence up to the blowing-up time of the limit equation.

The rigorous derivation of hydrodynamic (Euler) limits for weak solutions is far less understood (even in one-space dimension), due to the poor understanding of the existence theory for weak solutions of hyperbolic conservation laws. Nevertheless, there are available justifications of hydrodynamic limits from BGK-models to the isothermal Euler system in Lagrangian [26, 27] or Eulerian coordinates [3, 4] via compensated compactness. Moreover, the latest technology of the Bressan-Bianchini interaction functionals has been applied to justify the hydrodynamic limit from kinetic equations to systems of conservation laws [6]. All these analyses concern kinetic models with BGK-type collision operators and their extension to Boltzmann type collision operators appears nontrivial.

In this work, we establish the hydrodynamic limit using a relative entropy identity. The relative entropy method was initially developed in the context of uniqueness and stability for hyperbolic conservation laws by Dafermos [13] and DiPerna [14], and in a context of hydrodynamics for stochastic particle systems by Yau [31] and Olla-Varadhan-Yau [23]. It has been used in a variety of contexts to establish limits from entropy dissipating solutions of one physical theory to entropy conserving solutions of another. The reader is referred to [9, 10, 18, 20, 22, 25] for various applications. The approach used here is closest to the spirit of the articles Berthelin-Vasseur [5] and Tzavaras [28] concerning kinetic or relaxation limits for BGK-type of collision operators. For a Boltzmann-type collision operator one needs an estimation of the entropy dissipation (from below) in terms of a norm measuring the distance between the nonequilibrium kinetic density and the associated local Maxwellian. Villani in his well known study of the Cercignani conjecture [30] has studied the extent of validity of such inequalities in terms of the metric of the relative entropy. Instead, we estimate from below the entropy dissipation via an alternate metric (see Proposition 4.1). In addition, we use certain structural properties pertaining to the system (1.3) and due to the finite number of velocities. While the general form of the relative entropy identity remains valid for the case of the Boltzmann equation the latter two ingredients hinge on the fact that we deal with a discrete velocity model. Interestingly the resulting convergence result applies to the full compressible Euler system (1.3).

We remark that the analysis can be easily extended to the case of several space dimensions under the structural hypothesis (H). One could similarly establish a convergence result from a discrete velocity model equipped only with microscopic conservation of mass and momentum to an isothermal Euler system by replacing hypothesis (H) by the corresponding hypothesis that the collision kernel is two-dimensional and is spanned by the two associated collisional invariants.

We begin in Sect. 2 with a description of the model, an outline of the formalism of its hydrodynamic limit and the statement of the main result Theorem 2.3. In Sect. 3 we develop links between the kinetic and the macroscopic entropies and prove certain structural properties of the limit system, the entropy consistency property and hyperbolicity. Section 4 contains the key estimation of the entropy dissipation (Proposition 4.1), and Sect. 5 contains the derivation of the relative entropy identity and the conclusion of the proof of Theorem 2.3.

2 Description of the Model and Statement of Results

The interaction coefficients S_{ijkl} entering the definition of the collision operator (1.2) are assumed to satisfy the properties of symmetry and microreversibility,

$$S_{ijkl} = S_{jikl}, \quad S_{ijkl} = S_{ijlk}, \quad (2.4)$$

$$S_{ijkl} = S_{klij}, \tag{2.5}$$

and to describe the probability of the elastic collision $(i, j) \rightarrow (k, l)$ conserving the microscopic mass, momentum and energy

$$v_k + v_l = v_i + v_j, \quad v_k^2 + v_l^2 = v_i^2 + v_j^2 \quad \text{if } S_{ijkl} \neq 0. \tag{2.6}$$

For any $f \in \mathbb{R}^N$, we have from (2.6):

$$\sum_i Q_i = 0, \quad \sum_i v_i Q_i = 0, \quad \sum_i v_i^2 Q_i = 0, \tag{2.7}$$

which entail conservation laws for the total mass, momentum and energy. Consider the collision matrix $B \in \{-1, 0, 1\}^{N^2 \times N}$ given by:

$$B_{ij,i} = B_{ij,j} = -B_{ij,k} = -B_{ij,l} = 1 \quad \text{if } S_{ijkl} \neq 0, \\ B_{ij,k} = 0 \quad \text{everywhere else.}$$

Note that (2.7) implies that $(1, \dots, 1)$, (v_1, \dots, v_N) and (v_1^2, \dots, v_N^2) are in the kernel of B . We pose the additional hypothesis:

$$N(B) = \text{span} \{ (1, \dots, 1), (v_1, \dots, v_N), (v_1^2, \dots, v_N^2) \} \quad \text{and} \quad \dim N(B) = 3. \tag{H}$$

This implies that the only conserved quantities are precisely the mass, momentum and energy and that the model does not have any extraneous conservation laws. Finally, we define the entropy of f via the usual relation

$$\mathcal{H}(f) = \sum_{i=1}^N f_i \ln f_i.$$

For any C^1 function $F : \mathcal{D} \rightarrow \mathbb{R}$ defined on a convex set $\mathcal{D} \subset \mathbb{R}^k$, we define the associated relative function:

$$F(U_1|U_2) = F(U_1) - F(U_2) - F'(U_2)(U_1 - U_2).$$

We also set

$$s(y) = y \ln y,$$

and use in the sequel the following notations:

$$a * b = a_1 b_1 + a_2 b_2 + a_3 b_3 \quad a, b \in \mathbb{R}^3,$$

$$f \cdot g = \sum_{i=1}^N f_i g_i \quad f, g \in \mathbb{R}^N,$$

$$|f| = \sum_{i=1}^N |f_i| \quad f \in \mathbb{R}^N,$$

$$Pf = \sum_{i=1}^N (1, v_i, v_i^2) f_i \quad f \in \mathbb{R}^N,$$

$$|Pf| = \sum_{\beta=0}^2 \sum_{i=1}^N |v_i^\beta f_i| \quad f \in \mathbb{R}^N.$$

Our first lemma concerns the structure of Maxwellians associated to discrete velocity Boltzmann equations:

Lemma 2.1 *A vector $(M_1, \dots, M_N) \in (\mathbb{R}^+)^N$ verifies $Q(M, M) = 0$ if and only if there exists $a, b, c \in \mathbb{R}$ such that*

$$M_i = e^{a+bv_i+cv_i^2} \quad \text{for any } 1 \leq i \leq N.$$

Setting $\psi(b, c) = \sum_{i=1}^N e^{bv_i+cv_i^2}$, we express the Maxwellians in the form

$$M_i = \rho \frac{e^{bv_i+cv_i^2}}{\sum_{i=1}^N e^{bv_i+cv_i^2}}$$

and note the relations

$$\rho = \sum_{i=1}^N M_i = e^a \psi(b, c), \quad \sum_{i=1}^N v_i M_i = \rho \frac{\partial_b \psi}{\psi}(b, c), \quad \sum_{i=1}^N v_i^2 M_i = \rho \frac{\partial_c \psi}{\psi}(b, c). \tag{2.9}$$

For a given Maxwellian M , we define

$$(\rho, \rho u, \rho E) = PM,$$

that is to say

$$\begin{aligned} \rho &= \sum_{i=1}^N M_i = \sum_{i=1}^N e^{a+bv_i+cv_i^2} = e^a \psi(b, c), \\ \rho u &= \sum_{i=1}^N v_i M_i = \sum_{i=1}^N v_i e^{a+bv_i+cv_i^2} = \rho \frac{\partial_b \psi}{\psi}(b, c), \\ \rho E &= \sum_{i=1}^N v_i^2 M_i = \sum_{i=1}^N v_i^2 e^{a+bv_i+cv_i^2} = \rho \frac{\partial_c \psi}{\psi}(b, c). \end{aligned}$$

We notice that

$$u = \partial_b(\ln \psi), \quad E = \partial_c(\ln \psi),$$

and then we denote \mathcal{U} the set of admissible value of (u, E) , that is:

$$\mathcal{U} = \{(\partial_b(\ln \psi), \partial_c(\ln \psi)) \mid b, c \in \mathbb{R}\}.$$

The hydrodynamic limit system can be written formally as

$$\begin{cases} \partial_t \sum_i M_i + \partial_x \sum_i v_i M_i = 0, \\ \partial_t \sum_i v_i M_i + \partial_x \sum_i v_i^2 M_i = 0, \\ \partial_t \sum_i v_i^2 M_i + \partial_x \sum_i v_i^3 M_i = 0, \end{cases} \tag{2.10}$$

which leads to

$$\begin{cases} \partial_t \rho + \partial_x(\rho u) = 0, \\ \partial_t(\rho u) + \partial_x(\rho E) = 0, \\ \partial_t(\rho E) + \partial_x(\rho J) = 0, \end{cases} \tag{2.11}$$

where we set

$$J(u, E) = \frac{\partial_{bc} \psi}{\psi}. \tag{2.12}$$

The flux $J = J(u, E)$ is well defined thanks to the following lemma.

Lemma 2.2 *The function $\ln \psi$ is smooth and strictly convex and so the map: $T : (b, c) \rightarrow (u, E)$ defined by*

$$T(b, c) = \nabla_{(b,c)} \ln \psi(b, c)$$

is a C^1 diffeomorphism from \mathbb{R}^2 to \mathcal{U} .

We introduce also the entropy of the system:

$$\eta(\rho, \rho u, \rho E) = \mathcal{H}(M) = \sum_{i=1}^N M_i \ln M_i.$$

Conversely, for any $U = (\rho, \rho u, \rho E)$ with $\rho > 0$ and $(u, E) \in \mathcal{U}$, we define

$$M(U) = (M_i(U))_{i=1,\dots,N} = (e^{a+bv_i+cv_i^2})_{i=1,\dots,N}$$

with ρ, u, E and a, b, c related as in Lemma 2.1.

The article is devoted to the proof of the following theorem:

Theorem 2.3 *Let (ρ_0, u_0, E_0) , be a Lipschitzian function on \mathbb{R} with values in $\mathbb{R}^+ \times \mathcal{U}$ such that $U_0 = (\rho_0, \rho_0 u_0, \rho_0 E_0)$ and $\eta(\rho_0, \rho_0 u_0, \rho_0 E_0)$ lie altogether in $L^1(\mathbb{R})$ and $\partial_x U_0 \in L^2(\mathbb{R}) \cap L^\infty(\mathbb{R})$. Then, there exists a maximal time T^* such that the solution $(\rho, \rho u, \rho E)$ to the limit system (2.11) with initial values (ρ_0, u_0, E_0) stays Lipschitzian on $[0, T^*) \times \mathbb{R}$. Denote \bar{M} the Maxwellian associated to $(\rho, \rho u, \rho E)$. Consider $f_\varepsilon^0 \in (L^1(\mathbb{R}))^N$ such that each component is nonnegative and verifying $\mathcal{H}(f_\varepsilon^0)$ bounded in $L^1(\mathbb{R})$. We denote f_ε the solution of (1.1) with initial value f_ε^0 . If f_ε^0 converges strongly to \bar{M}^0 , Maxwellian associated to (ρ_0, u_0, E_0) in the sense that*

$$\int_{\mathbb{R}} \mathcal{H}(f_\varepsilon^0 | \bar{M}^0)(x) dx \xrightarrow{\varepsilon \rightarrow 0} 0,$$

then f_ε converges strongly to \bar{M} in the sense that for any $T < T^*$:

$$\sup_{0 \leq t \leq T} \int_{\mathbb{R}} \mathcal{H}(f_\varepsilon | \bar{M})(t, x) dx \xrightarrow{\varepsilon \rightarrow 0} 0,$$

where

$$\mathcal{H}(f|g) = \sum_i f_i \ln(f_i/g_i) - (f_i - g_i) \geq 0.$$

The proof is based on the results of Tzavaras [28] and Berthelin-Vasseur [5] on the relative entropy method, and on an estimation of the entropy-dissipation developed in Sect. 4.

3 Preliminaries

In this section, we gather certain structural properties of the model (1.1) and its hydrodynamic limit. Especially, we introduce a link between kinetic relative entropies and macroscopic ones, we show that the system is hyperbolic, entropy consistent and obtain properties on the domain \mathcal{U} .

First, we prove Lemmas 2.1 and 2.2.

Proof of Lemma 2.1 If $Q(M, M) = 0$ then in particular

$$D[M] = \sum_{i=1}^N \ln(M_i) Q_i(M) = \frac{1}{4} \sum_{ijkl} S_{ijkl} [\ln(M_k M_l) - \ln(M_i M_j)] (M_k M_l - M_i M_j) = 0.$$

But each terms of the last sum is nonnegative so, for any i, j, k, l such that $S_{ijkl} \neq 0$, we have $M_k M_l = M_i M_j$ which means:

$$\ln M_k + \ln M_l = \ln M_i + \ln M_j.$$

This implies that $\ln M = (\ln M_i)_{i=1, \dots, N}$ lies in $N(B)$. Hypothesis (H) implies that there exists (a, b, c) such that

$$\ln M_i = a + bv_i + cv_i^2 \quad \text{for any } 1 \leq i \leq N.$$

Conversely, note that if $\ln M$ is given by such a formula, then $M_k M_l = M_i M_j$ for any $ijkl$ verifying $S_{ijkl} \neq 0$ and $Q(M, M) = 0$ as well.

To show the second part of the statement. Note that from the definition of ψ we have $e^a = \frac{\rho}{\psi}$, that is to say $e^{-a} = \frac{\psi}{\rho}$. Noting that

$$\begin{aligned} \partial_b \psi &= \sum_{i=1}^N v_i e^{bv_i + cv_i^2} = e^{-a} \sum_{i=1}^N v_i M_i, \\ \partial_c \psi &= \sum_{i=1}^N v_i^2 e^{bv_i + cv_i^2} = e^{-a} \sum_{i=1}^N v_i^2 M_i, \end{aligned}$$

gives the result. □

We list the useful formulas

$$\begin{aligned} \sum_{i=1}^N e^{bv_i + cv_i^2} &= \psi(b, c), \\ \sum_{i=1}^N v_i e^{bv_i + cv_i^2} &= \psi(b, c)u, \end{aligned} \tag{3.1}$$

$$\sum_{i=1}^N v_i^2 e^{bv_i+cv_i^2} = \psi(b, c)E.$$

Proof of Lemma 2.2 The matrix of the second derivatives of $\ln \psi$ is:

$$\frac{1}{\psi^2} \begin{pmatrix} \psi \partial_{bb} \psi - (\partial_b \psi)^2 & \psi \partial_{bc} \psi - \partial_b \psi \partial_c \psi \\ \psi \partial_{bc} \psi - \partial_b \psi \partial_c \psi & \psi \partial_{cc} \psi - (\partial_c \psi)^2 \end{pmatrix}$$

which can be rewritten

$$\frac{1}{\psi} \begin{pmatrix} \sum_{i=1}^N (v_i - u)^2 e^{bv_i+cv_i^2} & \sum_{i=1}^N (v_i - u)(v_i^2 - E) e^{bv_i+cv_i^2} \\ \sum_{i=1}^N (v_i - u)(v_i^2 - E) e^{bv_i+cv_i^2} & \sum_{i=1}^N (v_i^2 - E)^2 e^{bv_i+cv_i^2} \end{pmatrix}.$$

Indeed, we have

$$\begin{aligned} \psi \partial_{bb} \psi - (\partial_b \psi)^2 &= \psi \sum_{i=1}^N v_i^2 e^{bv_i+cv_i^2} - \psi u \sum_{i=1}^N v_i e^{bv_i+cv_i^2} \\ &\stackrel{(3.12)}{=} \psi \sum_{i=1}^N (v_i - u)^2 e^{bv_i+cv_i^2}, \\ \psi \partial_{bc} \psi - (\partial_b \psi)(\partial_c \psi) &= \psi \sum_{i=1}^N v_i^3 e^{bv_i+cv_i^2} - \sum_{i=1}^N v_i e^{bv_i+cv_i^2} \sum_{i=1}^N v_i^2 e^{bv_i+cv_i^2} \\ &\stackrel{(3.1)}{=} \psi \sum_{i=1}^N v_i^3 e^{bv_i+cv_i^2} - \psi^2 u E \\ &\stackrel{(3.1)}{=} \psi \sum_{i=1}^N (v_i - u)(v_i^2 - E) e^{bv_i+cv_i^2}, \end{aligned}$$

and similarly for the last entry of the matrix.

The trace of this matrix is positive. Its determinant is also positive as can be seen by applying the Cauchy-Schwarz inequality,

$$\left[\sum_{i=1}^N (v_i - u)(v_i^2 - E) e^{bv_i+cv_i^2} \right]^2 \leq \left(\sum_{i=1}^N (v_i - u)^2 e^{bv_i+cv_i^2} \right) \left(\sum_{i=1}^N (v_i^2 - E)^2 e^{bv_i+cv_i^2} \right).$$

If the determinant is equal to 0 then equality holds in the Cauchy-Schwarz inequality which, in turn, implies that the vectors $(1, \dots, 1)$, (v_1, \dots, v_N) and (v_1^2, \dots, v_N^2) are linearly dependent. The latter is ruled out by hypothesis (H), and thus the matrix of the second derivatives of $\ln \psi$ is strictly positive, and $\ln \psi$ is strictly convex. The mapping T is a $C^{1,1}$ diffeomorphism from \mathbb{R}^2 to \mathcal{U} . □

We prove now the following lemma related to relative quantities.

Lemma 3.1 *Let $F : \mathcal{D} \rightarrow \mathbb{R}$ be a C^2 function on a convex set $\mathcal{D} \subset \mathbb{R}^k$. The function F is convex on \mathcal{D} if and only if the associated relative function is nonnegative on $\mathcal{D} \times \mathcal{D}$.*

Proof For any $U_1, U_2 \in \mathcal{V}$ we have:

$$F(U_1|U_2) = \int_0^1 \int_0^1 F''(U_1 + st(U_2 - U_1)) : [(U_1 - U_2) \otimes (U_1 - U_2)]t \, ds \, dt.$$

Hence, if F is convex then F'' is positive and so $F(U_1|U_2)$ is nonnegative. Conversely, for $|U_1 - U_2|$ small, we have:

$$F(U_1|U_2) = F''(U_2) : [(U_1 - U_2) \otimes (U_1 - U_2)] + o(|U_1 - U_2|^2).$$

Then, if $F(\cdot|\cdot)$ is nonnegative everywhere, then $F''(U_2)$ is a nonnegative matrix for any U_2 and F is convex. □

In particular, $s'(y) = 1 + \ln y$ leads to the usual relation:

$$s(y|z) = y \ln \frac{y}{z} - (y - z) \geq 0,$$

since s is convex.

Let us show now the following lemma which gives the link between the relative entropy at the kinetic level and at the macroscopic level.

Lemma 3.2 *For any $U = (\rho, \rho u, \rho E)$ with $\rho > 0$ and $(u, E) \in \mathcal{U}$, we set*

$$M(U) = (M_i(U))_{i=1,\dots,N} = (e^{a+bv_i+cv_i^2})_{i=1,\dots,N}$$

with ρ, u, E and a, b, c related as in Lemma 2.1. We then have

- (i) $PM(U) = U,$
- (ii) $\eta(U) = \mathcal{H}(M(U)) = \inf_{Pf=U} \mathcal{H}(f),$

and

- (iii) $\frac{\partial \eta}{\partial U}(U) * w = \frac{\partial \mathcal{H}}{\partial f}(M(U)) \cdot f,$ for any $w \in \mathbb{R}^3$ and any $f \in \mathbb{R}^N$ such that $w = Pf$.
Especially
- (iv) $\eta(U|\bar{U}) = \mathcal{H}(M(U)|M(\bar{U})),$

and

- (v) $\eta(U|\bar{U}) \leq \mathcal{H}(f|M(\bar{U})),$ for any $Pf = U.$

Proof

- (i) We have

$$PM(U) = \sum_{i=1}^N (1, v_i, v_i^2) M_i(U) = \sum_{i=1}^N (1, v_i, v_i^2) e^{a+bv_i+cv_i^2} = (\rho, \rho u, \rho E) = U.$$

(ii) By definition, we have

$$\eta(U) = \mathcal{H}(M(U)) = \sum_{i=1}^N s(M_i(U)).$$

For any f such that $Pf = U$, we have

$$\begin{aligned} 0 \leq \mathcal{H}(f|M(U)) &= \mathcal{H}(f) - \mathcal{H}(M(U)) - \partial_f \mathcal{H}(M(U)) \cdot (f - M(U)) \\ &= \mathcal{H}(f) - \mathcal{H}(M(U)) - \sum_{i=1}^N (1 + \ln M_i(U))(f_i - M_i(U)), \end{aligned}$$

with

$$\sum_{i=1}^N (1 + \ln M_i(U))(f_i - M_i(U)) = (1 + a, b, c) * P(f - M(U)) = 0$$

since $P(f - M(U)) = U - PM(U) = 0$. Hence:

$$\mathcal{H}(f) \geq \mathcal{H}(M(U)) \quad \text{for any } Pf = U,$$

which gives the result.

(iii) By differentiation of $U = PM(U)$ with respect to U , we get, with P linear,

$$Id = P \frac{\partial M}{\partial U}.$$

We denote

$$e_\beta = P \frac{\partial M}{\partial U_\beta}.$$

Let $f \in \mathbb{R}^N$ and $w \in \mathbb{R}^3$ such that $w = Pf$. Decomposing w on the basis (e_β) we have:

$$Pf = w = \sum_{\beta=1}^3 w_\beta e_\beta = \sum_{\beta=1}^3 w_\beta P \frac{\partial M}{\partial U_\beta},$$

and so

$$P \left(f - \sum_{\beta=1}^3 w_\beta \frac{\partial M}{\partial U_\beta} \right) = 0.$$

This gives the existence of g such that:

$$f = \sum_{\beta=1}^3 w_\beta \frac{\partial M}{\partial U_\beta} + g$$

$$Pg = 0.$$

But

$$\frac{\partial \eta}{\partial U_\beta} = \frac{\partial \mathcal{H}}{\partial f}(M(U)) \cdot \frac{\partial M}{\partial U_\beta}.$$

Hence:

$$\begin{aligned} \frac{\partial \eta}{\partial U} * w &= \sum_{\beta=1}^3 \frac{\partial \eta}{\partial U_{\beta}} w_{\beta} = \sum_{\beta=1}^3 w_{\beta} \frac{\partial \mathcal{H}}{\partial f}(M(U)) \cdot \frac{\partial M}{\partial U_{\beta}} \\ &= \frac{\partial \mathcal{H}}{\partial f}(M(U)) \cdot (f - g). \end{aligned}$$

We conclude with the argument that

$$\frac{\partial \mathcal{H}}{\partial f}(M(U)) \perp N(P).$$

This comes from the fact that

$$\eta(U) = \min_{Pf=U} \mathcal{H}(f) = \mathcal{H}(M(U))$$

(see [28, Proposition 2.1]).

(iv) We have

$$\begin{aligned} \eta(U|\bar{U}) &= \eta(U) - \eta(\bar{U}) - \partial_U \eta(\bar{U}) * (U - \bar{U}) \\ &= \mathcal{H}(M(U)) - \mathcal{H}(M(\bar{U})) - \partial_f \mathcal{H}(M(\bar{U})) \cdot (M(U) - M(\bar{U})) \\ &= \mathcal{H}(M(U)|M(\bar{U})) \end{aligned}$$

using (iii) with $w = U - \bar{U}$ and $f = M(U) - M(\bar{U})$.

(v) For f such that $U = Pf$, we have

$$\begin{aligned} \eta(U|\bar{U}) &= \mathcal{H}(M(U)) - \mathcal{H}(M(\bar{U})) - \partial_f \mathcal{H}(M(\bar{U})) \cdot (M(U) - M(\bar{U})) \\ &\leq \mathcal{H}(f) - \mathcal{H}(M(\bar{U})) - \partial_f \mathcal{H}(M(\bar{U})) \cdot (M(U) - M(\bar{U})) \\ &\leq \mathcal{H}(f|M(\bar{U})) - \partial_f H(M(\bar{U})) \cdot (M(U) - f). \end{aligned}$$

Now

$$\partial_f H(M(\bar{U})) \cdot g = \sum_{i=1}^N (1 + \ln M_i(U)) g_i = \sum_{i=1}^N (1 + a + bv_i + cv_i^2) g_i = 0$$

whenever $Pg = 0$. Since $P(M(U) - f) = 0$, we conclude. □

We can now show the main proposition of this section.

Proposition 3.3 *The system (1.3) is hyperbolic, admissible (in the sense of Berthelin-Vasseur [5]), i.e. there exists $C > 0$ such that*

$$|A(U|\bar{U})| \leq C\eta(U|\bar{U}) \quad \text{for any } \rho > 0, (u, E) \in \mathcal{U},$$

η is a convex entropy and

$$\eta(U|\bar{U}) = \mathcal{H}(M(U)|M(\bar{U})) = s(\rho|\bar{\rho}) + \rho \ln \psi((\bar{b}, \bar{c})|(b, c))$$

for any U, \bar{U} with ρ, u, E and a, b, c related as in Lemma 2.1. Finally, there exists a constant $C > 0$ such that

$$|u| + |E| \leq C \quad \text{for any } (u, E) \in \mathcal{U}.$$

Proof Let us first check that η is an entropy of the limit system with entropy flux $\sum_{i=1}^N v_i M_i(U) \ln M_i(U)$. Indeed

$$\begin{aligned} & \partial_t \sum_{i=1}^N M_i \ln M_i + \partial_x \sum_{i=1}^N v_i M_i \ln M_i \\ &= \sum_{i=1}^N (1 + \ln M_i)(\partial_t M_i + v_i \partial_x M_i) \\ &= \sum_{i=1}^N (1 + a + b v_i + c v_i^2)(\partial_t M_i + v_i \partial_x M_i) \\ &= (1 + a) \left(\partial_t \sum_{i=1}^N M_i + \partial_x \sum_{i=1}^N v_i M_i \right) + b \left(\partial_t \sum_{i=1}^N v_i M_i + \partial_x \sum_{i=1}^N v_i^2 M_i \right) \\ & \quad + c \left(\partial_t \sum_{i=1}^N v_i^2 M_i + \partial_x \sum_{i=1}^N v_i^3 M_i \right) \\ &= 0. \end{aligned}$$

Let us now calculate $\mathcal{H}(M|\bar{M})$ for two Maxwellians M, \bar{M} . We set

$$\begin{aligned} \ln M_i &= a + b v_i + c v_i^2 \\ \ln \bar{M}_i &= \bar{a} + \bar{b} v_i + \bar{c} v_i^2. \end{aligned}$$

Then,

$$\begin{aligned} \mathcal{H}(M|\bar{M}) &= \mathcal{H}(M) - \mathcal{H}(\bar{M}) - \partial_f \mathcal{H}(\bar{M}) \cdot (M - \bar{M}) \\ &= \sum_{i=1}^N M_i \ln M_i - \sum_{i=1}^N \bar{M}_i \ln \bar{M}_i - \sum_{i=1}^N (1 + \ln \bar{M}_i) \cdot (M_i - \bar{M}_i) \\ &= \sum_{i=1}^N M_i (\ln M_i - \ln \bar{M}_i) - \sum_{i=1}^N (M_i - \bar{M}_i) \\ &= (a - \bar{a}) \sum_{i=1}^N M_i + (b - \bar{b}) \sum_{i=1}^N v_i M_i + (c - \bar{c}) \sum_{i=1}^N v_i^2 M_i - \sum_{i=1}^N (M_i - \bar{M}_i) \\ &= (a - \bar{a})\rho + (b - \bar{b})\rho u + (c - \bar{c})\rho E - (\rho - \bar{\rho}) \\ &= \rho(\ln(\rho/\psi) - \ln(\bar{\rho}/\bar{\psi})) + (b - \bar{b})\rho u + (c - \bar{c})\rho E - (\rho - \bar{\rho}) \\ &= s(\rho|\bar{\rho}) + \rho [\ln \bar{\psi} - \ln \psi - \partial_b(\ln \psi)(b - \bar{b}) - \partial_c(\ln \psi)(c - \bar{c})] \\ &= s(\rho|\bar{\rho}) + \rho(\ln \psi)((\bar{b}, \bar{c})|(b, c)). \end{aligned}$$

The function s and $(-\ln \psi)$ are convex, and thus, thanks to Lemma 3.1,

$$\mathcal{H}(M|\overline{M}) \geq 0 \quad \text{for any } M, \overline{M}.$$

Lemma 3.2 gives that the relative entropy of η is nonnegative, and thanks to Lemma 3.1 again, we conclude that η is convex. Hence, the limit system is hyperbolic.

Note that for any $(u, E) \in \mathcal{U}$, since

$$u = \frac{\sum_{i=1}^N v_i M_i}{\sum_{i=1}^N M_i}, \quad E = \frac{\sum_{i=1}^N v_i^2 M_i}{\sum_{i=1}^N M_i},$$

we have

$$|u| \leq \sup_{i=1, \dots, N} |v_i|, \quad |E| \leq \sup_{i=1, \dots, N} |v_i^2|.$$

Hence \mathcal{U} is bounded in \mathbb{R}^2 . Let us write the limit system as

$$\partial_t U + \partial_x A(U) = 0,$$

where

$$A(\rho, \rho u, \rho E) = (\rho u, \rho E, \rho J(u, E)).$$

First note that the two first component of A are linear in U , so the associated relative quantity are 0. For the third one we calculate:

$$A_3(U|\overline{U}) = \rho J((u, E)|(\overline{u}, \overline{E})).$$

Thanks to the Taylor expansion, since $J \in C^2$ and \mathcal{U} is bounded, there exists a constant $C > 0$ such that for any $(u, E) \in \mathcal{U}$, we have

$$J((u, E)|(\overline{u}, \overline{E})) \leq C(|u - \overline{u}|^2 + |E - \overline{E}|^2).$$

We also have

$$\eta(U|\overline{U}) \geq \rho(\ln \psi)((\overline{b}, \overline{c})|(b, c)) \geq c\rho(|u - \overline{u}|^2 + |E - \overline{E}|^2),$$

with $c > 0$ thanks to the strict convexity of $\ln \psi$ and the boundedness of \mathcal{U} . Hence

$$|A(U|\overline{U})| \leq \frac{C}{c} \eta(U|\overline{U}) \quad \text{for any } \rho > 0, (u, E) \in \mathcal{U},$$

which means that the system is admissible. □

4 Estimation of the Dissipation

This section is dedicated to the estimation of the dissipation

$$D(f) = \frac{1}{4} \sum_{ijkl} S_{ijkl} \ln \left(\frac{f_k f_l}{f_i f_j} \right) (f_k f_l - f_i f_j) \geq 0 \tag{4.1}$$

via the proposition:

Proposition 4.1 *There exists a constant C such that for any $f \in \mathbb{R}^N$ we have*

$$\sum_{i=1}^N |f_i - M_i| \leq C\sqrt{D(f)}, \tag{4.2}$$

where $M = M(Pf)$ is the associated Maxwellian.

We first prove three lemmas.

Lemma 4.2 *Let $0 < \alpha < \beta$. For any $f \in \mathbb{R}^N$, we set $\rho = \sum_{i=1}^N f_i$, and $M = M(Pf)$. There exists $C_{\alpha\beta}$ such that for any $f \in \mathbb{R}^N$, if $0 < \alpha\rho \leq f_i \leq \beta\rho$ for any i , then*

$$D(f) \geq C_{\alpha\beta} \sum_{i=1}^N |f_i - M_i|^2. \tag{4.3}$$

Proof Since $D(f/\rho) = D(f)/\rho^2$ and $\sum_{i=1}^N |\frac{f_i}{\rho} - M_i(P(\frac{f}{\rho}))|^2 = \sum_{i=1}^N |\frac{f_i}{\rho} - \frac{M_i(Pf)}{\rho}|^2 = \frac{1}{\rho^2} \sum_{i=1}^N |f_i - M_i|^2$, we can assume that $\rho = 1$.

From $|\ln \frac{A}{B}| \leq \max(\frac{1}{A}, \frac{1}{B})|A - B|$ with $A = f_i f_j$ and $B = f_k f_l$, we get $D(f) \geq \overline{D}(f)$ with

$$\overline{D}(f) = \frac{\alpha^2}{4} \sum_{ijkl} S_{ijkl} (\ln f_k + \ln f_l - \ln f_i - \ln f_j)^2.$$

Since the kernel of $\overline{D}(f)$ is $V = \text{vect}((1, \dots, 1), (v_1, \dots, v_N), (v_1^2, \dots, v_N^2))$ from property (H), denoting by \mathbb{P} the linear projection from \mathbb{R}^N onto V , there exists C such that

$$\overline{D}(f) \geq C \sum_i |\ln f_i - \mathbb{P}(\ln f_i)|^2.$$

Now, since $\exp(\mathbb{P}(\ln f)) = M(\exp(\mathbb{P}(\ln f)))$, we have

$$\begin{aligned} f - M(f) &= \exp(\ln f) - \exp(\mathbb{P}(\ln f)) + M(\exp(\mathbb{P}(\ln f))) - M(\exp(\ln f)) \\ &= (\text{Id} - M) \circ \exp(\ln f) - (\text{Id} - M) \circ \exp(\mathbb{P}(\ln f)). \end{aligned}$$

Using that \exp is Lipschitz on every $]-\infty, R]$ and that $\mathbb{P} \ln f$ do not goes to $-\infty$, there exists $K_{\alpha\beta} > 0$ such that $(\text{Id} - M) \circ \exp$ is Lipschitz on $\ln[\alpha, \beta]$ and on $\ln(\mathbb{P}[\alpha, \beta])$. Thus

$$|f_i - M_i(f)| \leq K_{\alpha\beta} |\ln f_i - \mathbb{P} \ln f_i|$$

and therefore

$$\overline{D}(f) \geq \frac{C}{K_{\alpha\beta}^2} \sum_i |f_i - M_i|^2. \quad \square$$

Lemma 4.3 *There exists γ_1, C_1 such that for any $f \in \mathbb{R}^N$, setting $\rho = \sum_{i=1}^N f_i$, if there exists i_0 such that $f_{i_0} \leq \gamma_1 \rho$, then*

$$D(f) \geq C_1 \rho^2. \tag{4.4}$$

Proof Since $D(f/\rho) = D(f)/\rho^2$, we may assume with no loss of generality that $\rho = 1$.

The proof proceeds by contradiction. Let us assume that for any γ, C , there exists f and i_0 such that $f_{i_0} \leq \gamma\rho$ and $D(f) \leq C$. From Proposition 3.3, \mathcal{U} is bounded, that is to say $(u = \sum_{i=1}^N v_i f_i, E = \sum_{i=1}^N |v_i|^2 f_i)$ is bounded. Thus there exists γ such that $0 < \gamma < M_{i_0}$ for any i .

With this γ , for any $n \in \mathbb{N}^*$, taking $C = 1/n$, there exists f^n and $i_0(n)$ such that $f_{i_0(n)}^n \leq \gamma$ and $D(f^n) \leq 1/n$. Since $i_0(n)$ takes finitely many values, we can extract a subsequence such that $i_0(n)$ remains constant. For this index i_0 , we have for a subsequence $f_{i_0}^n \rightarrow f_{i_0} \in [0, \gamma]$, and extracting successively further subsequences, $f_j^n \rightarrow f_j \in [0, 1]$ for all other j . Now $D(f^n) \rightarrow 0$ gives $D(f) = 0$, and Lemma 2.1 implies that $f = M$ and then $\gamma < M_{i_0} = f_{i_0}$ which is a contradiction. \square

By similar arguments, we also prove that

Lemma 4.4 *There exists γ_2, C_2 such that for any $f \in \mathbb{R}^N$, setting $\rho = \sum_{i=1}^N f_i$, if there exists i_0 such that $f_{i_0} \geq \gamma_2\rho$, then*

$$D(f) \geq C_2\rho^2. \tag{4.5}$$

Based on these three properties, we can now show the Proposition 4.1.

Proof of Proposition 4.1. Let $\varepsilon > 0$ and set $I = \sum_{i=1}^N |f_i - M_i|$. If $\rho < \varepsilon$ then $I \leq 2\varepsilon$.

For $\rho \geq \varepsilon$, we select γ_1, γ_2 as in Lemmas 4.3 and 4.4 and distinguish three possibilities: either (i) $\gamma_1\rho < f_i < \gamma_2\rho$ for all indices i , or (ii) there exists i_0 so that $f_{i_0} > \gamma_2\rho$, or finally (iii) there is i_0 such that $f_{i_0} < \gamma_1\rho$. In each case I is estimated as follows:

$$\begin{aligned} \sum_{i=1}^N |f_i - M_i| &\leq \sum_{i=1}^N |f_i - M_i| \mathbb{1}_{\rho \leq \varepsilon} + \sum_{i=1}^N |f_i - M_i| \mathbb{1}_{\rho \geq \varepsilon} \\ &\leq 2\varepsilon + \sum_{i=1}^N |f_i - M_i| \mathbb{1}_{\exists i_0; f_{i_0} \leq \gamma_1\rho} \mathbb{1}_{\rho \geq \varepsilon} \\ &\quad + \sum_{i=1}^N |f_i - M_i| \mathbb{1}_{\exists i_0; f_{i_0} \geq \gamma_2\rho} \mathbb{1}_{\rho \geq \varepsilon} \\ &\quad + \sum_{i=1}^N |f_i - M_i| \mathbb{1}_{\forall i; \gamma_1\rho \leq f_i \leq \gamma_2\rho} \mathbb{1}_{\rho \geq \varepsilon} \\ &\leq 2\varepsilon + 2\rho \mathbb{1}_{\exists i_0; f_{i_0} \leq \gamma_1\rho} + 2\rho \mathbb{1}_{\exists i_0; f_{i_0} \geq \gamma_2\rho} \\ &\quad + \sqrt{\sum_i |f_i - M_i|^2 \mathbb{1}_{\forall i; \gamma_1\rho \leq f_i \leq \gamma_2\rho} \mathbb{1}_{\rho \geq \varepsilon}} \sqrt{N} \\ &\leq 2\varepsilon + 2\sqrt{\frac{D(f)}{C_1}} + 2\sqrt{\frac{D(f)}{C_2}} + \sqrt{\frac{ND(f)}{C_{\gamma_1\gamma_2}}} \end{aligned}$$

Finally, we take $\varepsilon \rightarrow 0$. \square

5 Hydrodynamic Limit

In this section, we prove Theorem 2.3. We denote by f_ε the solution of (1.1), by $U_\varepsilon = (\rho_\varepsilon, \rho_\varepsilon u_\varepsilon, \rho_\varepsilon E_\varepsilon) = P f_\varepsilon = \sum_{i=1}^N (1, v_i, v_i^2)(f_\varepsilon)_i$, by $M_\varepsilon = M(U_\varepsilon)$, by \bar{U} the smooth solution to the limit system and by $\bar{M} = \bar{M}(\bar{U})$ the associated Maxwellian.

Multiplying (1.1) by $\ln(f_\varepsilon)_i$ and summing in i gives

$$\partial_t \sum_{i=1}^N (f_\varepsilon)_i \ln(f_\varepsilon)_i + \partial_x \sum_{i=1}^N v_i (f_\varepsilon)_i \ln(f_\varepsilon)_i + \frac{D(f_\varepsilon)}{\varepsilon} = 0. \tag{5.1}$$

Thanks to Proposition 3.3, we have:

$$\partial_t \sum_{i=1}^N \bar{M}_i \ln \bar{M}_i + \partial_x \sum_{i=1}^N v_i \bar{M}_i \ln \bar{M}_i = 0. \tag{5.2}$$

We can now study the evolution of the relative entropy between f_ε and \bar{M} :

$$\begin{aligned} & \partial_t \mathcal{H}(f_\varepsilon | \bar{M}) + \partial_x \sum_{i=1}^N v_i s((f_\varepsilon)_i | \bar{M}_i) \\ &= \partial_t \sum_{i=1}^N s((f_\varepsilon)_i | \bar{M}_i) + \partial_x \sum_{i=1}^N v_i s((f_\varepsilon)_i | \bar{M}_i) \\ &= \partial_t \sum_{i=1}^N (f_\varepsilon)_i \ln(f_\varepsilon)_i + \partial_x \sum_{i=1}^N v_i (f_\varepsilon)_i \ln(f_\varepsilon)_i \\ & \quad - \partial_t \sum_{i=1}^N \bar{M}_i \ln \bar{M}_i - \partial_x \sum_{i=1}^N v_i \bar{M}_i \ln \bar{M}_i \\ & \quad - \partial_t \sum_{i=1}^N (1 + \ln \bar{M}_i)((f_\varepsilon)_i - \bar{M}_i) - \partial_x \sum_{i=1}^N v_i (1 + \ln \bar{M}_i)((f_\varepsilon)_i - \bar{M}_i). \end{aligned}$$

Since

$$\sum_{i=1}^N (1 + \ln \bar{M}_i)((f_\varepsilon)_i - \bar{M}_i) = \partial_f \mathcal{H}(\bar{M}) \cdot (f_\varepsilon - \bar{M}),$$

and using the notation $V : \mathbb{R}^N \rightarrow \mathbb{R}^N$ defined by:

$$V f_i = v_i f_i \quad 1 \leq i \leq N,$$

we also have

$$\sum_{i=1}^N v_i (1 + \ln \bar{M}_i)((f_\varepsilon)_i - \bar{M}_i) = \partial_f \mathcal{H}(\bar{M}) \cdot (V f_\varepsilon - V \bar{M}).$$

Combining this with (5.1) and (5.2), we get

$$\begin{aligned} & \partial_t \mathcal{H}(f_\varepsilon | \bar{M}) + \partial_x \sum_{i=1}^N v_i s((f_\varepsilon)_i | \bar{M}_i) + \frac{D(f_\varepsilon)}{\varepsilon} \\ &= -\partial_t \left(\partial_f \mathcal{H}(\bar{M}) \cdot (f_\varepsilon - \bar{M}) \right) - \partial_x \left(\partial_f \mathcal{H}(\bar{M}) \cdot (V f_\varepsilon - V \bar{M}) \right). \end{aligned}$$

Using Lemma 3.2, we get

$$\begin{aligned} & \partial_t \mathcal{H}(f_\varepsilon | \bar{M}) + \partial_x \sum_{i=1}^N v_i s((f_\varepsilon)_i | \bar{M}_i) + \frac{D(f_\varepsilon)}{\varepsilon} \tag{5.3} \\ &= -\partial_t \left(\partial_U \eta(\bar{U}) * P(f_\varepsilon - \bar{M}) \right) - \partial_x \left(\partial_U \eta(\bar{U}) * P(V f_\varepsilon - V \bar{M}) \right) \\ &= -\partial_t (\partial_U \eta(\bar{U})) * P(f_\varepsilon - \bar{M}) - \partial_U \eta(\bar{U}) * \partial_t (P(f_\varepsilon - \bar{M})) \\ &\quad - \partial_x (\partial_U \eta(\bar{U})) * P(V f_\varepsilon - V \bar{M}) - \partial_U \eta(\bar{U}) * \partial_x (P(V f_\varepsilon - V \bar{M})). \end{aligned}$$

For $k = 0, 1, 2$, multiplying (1.1) by v_i^k , summing over i and using (2.7), we have

$$\partial_t \sum_{i=1}^N v_i^k (f_\varepsilon)_i + \partial_x \sum_{i=1}^N v_i^{k+1} (f_\varepsilon)_i = 0,$$

that is to say

$$\partial_t P f_\varepsilon + \partial_x P (V f_\varepsilon) = 0. \tag{5.4}$$

Furthermore,

$$\partial_t \sum_{i=1}^N v_i^k \bar{M}_i + \partial_x \sum_{i=1}^N v_i^{k+1} \bar{M}_i = 0,$$

that is to say

$$\partial_t P(\bar{M}) + \partial_x P(V \bar{M}) = 0. \tag{5.5}$$

It gives

$$\begin{aligned} & \partial_t \mathcal{H}(f_\varepsilon | \bar{M}) + \partial_x \sum_{i=1}^N v_i s((f_\varepsilon)_i | \bar{M}_i) + \frac{D(f_\varepsilon)}{\varepsilon} \\ &= -\partial_{UU}^2 \eta(\bar{U}) \partial_t(\bar{U}) * P(f_\varepsilon - \bar{M}) - \partial_{UU}^2 \eta(\bar{U}) \partial_x(\bar{U}) * P(V f_\varepsilon - V \bar{M}) \\ &= \partial_{UU}^2 \eta(\bar{U}) A'(\bar{U}) \partial_x(\bar{U}) * P(f_\varepsilon - \bar{M}) - \partial_{UU}^2 \eta(\bar{U}) \partial_x(\bar{U}) * P(V f_\varepsilon - V \bar{M}) \\ &= \partial_{UU}^2 \eta(\bar{U}) \partial_x(\bar{U}) * \left(A'(\bar{U})(U_\varepsilon - \bar{U}) - P(V f_\varepsilon - V \bar{M}) \right) \\ &= \partial_{UU}^2 \eta(\bar{U}) \partial_x(\bar{U}) * \left(A'(\bar{U})(U_\varepsilon - \bar{U}) - P(V f_\varepsilon - V M_\varepsilon) - P(V M_\varepsilon - V \bar{M}) \right), \end{aligned}$$

where we used the fact that, since $\eta(U)$ is an entropy for (2.11), the flux $A(U)$ satisfies $(\partial_{uu}\eta)A' = (A')^T \partial_{uu}\eta$. Now

$$P(VM_\varepsilon - V\bar{M}) = \sum_{i=1}^N (1, v_i, v_i^2)v_i((M_\varepsilon)_i - \bar{M}_i) = A(U_\varepsilon) - A(\bar{M}),$$

therefore

$$\begin{aligned} \partial_t \mathcal{H}(f_\varepsilon | \bar{M}) + \partial_x \sum_{i=1}^N v_i s((f_\varepsilon)_i | \bar{M}_i) + \frac{D(f_\varepsilon)}{\varepsilon} \\ = -\partial_{UV}^2 \eta(\bar{U}) \partial_x(\bar{U}) * (A(U_\varepsilon | \bar{U}) + P(Vf_\varepsilon - VM_\varepsilon)). \end{aligned} \tag{5.6}$$

We exploit this evolution equation in order to get the bound. First we want to bound $D(f_\varepsilon)$ with respect to ε . Integrating (5.3) with respect to (t, x) gives

$$\begin{aligned} \int_{\mathbb{R}} \mathcal{H}(f_\varepsilon | \bar{M})(t, x) dx - \int_{\mathbb{R}} \mathcal{H}(f_\varepsilon^0 | \bar{M}^0)(x) dx + \int_0^t \int_{\mathbb{R}} \frac{D(f_\varepsilon)}{\varepsilon} dx ds \\ = - \int_{\mathbb{R}} \partial_U \eta(\bar{U}) * P(f_\varepsilon - \bar{M}) dx + \int_{\mathbb{R}} \partial_U \eta(\bar{U}) * P(f_\varepsilon^0 - \bar{M}^0) dx. \end{aligned}$$

For every $T < T^*$, there exists C_T such that $|\partial_U \eta(\bar{U})|(t, x) \leq C_T$ for any $x \in \mathbb{R}, 0 \leq t \leq T$. Thus we have

$$\int_0^t \int_{\mathbb{R}} \frac{D(f_\varepsilon)}{\varepsilon} dx ds \leq \int_{\mathbb{R}} \mathcal{H}(f_\varepsilon^0 | \bar{M}^0)(x) dx + C_T \int_{\mathbb{R}} |P(f_\varepsilon - \bar{M})| + |P(f_\varepsilon^0 - \bar{M}^0)| dx.$$

Integrating (5.4) and (5.5) with respect to (t, x) gives in particular

$$\int_{\mathbb{R}} |f_\varepsilon(t, x)| dx = \int_{\mathbb{R}} f_\varepsilon(t, x) dx = \int_{\mathbb{R}} f_\varepsilon^0(x) dx,$$

and

$$\int_{\mathbb{R}} |\bar{M}(t, x)| dx = \int_{\mathbb{R}} \bar{M}(t, x) dx = \int_{\mathbb{R}} \bar{M}^0(x) dx.$$

Thus

$$\begin{aligned} \int_{\mathbb{R}} |P(f_\varepsilon - \bar{M})| dx \\ \leq \left(1 + \sup_{i=1, \dots, N} |v_i| + \sup_{i=1, \dots, N} |v_i^2|\right) \left(\int_{\mathbb{R}} f_\varepsilon^0(x) dx + \int_{\mathbb{R}} \bar{M}^0(x) dx\right), \end{aligned}$$

and

$$\int_0^t \int_{\mathbb{R}} D(f_\varepsilon) dx ds \leq C_T^0 \varepsilon, \quad \text{for } 0 \leq t \leq T, \tag{5.7}$$

with

$$C_T^0 = \sup_{\varepsilon} \left(\int_{\mathbb{R}} \mathcal{H}(f_{\varepsilon}^0 | \overline{M}^0)(x) dx \right) + 4C_T \max(1, \sup_{i=1, \dots, N} |v_i^2|) \sup_{\varepsilon} \left(\int_{\mathbb{R}} f_{\varepsilon}^0(x) dx + \int_{\mathbb{R}} \overline{M}^0(x) dx \right).$$

We turn now to the estimation of $\mathcal{H}(f_{\varepsilon} | \overline{M})$ with respect to ε . For every $T < T^*$, there exists \tilde{C}_T such that

$$|\partial_{\overline{U}U}^2 \eta(\overline{U})|(t, x) \leq \tilde{C}_T, \quad |\partial_x \overline{U}|(t, x) \leq \tilde{C}_T,$$

for any $x \in \mathbb{R}, 0 \leq t \leq T$ and

$$\int_0^T \int_{\mathbb{R}} |\partial_x \overline{U}|^2(s, x) dx ds \leq \tilde{C}_T. \tag{5.8}$$

Then integrating (5.6) with respect to (t, x) gives

$$\begin{aligned} & \int_{\mathbb{R}} \mathcal{H}(f_{\varepsilon} | \overline{M})(t, x) dx - \int_{\mathbb{R}} \mathcal{H}(f_{\varepsilon}^0 | \overline{M}^0)(x) dx + \int_0^t \int_{\mathbb{R}} \frac{D(f_{\varepsilon})}{\varepsilon} dx ds \\ &= - \int_0^t \int_{\mathbb{R}} \partial_{\overline{U}U}^2 \eta(\overline{U}) \partial_x(\overline{U}) * \left(A(U_{\varepsilon} | \overline{U}) + P(V f_{\varepsilon} - V M_{\varepsilon}) \right) dx ds. \end{aligned}$$

Thanks to Proposition 3.3 and Lemma 3.2, we get

$$|A(U_{\varepsilon} | \overline{U})| \leq C_1 \eta(U_{\varepsilon} | \overline{U}) \leq C_1 \mathcal{H}(f_{\varepsilon} | \overline{M}).$$

Thanks to Proposition 4.1, we get

$$|PV(f_{\varepsilon} - M_{\varepsilon})|^2 \leq C_2 \left(\sum_{i=1}^N |(f_{\varepsilon})_i - (M_{\varepsilon})_i| \right)^2 = C_2 |f_{\varepsilon} - M_{\varepsilon}|^2 \leq C_3 D(f_{\varepsilon}).$$

Thus, it gives

$$\begin{aligned} & \int_{\mathbb{R}} \mathcal{H}(f_{\varepsilon} | \overline{M})(t, x) dx \\ & \leq \int_{\mathbb{R}} \mathcal{H}(f_{\varepsilon}^0 | \overline{M}^0)(x) dx + (\tilde{C}_T)^2 C_1 \int_0^t \int_{\mathbb{R}} \mathcal{H}(f_{\varepsilon} | \overline{M})(s, x) dx ds \\ & \quad + \left(\int_0^t \int_{\mathbb{R}} |\partial_{\overline{U}U}^2 \eta(\overline{U}) \partial_x(\overline{U})|^2 dx ds \right)^{1/2} \left(\int_0^t \int_{\mathbb{R}} |PV(f_{\varepsilon} - M_{\varepsilon})|^2 dx ds \right)^{1/2} \\ & \leq \int_{\mathbb{R}} \mathcal{H}(f_{\varepsilon}^0 | \overline{M}^0)(x) dx + \tilde{C}_T^2 C_1 \int_0^t \int_{\mathbb{R}} \mathcal{H}(f_{\varepsilon} | \overline{M})(s, x) dx ds \\ & \quad + \tilde{C}_T^{3/2} C_3^{1/2} \left(\int_0^t \int_{\mathbb{R}} D(f_{\varepsilon})(s, x) dx ds \right)^{1/2} \\ & \leq \int_{\mathbb{R}} \mathcal{H}(f_{\varepsilon}^0 | \overline{M}^0)(x) dx + \tilde{C}_T^2 C_1 \int_0^t \int_{\mathbb{R}} \mathcal{H}(f_{\varepsilon} | \overline{M})(s, x) dx ds + \tilde{C}_T^{3/2} C_3^{1/2} \sqrt{C_T^0 \varepsilon} \end{aligned}$$

using (5.7). Setting $w_\varepsilon(t) = \int_{\mathbb{R}} \mathcal{H}(f_\varepsilon | \overline{M})(t, x) dx$, it writes

$$w_\varepsilon(t) \leq w_\varepsilon(0) + C_4 \int_0^t w_\varepsilon(s) ds + C_5 \sqrt{\varepsilon}.$$

Using Gronwall's lemma, we get

$$\sup_{0 \leq t \leq T} \int_{\mathbb{R}} \mathcal{H}(f_\varepsilon | \overline{M})(t, x) dx \leq \left(\int_{\mathbb{R}} \mathcal{H}(f_\varepsilon^0 | \overline{M}^0) dx + C_5 \sqrt{\varepsilon} \right) e^{C_4 T}.$$

References

1. Beale, J.T.: Large-time behavior of discrete velocity Boltzmann equations. *Commun. Math. Phys.* **106**(4), 659–678 (1986)
2. Bellouquid, A.: On the asymptotic analysis of kinetic models towards the compressible Euler and acoustic equations. *Math. Models Methods Appl. Sci.* **14**(6), 853–882 (2004)
3. Berthelin, F., Bouchut, F.: Kinetic invariant domains relaxation limit from a BGK model to isentropic gas dynamics. *Asymptot. Anal.* **31**(2), 153–176 (2002)
4. Berthelin, F., Bouchut, F.: Relaxation to isentropic gas dynamics for a BGK system with single kinetic entropy. *Methods Appl. Anal.* **9**(2), 313–327 (2002)
5. Berthelin, F., Vasseur, A.: From kinetic equations to multidimensional isentropic gas dynamics before shocks. *SIAM J. Math. Anal.* **36**(6), 1807–1835 (2005)
6. Bianchini, S.: Hyperbolic limit of the Jin-Xin relaxation model. *Commun. Pure Appl. Math.* **59**(5), 688–753 (2006)
7. Bony, J.-M.: Solutions globales bornées pour les modèles discrets de l'équation de Boltzmann, en dimension 1 d'espace. In: Journées "Équations aux dérivées partielles", Saint Jean de Monts, 1987, pp. 1–10. École Polytechnique, Palaiseau (1987)
8. Bony, J.-M.: Existence globale à données de Cauchy petites pour les modèles discrets de l'équation de Boltzmann. *Commun. Part. Differ. Equ.* **16**(4-5), 533–545 (1991)
9. Brenier, Y.: Convergence of the Vlasov-Poisson system to the incompressible Euler equations. *Commun. Part. Differ. Equ.* **25**(3-4), 737–754 (2000)
10. Brenier, Y., Natalini, R., Puel, M.: On a relaxation approximation of the incompressible Navier-Stokes equations. *Proc. Am. Math. Soc.* **132**(4), 1021–1028 (2004)
11. Cabannes, H., Gatignol, R., Luo, L.-S.: The discrete Boltzmann equation (theory and applications). Henri Cabannes, Paris, 2003. Revised from the lecture notes given at the University of California, Berkeley, CA, 1980. Available electronically at http://lapasserelle.com/henri_cabannes/Cours_de_Berkeley.pdf
12. Caflisch, R.E., Papanicolaou, G.C.: The fluid dynamical limit of a nonlinear model Boltzmann equation. *Commun. Pure Appl. Math.* **32**(5), 589–616 (1979)
13. Dafermos, C.M.: The second law of thermodynamics and stability. *Arch. Ration. Mech. Anal.* **70**(2), 167–179 (1979)
14. DiPerna, R.J.: Uniqueness of solutions to hyperbolic conservation laws. *Indiana Univ. Math. J.* **28**(1), 137–188 (1979)
15. Feldman, M., Ha, S.-Y.: Nonlinear functionals of multi-D discrete velocity Boltzmann equations. *J. Stat. Phys.* **114**(3-4), 1015–1033 (2004)
16. Golse, F., Levermore, C.D.: Hydrodynamic Limits of Kinetic Models. *Topics in Kinetic Theory*, Fields Inst. Commun. pp. 1–75. Am. Math. Soc., Providence (2005)
17. Golse, F., Saint-Raymond, L.: The Navier-Stokes limit of the Boltzmann equation for bounded collision kernels. *Invent. Math.* **155**, 81–161 (2004)
18. Goudon, T., Nieto, J., Poupaud, F., Soler, J.: Multidimensional high-field limit of the electrostatic Vlasov-Poisson-Fokker-Planck system. *J. Differ. Equ.* **213**(2), 418–442 (2005)
19. Ha, S.-Y., Tzavaras, A.E.: Lyapunov functionals and L^1 -stability for discrete velocity Boltzmann equations. *Commun. Math. Phys.* **239**(1-2), 65–92 (2003)
20. Jourdain, B., Le Bris, C., Lelièvre, T., Otto, F.: Long-time asymptotics of a multiscale model for polymeric fluid flows. *Arch. Ration. Mech. Anal.* **181**(1), 97–148 (2006)
21. Kawashima, S.: Large-time behavior of solutions of the discrete Boltzmann equation. *Commun. Math. Phys.* **109**(4), 563–589 (1987)
22. Lattanzio, C., Tzavaras, A.E.: Structural properties of stress relaxation and convergence from viscoelasticity to polyconvex elastodynamics. *Arch. Ration. Mech. Anal.* **180**(3), 449–492 (2006)

23. Olla, S., Varadhan, S.R.S., Yau, H.-T.: Hydrodynamical limit for a Hamiltonian system with weak noise. *Commun. Math. Phys.* **155**(3), 523–560 (1993)
24. Płatkowski, T., Illner, R.: Discrete velocity models of the Boltzmann equation: a survey on the mathematical aspects of the theory. *SIAM Rev.* **30**(2), 213–255 (1988)
25. Saint-Raymond, L.: Convergence of solutions to the Boltzmann equation in the incompressible Euler limit. *Arch. Ration. Mech. Anal.* **166**(1), 47–80 (2003)
26. Serre, D.: Relaxations semi-linéaire et cinétique des systèmes de lois de conservation. *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* **17**(2), 169–192 (2000). [Semilinear and kinetic relaxations of systems of conservation laws]
27. Tzavaras, A.E.: Materials with internal variables and relaxation to conservation laws. *Arch. Ration. Mech. Anal.* **146**(2), 129–155 (1999)
28. Tzavaras, A.E.: Relative entropy in hyperbolic relaxation. *Commun. Math. Sci.* **3**(2), 119–132 (2005)
29. Ukai, S., Asano, K.: The Euler limit and initial layer of the nonlinear Boltzmann equation. *Hokkaido Math. J.* **12**, 311–332 (1983)
30. Villani, C.: Cercignani’s conjecture is sometimes true and always almost true. *Commun. Math. Phys.* **234**, 455–490 (2003)
31. Yau, H.-T.: Relative entropy and hydrodynamics of Ginzburg-Landau models. *Lett. Math. Phys.* **22**(1), 63–80 (1991)
32. Yu, S.-H.: Hydrodynamic limits with shock waves of the Boltzmann equation. *Commun. Pure Appl. Math.* **58**, 409–443 (2005)