

THE THERMAL CHLORINATION OF METHANE BY A FLOW METHOD

Sir:

We have recently been investigating the thermal (non-catalytic) chlorination of methane by a flow method. The following conclusions as to the kinetics of the reaction may be of interest.

The reaction takes place readily in pyrex glass reaction tubes above about 200° at 1 atm. total pressure. The rate is unaffected if the reaction tube is packed with broken glass, or if the internal surface of the tube is coated with potassium chloride. The reaction is, therefore, homogeneous. The effects of changes in concentration of reactants indicate that it is also bimolecular.

At the lower temperatures (225–300°) the reaction is inhibited by oxygen. Thus at 225° and five minutes' heating, 30% of the chlorine reacts when the reaction mixture consists of 60% methane and 40% chlorine at 1 atm. With 5% of oxygen present there is no measurable reaction. At 300° the suppression by oxygen is still considerable, but at 375°, at which temperature the rate of the unsuppressed reaction is very high, 5% of oxygen has no measurable effect. This effect of oxygen points to a chain mechanism. This is borne out by a discrepancy between the observed reaction rate at any temperature and that calculated by the collision theory of bimolecular gas reactions. About 1000 more molecules react than the theory predicts. A recent study of the photochemical chlorination of methane by Coehn and Cordes [*Z. physik. Chem.*, Abt B, **9**, 1 (1930)] has demonstrated that a chain mechanism applies here also, chains being some 10⁴ molecules in length at the maximum.

This work is being extended and will be later reported in full.

FRICK CHEMICAL LABORATORY
PRINCETON UNIVERSITY
PRINCETON, NEW JERSEY

RECEIVED DECEMBER 10, 1930
PUBLISHED JANUARY 12, 1931

ROBERT N. PEASE¹
GEORGE F. WALZ²

ADSORPTION AND PROMOTER ACTION IN CATALYSIS

Sir:

The results of Elissafoff's experiments [*Z. Electrochem.*, **21**, 352 (1915)] have been quoted by Hugh S. Taylor in his "Treatise upon Physical Chemistry" and also by Rudel and Haring [*Ind. Eng. Chem.*, **22**, 1234–1237 (1930)] as an example of adsorption in catalysis. The first investigator found that copper sulfate in the presence of glass wool decomposed hydrogen peroxide more rapidly than either substance alone, and that the effect was greater than the sum of the individual effects. Since separate experiments showed that a glass powder, of the same composition as the

¹ Research Associate, American Petroleum Institute.

² Research Assistant, American Petroleum Institute.

glass wool, removed copper from solution, Elissaffoff concluded that the copper was adsorbed at the interface and that the increased concentration of copper was responsible for the increase of rate. This conclusion was justified by close parallelism between the increased rate of reaction when the copper concentration was increased tenfold, and the increased amount of copper removed from a solution of similarly increased copper concentration.

At the time my attention was called to this article, I conducted a few brief tests with results incompatible with those of Elissaffoff. The experiments could not be amplified for lack of time, and therefore it seems best to indicate them briefly now. Hydrogen peroxide solutions were decomposed in quartz vessels by freshly prepared copper sulfate solutions containing one millimole per liter of copper sulfate. It was found that the velocity constant was nearly first order and not easily reproduced in the presumably neutral solutions employed. It is felt that the data from one series of experiments may be compared among themselves, however. It was also found that the addition of glass wool to a solution containing hydrogen peroxide and copper sulfate did increase the rate of decomposition but to a less extent than Elissaffoff observed. The reaction velocities for the promoted reaction increased appreciably with time, which points to a dependence of rate upon an adsorbed catalyst. However, it was found that a filtered extract of the glass wool increased the rate quite as much as the solid material. Also it was found that increasing the amount of glass wool ten-fold increased the rate of reaction very little. From these observations it was concluded that the glass wool no doubt removed copper from solution as Elissaffoff says, but not as copper sulfate. The copper probably formed a basic salt upon the surface of the glass wool, and then solid copper peroxide, a reaction which is favored by the alkali from the glass. The reaction in the absence of glass wool can be and probably is homogeneous; thus it seems probable that the mechanism of the reaction is different under "promotion" conditions than otherwise, and that this reaction is not an example of the effect of adsorption upon catalytic reaction rates.

DEPARTMENT OF CHEMISTRY
UNIVERSITY OF ILLINOIS
URBANA, ILLINOIS

RECEIVED DECEMBER 5, 1930
PUBLISHED JANUARY 12, 1931

A. C. ROBERTSON

THE OXIDATION OF CARBON MONOXIDE BY DISSOCIATED WATER VAPOR

Sir:

In a recent publication Harteck and Kopsch¹ have shown that atomic oxygen, from a discharge tube, has very little effect on carbon monoxide;

¹ Harteck and Kopsch, *Z. Elektrochem.*, 36, 714 (1930).