

nitrothiophenol. Here the intermediate formation of a sulfur bromide is hardly possible and the mechanism originally proposed by Kohlhasse seems the only one available. It is quite probable that in these reductions of all RSO_2X derivatives by phosphorus tribromide, *both* mechanisms may be operative, the relative speeds depending on conditions and on the nature of R and X, although the mechanism of Hunter and Sorenson is undoubtedly the chief one in *most* cases where a thiol is produced.

SCHOOL OF CHEMISTRY
UNIVERSITY OF MINNESOTA
MINNEAPOLIS, MINNESOTA

RECEIVED NOVEMBER 7, 1932
PUBLISHED MARCH 7, 1933

Notes on the Thompson-Oakdale Method for the Determination of Halogen in Organic Compounds

BY J. J. THOMPSON AND U. O. OAKDALE

The Thompson-Oakdale method¹ for the determination of halogen has been used with excellent results in hundreds of instances in our laboratories. However, the fact is not always recognized that the alkaline solution of the halide contains a considerable amount of sodium sulfite and consequently erroneous results may be obtained in the gravimetric estimation of the halogen due to the fact that the precipitated silver halide is apt to be contaminated, possibly with silver sulfide formed by decomposition of silver sulfite. This difficulty may be obviated in the following manner. The alkaline solution to which sodium arsenite has been added in accordance with the original directions is poured into a beaker, cooled thoroughly and after the addition of about 5 cc. of superoxol or perhydrol, to oxidize the alkaline sulfite to sulfate, the mixture is stirred and boiled for a few minutes. The solution is then cooled, acidified with nitric acid and the halide precipitated in the usual manner.

In view of our observations on the use of this method by students it seems necessary to emphasize the fact that *all* of the halogen, especially in the case of iodine, cannot be driven through the condenser into the alkaline solution unless the mixture in the Kjeldahl flask is boiled in such a manner that a continuous stream of steam passes through the apparatus for a short time.

The potassium persulfate used must always be tested for perchlorate since many commercial brands of persulfate contain this impurity. Potassium persulfate, free from perchlorate, is now available commercially (Mallinckrodt Chemical Works).

COLLEGE OF PHARMACY
DEPARTMENT OF CHEMISTRY
UNIVERSITY OF MICHIGAN
ANN ARBOR, MICHIGAN

RECEIVED DECEMBER 16, 1932
PUBLISHED MARCH 7, 1933

(1) Thompson and Oakdale, *THIS JOURNAL*, **52**, 1195 (1930).