

TABLE I

	Compound	Reaction temp., °C.	Yield, %	B. p., °C.	(Mm.)	M. p., °C.
1	2-Butyryl-6-ethylphenol	120	71	170-175	(1)	91-92
2	2-Butyryl-4-ethylphenol	120	50	112-113	(1)
3	2-Propionyl-6-propylphenol	120	87	164	(3)	78-79
4	2-Propionyl-4-propylphenol	120	61	115-118	(3)	(Phenylhydrazone, 96)
5	2-Butyryl-4,6-dimethylphenol	120	45	121	(2)	(Phenylhydrazone, 168-170)
6	2-Butyryl-4,5-dimethylphenol	120	89	40-41
7	4-Butyryl-2,5-dimethylphenol	25	70	131-132
in C ₆ H ₅ NO ₂						
8	3,3'-Diacetyl-4,4'-dihydroxydiphenyl	120	75	219-219.5
9	3,3'-Dipropionyl-4,4'-dihydroxydiphenyl	120	55	140-141
10	4-Caproylcatechol ^a	140	62	93-94
11	2-Valerylhydroquinone	180	70	174-176	(2)

	Physical form	Solvent for crystallization	Formula	Analyses, %			
				C	Calcd.	H	Found.
1	Colorless plates	Bz-pet. ether	C ₁₂ H ₁₆ O ₂	75.0	8.4	74.9	8.2
2	Oil	C ₁₂ H ₁₆ O ₂	75.0	8.4	74.9	8.6
3	Colorless plates	Pet. ether	C ₁₂ H ₁₆ O ₂	75.0	8.4	74.9	8.4
4	Oil	C ₁₂ H ₁₆ O ₂	75.0	8.4	74.5	8.5
5	Oil	C ₁₂ H ₁₆ O ₂	75.0	8.4	75.1	8.3
6	Colorless prisms	Dil. alc.	C ₁₂ H ₁₆ O ₂	75.0	8.4	75.2	8.7
7	Colorless prisms	Benzene	C ₁₂ H ₁₆ O ₂	75.0	8.4	74.8	8.6
8	Yellow needles	HAc	C ₁₆ H ₁₄ O ₄	71.1	5.2	70.8	5.0
9	Yellow needles	HAc	C ₁₈ H ₁₈ O ₄	72.5	6.1	72.6	6.4
10	Colorless plates	Bz	C ₁₂ H ₁₆ O ₂	75.0	8.4	74.8	8.6
11	Oil	C ₁₁ H ₁₄ O ₂	68.0	7.3	68.4	7.2

^a Coulthard, Marshall and Pyman, *J. Chem. Soc.*, 280 (1930).

TABLE II

Compound	B. p., °C.	(Mm.)	M. p., °C.	Yield, %	Analyses, %			
					C	Calcd.	H	Found
2-Butyl-6-ethylphenol	108-112	(2)	46	80.9	10.2	81.0	10.5
2-Butyl-4-ethylphenol	112-118	(5)	31	80.9	10.2	80.5	10.1
2,6-Dipropylphenol ^a	114-116	(5)	63	80.9	10.2	80.8	10.2
2,4-Dipropylphenol ^a	125-126	(10)	34	80.9	10.2	81.0	10.5
2-Butyl-4,6-dimethylphenol	106-108	(3)	80	80.9	10.2	80.5	10.3
2-Butyl-4,5-dimethylphenol	135-140	(8)	43-45	68	80.9	10.2	80.6	10.2
4-Butyl-2,5-dimethylphenol	130-132	(4)	70	80.9	10.2	80.7	10.9
3,3'-Diethyl-4,4'-dihydroxydiphenyl ^b	119-120	55	79.3	7.5	79.3	7.9
3,3'-Dipropyl-4,4'-dihydroxydiphenyl ^b	113-114	35	79.9	8.2	79.9	8.4
4-Hexylcatechol	169	(4.5)	25-26	50	74.3	9.5	74.2	9.8
2-Amylhydroquinone ^c	176-180	(3)	85-86	10	73.3	9.0	73.1	9.0

^a Prepared by Claisen [*Ann.*, **418**, 69 (1919)] by the reduction of the corresponding diallylphenols.

^b White needles from petroleum ether.

^c White plates from benzene.

from 150-170° was obtained consisting mainly of hydroquinone and finally 5 g. (10%) of amylydroquinone boiling at 176-180° at 3 mm. The reason for this behavior has not been investigated as yet. The physical constants of these phenols are given in Table II.

DEPARTMENT OF PHARMACOLOGY
VANDERBILT UNIVERSITY SCHOOL OF MEDICINE
NASHVILLE, TENNESSEE

RECEIVED JUNE 11, 1934

Isobutyrylformic Acid Phenylhydrazone

BY LYMAN C. CRAIG

In an earlier paper¹ we have reported the isolation of isobutyrylformic acid from the reaction mixture which results on treatment of ergotinine with aqueous alkali. The phenylhydrazone of the ketonic acid was prepared for final identifi-

(1) W. A. Jacobs and L. C. Craig, *J. Biol. Chem.*, **104**, 547 (1934).

cation. Although the analytical figures obtained with this derivative were satisfactory, its melting point (152°) proved to be higher than that recorded in the literature (128° ² and 143° ³). In order to clear up this discrepancy, we have since prepared the phenylhydrazone* of synthetic isobutyrylformic acid. The latter melts at 152° , thus confirming our observation on the substance obtained from ergotinine. A mixed melting point showed no depression.

The method of synthesis was essentially that reported by Tschelinzeff⁴ with a slight variation in the method of hydrolysis of the intermediate isobutyryl cyanide. It was found necessary to heat

(2) D. Tschtschenke, *Bull. soc. chim.*, [4] **37**, 623 (1925).

(3) E. Abderhalden and E. Rossner, *Z. physiol. Chem.*, **163**, 261 (1926).

(4) W. Tschelinzeff and W. Schmidt, *Ber.*, **62**, 2210 (1929).

the cyanide on the steam-bath for two hours in 1:1 hydrochloric acid in order to effect hydrolysis. The hydrolysis mixture was extracted with ether, the ether extract dried with sodium sulfate and evaporated to a sirup. If this sirup is used to prepare the hydrazone it does not crystallize. However, if the sirup is shaken with a little water, in which it is not completely soluble, and the aqueous layer used to prepare the hydrazone, the product crystallizes nicely. Repeated recrystallizations from dilute alcohol failed to raise the melting point above 152° .

Anal. Calcd. for $C_{11}H_{14}O_2N_2$: C, 64.08; H, 6.84; N, 13.59. Found: C, 64.15; H, 6.74; N, 13.66.

FROM THE LABORATORIES OF
THE ROCKEFELLER INSTITUTE
FOR MEDICAL RESEARCH
NEW YORK, N. Y.

RECEIVED JUNE 28, 1934

COMMUNICATIONS TO THE EDITOR

ORGANIC REACTIONS WITH LIQUEFIED HYDROGEN CHLORIDE

Sir:

Information is accumulating in our laboratories relative to the reaction of liquefied hydrogen chloride with various types of organic compounds. In the majority of cases the reactions proceed rapidly indeed, partly because of the fact that the liquefied hydrohalides, in particular hydrogen chloride, are excellent solvents for a wide range of compounds. Many of the resulting solutions are good conductors of electricity.

The reactions observed so far are additions to multiple bonds, dehydration, or both; acidolysis appears not to be as frequent as might be anticipated. Reactions of hydrohalides as exemplified by our experiments are interesting in view of theoretical considerations concerned with addition to multiple bonds. As example of these reactions the following are typical: (1) vinyl acetate reacts with liquid hydrogen chloride at atmospheric pressure forming α -chloroethyl acetate in 50 to 70% yields. The literature contains two boiling points for this compound, 119 – 121° (740 mm.¹) and 113 – 116° (740 mm.²); different preparations resulted in products which boiled at either one or the other

(1) Descade, *Ann. chim.*, [7] **29**, 488 (1903).

(2) Ulich and Adams, *THIS JOURNAL*, **43**, 660 (1921).

of the two points. This is being investigated. (2) Allyl alcohol (b. p. 96.5 – 97.5°) is converted to allyl ether (b. p. 91 – 93°) with a high yield. (3) Acetaldehyde yields α, α' -dichloroethyl ether boiling at 112.5 – 114° in yields of 70%, n_D^{25} 1.4186, d_4^{25} 1.106. This compound was identified by hydrolysis to acetaldehyde in the presence of *p*-nitrophenylhydrazine and isolation of the hydrazone. (4) Good yields of ethylene chlorohydrin are obtained from the reaction of ethylene oxide with liquefied hydrogen chloride.

A mixture of one mole of isopropyl acetate and one mole of *n*-butyl alcohol was treated with liquid hydrogen chloride at atmospheric pressure, from which was isolated some isopropyl alcohol and butyl acetate. This is being investigated on a series of such compounds with the purpose of obtaining data relative to negativity of the groups tested.

A number of cases have been found in which no action takes place. Cinnamic acid is a typical example. This work is being extended. A complete report will be made later.

NORTHWESTERN UNIVERSITY ERICH GEBAUER-FUELNEGG
DEPARTMENT OF CHEMISTRY
EVANSTON, ILL.

BORG LABORATORIES
CHICAGO, ILL.

EUGENE MOFFETT

RECEIVED JULY 11, 1934