

The Preparation of Certain Nitrogen-Substituted Sulfon-*m*-toluidides and Sulfon-*p*-toluidides

BY GEORGE H. YOUNG

This paper is a continuation of an investigation of *N*-alkyl-*p*-toluene sulfonanilides.¹ Yields of 80–98% of *m*-sulfontoluidides and 60–91% of *p*-isomers were obtained. Recrystallized from methanol (the ethyl *m*-toluidide from 5:1 ethanol-ether), these compounds deposit as blunt needles or thick prisms. They are insoluble in water, difficultly soluble in ether and soluble in most other common solvents, with the meta derivatives usually having the greater solubility.

TABLE I

THE SULFON- <i>m</i> -TOLUIDIDES				
<i>p</i> -Toluene sulfon- <i>m</i> -toluidide	Formula	M. p., °C.	Sulfur analyses, %	
			Calcd.	Found
<i>N</i> -Methyl	C ₁₅ H ₁₇ O ₂ NS	63	11.64	11.61
Ethyl	C ₁₆ H ₁₉ O ₂ NS	60.5–61	11.07	11.08
<i>n</i> -Propyl	C ₁₇ H ₂₁ O ₂ NS	51	10.56	10.71
Isoamyl	C ₁₉ H ₂₅ O ₂ NS	76.5	9.55	9.61

TABLE II

THE SULFON- <i>p</i> -TOLUIDIDES				
<i>p</i> -Toluene sulfon- <i>p</i> -toluidide	Formula	M. p., °C.	Sulfur analyses, %	
			Calcd.	Found
<i>N</i> -Isopropyl	C ₁₇ H ₂₁ O ₂ NS	107	10.56	10.76
Isobutyl	C ₁₈ H ₂₃ O ₂ NS	73	10.09	10.04
<i>n</i> -Amyl	C ₁₉ H ₂₅ O ₂ NS	59.5–60	9.55	9.70
Isoamyl	C ₁₉ H ₂₅ O ₂ NS	89.5–90	9.55	9.43

The writer gratefully acknowledges the advice and assistance of the late Dr. W. J. Keith, who suggested this research.

(1) Young, *THIS JOURNAL*, **56**, 2167, 2783 (1934).

SCHOOL OF CHEMISTRY AND PHYSICS
PENNSYLVANIA STATE COLLEGE
STATE COLLEGE, PENNA. RECEIVED JANUARY 28, 1935

Preparation of 5-Keto-gluconic Acid by Bromine Oxidation

BY ELMER W. COOK AND RANDOLPH T. MAJOR

In the oxidation of glucose with bromine generated electrolytically according to the Isbell¹ method it was noticed that a reducing substance remained even after electrolysis had been continued longer than was necessary theoretically to convert the glucose to gluconic acid.

Sheppard and others² have observed likewise the presence of a reducing substance during the oxidation of glucose or gluconic acid with bromine.

This reducing substance has been isolated and

(1) Isbell and Frush, *Bur. Standards J. Research* **6**, 1145 (1931).

(2) Everett, Edwards and Sheppard, *J. Biol. Chem.*, **104**, 11 (1934); Sheppard and Everett, *ibid.*, *Proc. Am. Soc. Biol. Chemists*, XXVIII, Annual Meeting, p. LXXX, 1934.

proved to be the 5-keto-gluconic acid first described by Boutroux.³

Experimental

The oxidation of glucose was carried out as described by Isbell. The electrolyte consisted of 7 g. of glucose, 5 g. of calcium carbonate, and 5 g. of calcium bromide in 140 cc. of water. Electrodes of platinum were used. The electrolysis was continued for four ampere hours at a current density of 2.5 amperes per sq. dm.

At the end of the electrolysis the calcium carbonate was filtered and the resulting solution concentrated to 30 cc. Alcohol was added to a slight permanent turbidity and the solution allowed to stand for two to three weeks. The precipitate which had formed slowly during this time was filtered and dissolved in boiling water. After standing for forty-eight hours a calcium salt had settled out in the form of a fine crystalline crust consisting of small rhombohedra; yield 0.8 g. The properties⁴ of this salt agreed in every respect with those of the well-known calcium 5-keto-gluconate. It reduced Fehling's solution, dissolved in about 50 parts of boiling water, and lost practically no weight when dried *in vacuo* at 85–90° (distinction from calcium saccharate). A 2% solution obtained by dissolving the calcium salt in the calculated amount of hydrochloric acid gave α^{20}_D -14° , based on the free acid.

Anal. Calcd. for Ca(C₆H₉O₇)₂·3H₂O: Ca, 8.34; C, 29.98; H, 5.04. Found: Ca, 8.46; C, 30.18; H, 5.18.

(3) Boutroux, *Compt. rend.*, **102**, 924 (1886); **127**, 1224 (1898).

(4) Ruff, *Ber.*, **32**, 2270 (1899); Kiliani, *ibid.*, **55**, 2820 (1922); Barch, *THIS JOURNAL*, **56**, 3653 (1933).

LABORATORY FOR PURE RESEARCH
MERCK & Co., INC.
RAHWAY, N. J.

RECEIVED FEBRUARY 7, 1935

The Indium Salts of Some Organic Acids

BY JOHN B. EKELEY AND WAYNE W. JOHNSON

Indium hydroxide forms basic salts when treated with propionic, *n*-butyric, trichloroacetic, benzoic or with ortho, meta or para-toluic acids.

The propionate and the *n*-butyrate were prepared by dissolving weighed quantities of the hydroxide in a slight excess of the pure acids and the excess acid evaporated off on the water-bath until the salts were precipitated. The precipitates were boiled with alcohol, filtered, washed with ether and dried over sulfuric acid. These salts are white powders insoluble in water, alcohol, and ether; yield quantitative.

The trichloroacetate was prepared in the same manner as the two preceding salts, except that, since the salt is soluble in water, alcohol and ether, the excess of acid was removed by drying the product to constant weight at 110°; yield quantitative.

The benzoate was prepared by precipitating