

investigation, it is necessary to defer publication of the details of the work until a later date.

LABORATORY FOR ENDOCRINE RESEARCH H. JENSEN
SCHOOL OF MEDICINE
JOHNS HOPKINS UNIVERSITY
BALTIMORE, MARYLAND

RECEIVED MARCH 10, 1937

**STEROLS. XII. SYNTHETIC PREPARATION OF
EPI-ALLO-PREGNANOLONE, THE ANDROGENIC
PRINCIPLE OF HUMAN PREGNANCY URINE**

Sir:

That human pregnancy urine contains a substance which has male hormone activity has been reported by several workers. Because of the similarity in physiological activities in the male hormone tests between human pregnancy and male urine extracts, it has been assumed that the male principle in pregnancy urine was androsterone. Heretofore this androgenic principle has not been isolated.

Recently, Marker, Kamm and McGrew [THIS JOURNAL, **59**, 616 (1937)] reported the isolation and identification of *epi*-allo-pregnanolone from the sterol fraction of human pregnancy urine after the removal of theelin and theelol. At the time of this report we did not have the physiological assays on this compound. Since then it has been found that the substance isolated from human pregnancy urine promotes the growth of the seminal vesicle in the rat tests. Preliminary tests have shown this to have about the same activity as androsterone.

We have now prepared synthetically *epi*-allo-pregnanolone from 3-chloro-*allo*-cholanolic acid, a by-product from the preparation of chloroandrosterone [Marker, Whitmore and Kamm, THIS JOURNAL **57**, 2358 (1935)]. This was prepared by the stepwise degradation of the side-chain according to the method of Wieland [*Z. physiol. Chem.*, **161**, 80 (1926)].

3-Chloro-*allo*-cholanolic acid, m. p. 180°, was converted into its methyl ester, m. p. 133°. *Anal.* Calcd. for $C_{26}H_{41}O_2Cl$: C, 73.2; H, 10.1; Found: C, 72.8; H, 10.3. This was treated with a phenyl Grignard reagent and the resulting carbinol melted at 171°. *Anal.* Calcd. for $C_{30}H_{49}OCl$: C, 81.1; H, 9.2. Found: C, 81.1; H, 9.3. It was oxidized by chromic acid to 3-chloro-*allo*-*nor*-cholanolic acid. *Anal.* Calcd. for $C_{23}H_{37}O_2Cl$: C, 72.5; H, 9.8. Found: C, 72.8; H, 9.6. This acid was converted to its methyl ester, m. p. 178°.

Anal. Calcd. for $C_{24}H_{39}O_2Cl$: C, 73.0; H, 10.0. Found: C, 73.3; H, 10.2. By treating with a phenyl Grignard reagent, 3-chloro-*allo*-bis-*nor*-cholanyldiphenylcarbinol was obtained, m. p. 183°. *Anal.* Calcd. for $C_{35}H_{47}OCl$: C, 80.9; H, 9.1. Found: C, 80.8; H, 9.3. This carbinol was oxidized by chromic acid to 3-chloro-bis-*nor*-*allo*-cholanolic acid, m. p. 231°. *Anal.* Calcd. for $C_{22}H_{35}O_2Cl$: C, 71.9; H, 9.6. Found: C, 71.8; H, 9.3. This acid was converted to its methyl ester, m. p. 150°. *Anal.* Calcd. for $C_{23}H_{37}O_2Cl$: C, 72.5; H, 9.8. Found: C, 72.5; H, 9.8. By treating with a phenyl Grignard reagent the carbinol was obtained, m. p. 146°. *Anal.* Calcd. for $C_{34}H_{45}OCl$: C, 80.8; H, 9.0. Found: C, 80.9; H, 9.2. This carbinol was dehydrated by refluxing with acetic acid and acetic anhydride. The resulting product was ozonized and the ozonolysis product treated with potassium acetate to convert the chloro derivative into an *epi*-OH compound [Marker, Whitmore and Kamm, THIS JOURNAL, **57**, 2358 (1935)]. The hydroxy ketone was purified by means of its half succinic ester and then as the semicarbazone. It gave *epi*-allo-pregnanolone, m. p. 170°. *Anal.* Calcd. for $C_{21}H_{34}O_2$: C, 79.2; H, 10.7. Found: C, 78.9; H, 10.9. This was identical with the natural product which we isolated from human pregnancy urine.

THE SCHOOL OF
CHEMISTRY AND PHYSICS
PENNSYLVANIA STATE COLLEGE
STATE COLLEGE, PA.
PARKE, DAVIS & CO.
RESEARCH LABORATORIES
DETROIT, MICH.

RUSSELL E. MARKER
OLIVER KAMM
DAVID M. JONES
EUGENE L. WITTE
THOMAS S. OAKWOOD
HARRY M. CROOKS

RECEIVED MARCH 18, 1937

STEROLS. XIII. THE DIHYDROEQUILENINS

Sir:

Recently Wintersteiner, Schwenk, Hirschmann and Whitman [THIS JOURNAL, **58**, 2652 (1936)] in their studies on the isolation of the δ -follicular hormone from the phenolic fraction of mare's urine obtained as the δ -follicular fraction a molecular compound of two components, one of which formed a picrate. This was isolated and identified as dihydroequilenin. They suggested that possibly the high estrogenic activity of the originally reported δ -follicular hormone [Wintersteiner, Schwenk and Whitman, *Proc. Soc. Exptl. Biol. Med.*, **32**, 1087 (1935)] is due to the other