


dehydrotigogenone in 20 cc. of peanut oil. The 4-dehydrotigogenone was carefully purified and gave no precipitate with digitonin. The feces were collected for six days and treated essentially by the method of Schoenheimer, Rittenberg and Graff. From the fraction precipitated with digitonin 0.2 g. of diosgenin, m. p. 207–209°, was obtained. This was also identified as its acetate, m. p. 199–200°. The fraction (7.2 g.) of non-saponifiable material not precipitated with digitonin gave 3.2 g. of unchanged 4-dehydrotigogenone.



Since diosgenin could only have been produced from the administered 4-dehydrotigogenone, this experiment definitely establishes the bio-reduction of a  $\Delta^4$ -3-keto-steroid to a  $\Delta^5$ -3-hydroxy-steroid.

The experiments of Callow [*Biochem. J.*, **33**, 559 (1939)] cited by Fieser are subject to the same objection raised by Fieser in connection with Schoenheimer's work since the administered testosterone propionate was not "labelled." In addition it should be pointed out that negative results in urine work can hardly be regarded as significant.

It does not seem to be generally known that cholestenone has been converted to cholesterol by purely chemical methods. This has been reported by several authors [Wagner-Jauregg and Werner, *Z. physiol. Chem.*, **208**, 72 (1932); Lettré, *ibid.*, **221**, 73 (1933)].

RUSSELL E. MARKER  
THE SCHOOL OF CHEMISTRY AND PHYSICS R. B. WAGNER  
THE PENNSYLVANIA STATE COLLEGE D. L. TURNER  
STATE COLLEGE, PENNA. EMERSON L. WITTEBECKER

RECEIVED MAY 16, 1941

### IMPROVED COMPUTATIONS ON CONJUGATION AND HYPERCONJUGATION

Sir:

Using molecular orbital theory, we recently computed conjugation and hyperconjugation energies and corresponding bond distance changes in good agreement with observation, but could explain observed hyperconjugation shifts in ultraviolet spectra only qualitatively.<sup>1</sup> New computations, using a better approximation, give satisfactory quantitative agreements with spectra as well as energies and distances, for both conjugation and hyperconjugation.

The secular equation for a molecule containing two conjugated bonds is (*cf.* (6) of ref. 1)

$$\begin{vmatrix} \gamma^* - x & \gamma^* - S^*E & 0 & 0 \\ \gamma^* - S^*E & x & \gamma - SE & 0 \\ 0 & \gamma - SE & x & \gamma^* - S^*E \\ 0 & 0 & \gamma^* - S^*E & x \end{vmatrix} = 0 \quad (1)$$

where  $x \equiv \alpha - E$ . Asterisks denote integrals for multiple bonds. In solving this equation it has been customary to neglect all  $SE$ 's, but this procedure is invalid, since  $\gamma$  and  $SE$  are approximately equal (*sec.* 7). Nevertheless it has seemingly given useful results.

To understand how, we write

$$\gamma - SE = (\gamma - S\alpha) + S(\alpha - E) \equiv \beta + Sx \quad (2)$$

and (1) becomes

$$\begin{vmatrix} x & \beta^* + S^*x & 0 & 0 \\ \beta^* + S^*x & x & \beta + Sx & 0 \\ 0 & \beta + Sx & x & \beta^* + S^*x \\ 0 & 0 & \beta^* + S^*x & x \end{vmatrix} = 0 \quad (3)$$

In general,  $x$  is of the order of the  $\beta$ 's, and since  $S$  for unsaturation electrons is about 0.25, the  $Sx$ 's are of smaller magnitude than the  $\beta$ 's. However, for the quasi-unsaturation electrons of hyperconjugation,  $S$  is about 0.6 and the  $Sx$ 's are more important.

Equation (1) neglecting  $SE$  is formally identical with (3) neglecting  $Sx$ . Thus while previous workers seemingly solved (1) with the poor approximation  $SE = 0$ , we see that in effect they solved (3) with the better approximation  $Sx = 0$ , if we merely reinterpret their empirical resonance parameter as  $\beta$  instead of  $\gamma$ . This was done in our paper.

Our present procedure solves (3) exactly.<sup>2</sup> The numerical labor is increased, but no real difficul-

(1) *THIS JOURNAL*, **63**, 41 (1941).

(2) Eqs. (1), (3) themselves involve an approximation in that relatively small matrix elements of non-neighboring atoms are replaced by zero. Our ability to fit spectroscopic as well as thermal data supports the assumption that these may still be neglected.

ties are introduced. By solving exactly the secular equation in form (3) instead of (1), we obtain resonance and excitation energies independent of  $\alpha$  without having to assume the  $\beta$ 's and  $S$ 's constant for all distances and for different atoms. Each  $S$  is calculated using Slater functions. The parameters  $\rho$ ,  $\eta$ , and  $\beta_{1.33}$  (see (24) of ref. 1) are determined, as before, to fit experimental data. The variation of  $\rho$  with distance now found is less than half as rapid as before;  $-\beta_{1.33}$  is increased from 44.5 to about 56 kcal.; and  $\eta$  is reduced from 4 to about 1.9.<sup>3</sup>

The improvement in our computations has been obtained without sacrificing essential simplicity and ease of computation; the results support the general validity of the molecular orbital approximation.

(3) In sec. 14, we indicated that 4 is a reasonable value for  $\eta$ . Actually the present value is much more reasonable, since corrections previously neglected (cf. footnote 30) reduce the estimated  $\eta$  to about 1.5.

RYERSON PHYSICAL LABORATORY  
UNIVERSITY OF CHICAGO  
CHICAGO, ILLINOIS

ROBERT S. MULLIKEN  
CAROL A. RIEKE

RECEIVED APRIL 12, 1941

#### THE ACTION OF METHYLAMINE WITH NITROUS ACID

Sir:

In view of the experience of one of us (L. U. S.) in the successful quantitative determination of methylamine by the evolution of nitrogen on treatment with nitrous acid [Van Slyke, *J. Biol. Chem.*, **9**, 185 (1911)] it seems strange that methylamine should differ so greatly from *n*-butylamine in its reactivity with nitrous acid in aqueous hydrochloric acid solution [Whitmore, *et al.*, *THIS JOURNAL*, **54**, 3441 (1932); **63**, 1118 (1941)]. Apparently, the difference is due to the difference between an aqueous hydrochloric acid solution and an acetic acid solution containing no mineral acid and relatively little water. We now find that a solution of 1 mole of methylamine in 1 liter of glacial acetic acid when added to 5 moles of powdered sodium nitrite gives no appreciable action. On addition of 100 ml. of water, reaction starts and continues until about half a mole of nitrogen has been evolved. The reaction then slackens. An additional 100 ml. of water starts the reaction and finally gives, on heating, nearly half a mole of nitrogen. The gases were passed through dry-ice traps and a scrubber containing alkaline potassium permanganate. The chief organic product

of the reaction was methyl acetate. The only other product isolated was a trace of methyl nitrite.

Apparently, the excess of nitrous acid and the minimum amount of water cut down the hydrolysis of methylamine nitrite earlier observed. It is also probable that the acetate ion with its cloud of electrons can make a more effective attack on the side of the carbon opposite the nitrogen than can hydroxyl ions, water molecules or chloride ions.

This work is being continued.

ROHM AND HAAS COMPANY  
PHILADELPHIA, PENNSYLVANIA  
THE PENNSYLVANIA STATE COLLEGE  
STATE COLLEGE, PENNSYLVANIA

L. U. SPENCE  
FRANK C. WHITMORE  
J. D. SURMATIC

RECEIVED MAY 6, 1941

#### *p*-AMINO BENZOIC ACID AND TYROSINASE ACTIVITY

Sir:

Since *p*-aminobenzoic acid has been reported to have chromotrichial activity [*Science*, **93**, 164 (1941); *J. Biol. Chem.*, **138**, 441 (1941)], we investigated its influence on dopa reaction and noted [*Proc. Soc. Exp. Biol. Med.*, **47**, May (1941)] it to modify enzymatic formation of melanin. Using the Warburg apparatus [kindly placed at our disposal by Reverend J. B. Meunzen, F.G., at Fordham University], we determined its effect on the kinetics of tyrosinase action. In numerous experiments, the detailed data of which are about to be published, we found that the aerobic oxidation of tyrosine and that of dopa is retarded by *p*-aminobenzoic acid, but that of *p*-cresol is accelerated. Qualitatively, aniline has the same influence as *p*-aminobenzoic acid; quantitatively, the effect of the latter is greater than that of the former.

In a typical set of experiments a reaction mixture was employed consisting of 0.4 ml. of 0.0463 *M* *p*-cresol, 1 ml. of McIlvaine buffer (*pH* 6.5), 0.5 ml. (2.5 mg.) of gelatin solution, 1 ml. of appropriately diluted enzyme solution [kindly furnished by Dr. J. M. Nelson, Columbia University; isolated from *psalliota campestris* and containing approximately 200 hydroquinone-catechol units and 50 cresolate units per ml. measured at 25°], and 1.1 ml. of water or 0.1 ml. of water and 1.0 ml. of 0.01 *M* solution of the test substance. All the determinations were made at a temperature of 37.2°, the rates of oxygen uptake were calcu-