

and crystallized from mixtures of polar and non-polar solvents. Diethoxyethylguanidine sulfate crystallized best from *n*-butanol-ethyl acetate. The others were crystallized from alcohol-ether.

N-Benzyl-N- α -naphthylguanidine Hydrochloride.—Benzyl- α -naphthylamine was refluxed in amyl alcohol with 1 mol of cyanamide and 1.2 mols of hydrogen chloride. On cooling, ether was added and the hydrochlorides separated as a purple sirup. Attempts to crystallize having failed, the mixture was dissolved in water and ammonia added. Some unreacted benzylnaphthylamine separated together with most of the color. The aqueous layer was then basified with sodium hydroxide and the precipitated guanidine taken into ether. After drying over potassium carbonate, alcoholic hydrogen chloride was added. The salt was recrystallized from alcohol-ether mixtures.

N,N'-Dihomoanisylguanidine Hydrochloride.—To a

$[p\text{-XC}_6\text{H}_4\text{OCH}_2\text{CH}_2\text{OCH}_2\text{CH}_2\text{N}^+(\text{CH}_2)_2\text{R}]\text{Y}^-$					Analyses, %			
R	X	Y	M. p., °C.	Formula	Calcd.		Found	
					C	H	C	H
CH ₃	Cl	I	152	C ₁₇ H ₂₀ O ₂ NCI	46.19	6.62	46.20	6.42
C ₆ H ₅ CH ₂	H	Cl	122-123	C ₂₃ H ₂₄ O ₂ NCl	3.58 ^a	9.05 ^b	3.91 ^a	9.21 ^b
<i>p</i> -ClC ₆ H ₄ CH ₂	H	Cl	166-166.5	C ₂₃ H ₁₈ O ₂ NCl ₂	64.76	7.80	64.85	7.80
<i>p</i> -ClC ₆ H ₄ CH ₂	Cl	Cl	160	C ₂₃ H ₁₂ O ₂ NCl ₃	59.92	7.00	60.12	7.01
<i>p</i> -BrC ₆ H ₄ CH ₂	Cl	Cl	156.5-157	C ₂₃ H ₁₂ O ₂ NCl ₂ Br	54.64	6.38	54.95	6.35

^a N, ^b Cl.

solution of 2 mols of homoanisylamine in absolute ether was added, with shaking and ice-cooling, a solution of 1 mol of cyanogen bromide in ether. After standing one-half hour the ether was evaporated by a stream of dry air, a little absolute alcohol added to homogenize the mixture and the whole was heated three hours in an oil-bath at 150°. The material was then dissolved in water, the base liberated with alkali, and extracted with ether. On drying over potassium carbonate a crystalline solid appeared on the surface of the drying agent. The ether was decanted off, the solid dissolved in chloroform and transformed into the hydrochloride by alcoholic hydrogen chloride. It was recrystallized from alcohol-ether, forming lustrous plates.

N-Methyl-N'- α -naphthylguanidine.—Methylamine and α -naphthylisothiocyanate yielded N,N'-methyl-naphthylthiourea. Methylation with methyl sulfate gave the S-methyl derivative which was desulfurized in the usual manner with lead oxide and ammonia.

Data on these compounds are presented in the table.

Substance	M. p., °C.	Empirical formula	Analyses, %			
			Calcd.		Found	
			C	H	C	H
β -N-Morpholinoethylguanidine sulfate	197	C ₇ H ₁₆ ON ₄ (H ₂ SO ₄) _{1/2}	37.99	7.75	38.33	7.52
β,β -Diethoxyethylguanidine sulfate	154	C ₇ H ₁₇ O ₂ N ₃ (H ₂ SO ₄) _{1/2}	37.47	8.09	37.38	7.99
N,N-Dicyclohexylguanidine sulfate	195	C ₁₃ H ₂₅ N ₃ (H ₂ SO ₄) _{1/2}	57.28	9.62	57.05	9.28
N-Benzyl-N-methylguanidine sulfate	252 dec.	C ₉ H ₁₃ N ₃ (H ₂ SO ₄) _{1/2}	50.90	6.65	50.87	6.59
δ -Phenoxybutylguanidine sulfate	199-199.5	C ₁₁ H ₁₇ ON ₃ (H ₂ SO ₄) _{1/2}	51.53	7.08	51.32	7.33
N-Benzyl-N- α -naphthylguanidine hydrochloride	223-224	C ₁₈ H ₁₇ N ₃ ·HCl	69.33	5.82	69.19	5.80
N,N'-Dihomoanisylguanidine hydrochloride	125.5-126.5	C ₁₉ H ₂₆ O ₂ N ₃ ·HCl	62.69	7.21	62.63	7.05
N- α -Naphthyl-N'-methylguanidine hydrochloride	220-220.5 dec.	C ₁₂ H ₁₃ N ₃ ·HCl	61.12	5.99	61.06	5.99

BURROUGHS WELLCOME & Co., U. S. A. JOHANNES S. BUCK
EXPERIMENTAL RESEARCH LABORATORIES
TUCKAHOE, NEW YORK

RICHARD BALTZLY
CLAYTON W. FERRY

RECEIVED MAY 18, 1942

Some Quaternary Salts from β -Dimethylamino- β' -cymoxydiethyl Ether

A concentrated aqueous solution of sodium thymolate or sodium *p*-chlorothymolate was heated under reflux with β,β' -dichlorodiethyl ether. After separation of the aqueous layer, unreacted dichlorodiethyl ether was removed *in vacuo*, and the residue was heated for seven hours at 145° (*p* = ca. 150 lb.) in a glass-lined bomb with 33% methanolic dimethylamine. After removal of volatile materials on the steam-bath under diminished pressure, the residual tertiary amines were partially purified by solution in acid and extraction with ether; on liberation with alkali, they were obtained as oils which could not be distilled *in vacuo*, but were converted directly into quaternary salts by warming on the steam-bath with the appropriate halides. The salts crystallized from acetone or alcohol on addition of ether.

BURROUGHS WELLCOME AND CO., U. S. A.
EXPERIMENTAL RESEARCH LABORATORIES
TUCKAHOE, NEW YORK

CLAYTON W. FERRY
ALAN E. ARDIS
JOHANNES S. BUCK

RECEIVED MAY 18, 1942

N,N-Dimethylethylenediamine and Some Derivatives

The readily available dimethyl glycine nitrile¹ can be reduced by sodium and absolute alcohol to give N,N-dimethylethylenediamine. This is a colorless liquid boiling, when anhydrous, at 107°. As its dehydration is difficult and its dihydrochloride (melting around 160°) is also hygroscopic, it is better characterized through a derivative. The following compounds were prepared as outlined, nitro derivatives being reduced with Adams catalyst in alcoholic solution containing hydrogen chloride. β -[*p*-Nitrobenzoylamidoethyl] dimethylamine hydrochloride (I), formed from *p*-nitrobenzoyl chloride and the

diamine, was hydrogenated catalytically yielding β -[*p*-aminobenzoylamidoethyl]-dimethylamine dihydrochloride (II). The methochloride of I was reduced to β -[*p*-amino-

¹ J. v. Braun, *Ber.*, **40**, 3937 (1907).

benzoylamidoethyl]-trimethylammonium chloride, hydrochloride (III).

The diamine reacted with *p*-nitrophenyl isocyanate to give β -[*p*-nitrophenylureidoethyl]-dimethylamine; m. p. of the hydrochloride, 247–248.5°. From this were obtained β -[*p*-aminophenylureidoethyl]-dimethylamine di-

Some Unsymmetrical Disubstituted Ureas

The substances, data on which are presented in the subjoined table, were prepared by the action of nitrourea on the corresponding secondary amines.¹ They crystallize in colorless prisms from alcohol or benzene–petrol ether.

UNSYMMETRICAL DISUBSTITUTED UREAS R,R'NCONH₂

Substituents		Formula	M. p., °C.	Analyses, %	
R	R'			Calcd. N	Found N
CH ₃	(<i>n</i>)C ₆ H ₁₃	C ₈ H ₁₈ ON ₂	75	17.72	17.92
(4)CH ₃ OC ₆ H ₄	C ₂ H ₅ (CH ₃)CH	C ₁₂ H ₁₈ O ₂ N ₂	140	12.61	12.70
(4)CH ₃ OC ₆ H ₄	C ₂ H ₅ (CH ₃)CHCH ₂	C ₁₃ H ₂₀ O ₂ N ₂	130	11.86	12.17
(4)CH ₃ OC ₆ H ₄	(CH ₃) ₃ CCH ₂	C ₁₃ H ₂₀ O ₂ N ₂	155	11.86	12.13
(4)CH ₃ OC ₆ H ₄	(CH ₃) ₂ CHCH ₂ (CH ₃)CH	C ₁₄ H ₂₂ O ₂ N ₂	110	11.19	11.25

hydrochloride (IV) and the corresponding trimethylammonium chloride, hydrochloride (V). The diamine with *p*-nitrophenylacetyl chloride gave the *p*-nitrophenylacetamide, m. p. of the hydrochloride, 190–192.5°, and reduction of this yielded β -[*p*-aminophenylacetamidoethyl]-dimethylamine dihydrochloride (VI). Reduction of the methochloride of the nitro compound gave β -[*p*-

(1) Cf. Buck and Ferry, *THIS JOURNAL*, **58**, 854 (1936).

BURROUGHS WELLCOME & CO., U. S. A.
EXPERIMENTAL RESEARCH LABORATORIES
TUCKAHOE, NEW YORK

JOHANNES S. BUCK
WALTER S. IDE
RICHARD BALTZLY

RECEIVED JUNE 6, 1942

DERIVATIVES OF N,N-DIMETHYLETHYLENEDIAMINE

No.	Formula	M. p., °C.	Analyses, %			
			Calcd.		Found	
			C	H	C	H
I	O ₂ NC ₆ H ₄ CONHCH ₂ CH ₂ NMe ₂ ·HCl	182.5–183.5	48.24	5.89	48.09	5.90
II	H ₂ NC ₆ H ₄ CONHCH ₂ CH ₂ NMe ₂ ·2HCl	190–191	47.12	6.84	47.23	6.89
III	H ₂ NC ₆ H ₄ CONHCH ₂ CH ₂ NMe ₂ ·Cl·HCl	dec. > 230	48.96	7.20	49.37	7.39
IV	H ₂ NC ₆ H ₄ NHCONHCH ₂ CH ₂ NMe ₂ ·2HCl	182–184 dec.	44.73	6.83	44.88	6.87
V	H ₂ NC ₆ H ₄ NHCONHCH ₂ CH ₂ NMe ₂ ·Cl·HCl	186	46.59	7.17	46.59	7.40
VI	H ₂ NC ₆ H ₄ CH ₂ CONHCH ₂ CH ₂ NMe ₂ ·2HCl	209.5–210.5	48.96	7.20	49.12	7.13
VII	H ₂ NC ₆ H ₄ CH ₂ CONHCH ₂ CH ₂ NMe ₂ ·Cl·HCl	155–156 dec.	50.63	7.52	50.49	7.61
VIII	C ₆ H ₅ NHCSNHCH ₂ CH ₂ NMe ₂	83–83.5	59.15	7.68	58.97	7.69
IX	H ₂ NC ₆ H ₄ SO ₂ NHCH ₂ CH ₂ NMe ₂ ·2HCl	211.5–213 dec.	37.96	6.06	37.86	6.02

aminophenylacetamidoethyl]-trimethylammonium chloride hydrochloride (VII).

With phenyl isothiocyanate the diamine formed *N*-phenyl-*N'*- β -dimethylaminoethyl thiourea (VIII) and with *p*-acetamidobenzenesulfonyl chloride it formed the *p*-acetamidobenzenesulfonamide which was hydrolyzed with hydrochloric acid to β -[*p*-aminophenylsulfonamidoethyl]-dimethylamine dihydrochloride (IX). The thiourea was

Some N-Substituted Barbituric Acids

The subjoined table contains data on five new compounds of this type. 1-*p*-Nitrophenyl-5-*i*-butyl-5-ethyl barbituric acid was obtained by nitration¹ of 1-phenyl-5-*i*-butyl-5-ethyl barbituric acid² and in turn was reduced catalytically¹ to the *p*-amino derivative. The other three substances were prepared by the conventional method from the corresponding ureas and malonic esters. All

DERIVATIVES OF BARBITURIC ACID R,R' $\begin{matrix} \diagup \text{CO-NR}'' \\ \diagdown \text{CO-NH} \end{matrix}$ CO

Substituents			Formula	M. p., °C.	Analyses, %					
R	R'	R''			Calcd.		Found			
					C	H	N	C	H	N
C ₂ H ₅	C ₂ H ₅	(<i>n</i>)C ₆ H ₁₃	C ₁₄ H ₂₄ O ₃ N ₂	41	62.64	9.02		62.79	9.09	
C ₂ H ₅	(<i>n</i>)C ₄ H ₉	(4)C ₂ H ₅ C ₆ H ₄	C ₁₈ H ₂₄ O ₃ N ₂	107	68.31	7.65		68.21	7.81	
C ₂ H ₅	(CH ₃) ₂ CHCH ₂	(4)H ₂ NC ₆ H ₄	C ₁₆ H ₂₁ O ₃ N ₃	153	63.33	6.98		63.58	7.32	
H	H	(2)C ₂ H ₅ OC ₆ H ₄	C ₁₂ H ₁₂ O ₄ N ₂	193.5			11.29			11.47
C ₂ H ₅	(CH ₃) ₂ CHCH ₂	(4)O ₂ NC ₆ H ₄	C ₁₆ H ₁₉ O ₅ N ₃	188			12.61			12.89

crystallized from benzene–hexane; the hydrochlorides from absolute alcohol.

crystallized in colorless prisms, the 1-*n*-hexyl derivative from hexane, the others from alcohol.

(1) Cf. Buck, *THIS JOURNAL*, **59**, 1249 (1937).

(2) Hjort and Dox, *J. Pharmacol.*, **35**, 155 (1929).

BURROUGHS WELLCOME & CO., U. S. A.
EXPERIMENTAL RESEARCH LABORATORIES
TUCKAHOE, NEW YORK

RICHARD BALTZLY
JOHANNES S. BUCK
WALTER S. IDE

BURROUGHS WELLCOME & CO., U. S. A.
EXPERIMENTAL RESEARCH LABORATORIES
TUCKAHOE, NEW YORK

JOHANNES S. BUCK
WALTER S. IDE
RICHARD BALTZLY

RECEIVED JUNE 6, 1942

RECEIVED JUNE 6, 1942