

We have found further that the additional carbon atom is present as a methoxyl group. Methoxyl determinations in this Laboratory on bethogenin were as follows: Calcd. for  $C_{28}H_{44}O_4$ , 6.97. Found: 6.68, 7.25, 6.82, 6.65. For bethogenin acetate, Calcd. for  $C_{30}H_{46}O_5$ , 6.37. Found, 6.15, 6.32. For bethogenin benzoate, Calcd. for  $C_{35}H_{48}O_5$ , 5.65. Found: 5.11.

Although bethogenin reacts with hydroxylamine in alcoholic pyridine solution<sup>1</sup> with the introduction of two nitrogen atoms, we find that bethogenin does not show typical carbonyl absorption in the ultraviolet. The absorption in alcohol rises almost regularly from  $\lambda$  3500 to  $\lambda$  2300 with only an inflection at  $\lambda$  2850,  $\log \epsilon$  at this point being about 0.3. After treatment with hydrogen bromide in acetic acid, however, a product is obtained which gives an absorption spectrum typical of the isolated carbonyl group;  $\lambda_{\max.} = 2850$ ,  $\log \epsilon = 1.77$ . This would indicate that bethogenin is an enol ether but we are not prepared to suggest a structure for bethogenin until work in progress has been completed.

DEPARTMENT OF CHEMISTRY  
STANFORD UNIVERSITY  
STANFORD UNIV., CALIF.

C. R. NOLLER  
M. R. BARUSCH

RECEIVED MAY 6, 1943

**THE SYNTHESIS OF CONDENSED RING COMPOUNDS. X. *cis*-9-METHYL-8-KETO-2-OCTALIN AND *cis*-10-METHYL-1-VINYL-1,7-NAPHTHITADIENE<sup>1</sup>**

Sir:


The addition of 1-methylcyclohexen-6-one to butadiene has yielded *cis*-9-methyl-8-keto-2-octalin (*cis*-10-methyl-7-naphthiten-1-one<sup>2</sup>), I, b. p. 67° at 0.5 mm.,  $n_D^{20}$  1.5042. Semicarbazone, m. p. 224.8–225° (dec., slight darkening at 224°). *Anal.* Calcd. for  $C_{12}H_{18}ON_3$ : N, 18.99; N found, 18.71, 18.93. In the presence of Adams catalyst, 0.94 millimole of the methyloctalone absorbed 1.88 millimoles of hydrogen.

The methyloctalone was converted by way of 5,10-*cis*-1-ethynyl-10-methyl-7-naphthiten-1-ol, b. p. 93–94° at 0.5 mm.,  $n_D^{20}$  1.5260, and 5,10-*cis*-10-methyl-1-vinyl-7-naphthiten-1-ol, b. p. 92° at 0.55 mm.,  $n_D^{20}$  1.5218, to *cis*-10-methyl-1-vinyl-1,7-naphthitadiene, II. The best reagent found so far for accomplishing the last step is

(1) This work was supported by an allotment from the Special Research Fund (Bankhead-Jones Act of June 29, 1935). Not subject to copyright.

(2) For nomenclature see Butz and Joshel, *THIS JOURNAL*, **64**, 1311 (1942); decahydronaphthalene becomes *naphthitane* and carbon atoms are numbered in sequence.

phosphorus tribromide and pyridine. II, b. p. 66–67° at 0.45 mm.,  $n_D^{20}$  1.5284, maximum absorption at 2380 Å.,  $\epsilon$  9400 in hexane and at 2380 Å.,  $\epsilon$  10,000 in ethanol.<sup>3</sup> Hydrogen absorbed by 0.37 millimole of triene, 26.2 ml. at standard temperature and pressure; calcd. for three double bonds, 26.75 ml.



*p*-Naphthoquinone was heated with two equivalents of the triene at 110° in a sealed tube for forty-five minutes. Addition of methanol to the reaction mixture gave crystals, m. p. 183–185°. This material was recrystallized from ethanol with Norite and then from ethanol containing a little petroleum ether; yellow crystals, m. p. 191.8–192.8°. *Anal.* Calcd. for  $C_{23}H_{24}O_2$ : C, 83.10; H, 7.28. Found: C, 82.87, 82.83; H, 7.12, 7.15. This compound may have the structure III.

We are applying these reactions to the synthesis of steroids.

(3) Obtained through the kindness of Mr. Harry Bastron. Compare with the maxima of related compounds in Booker, Evans and Gillam, *J. Chem. Soc.*, 1453 (1940).

BUREAU OF ANIMAL INDUSTRY WALTER NUDENBERG  
AGRICULTURAL RESEARCH ADMINISTRATION  
UNITED STATES DEPARTMENT OF AGRICULTURE  
BELTSVILLE, MARYLAND LEWIS W. BUTZ

RECEIVED JUNE 11, 1943

**ORIENTATION IN STRETCHED FILMS OF AMYLOSE TRIACETATE**

Sir:

Starch (corn, wheat, and potato) can be separated by either water extraction or butanol precipitation<sup>1</sup> into two components, amylose and amylopectin. Amylose is the component extracted by water from swollen starch granules or precipitated by butanol from an autoclaved starch paste while amylopectin is the starch component remaining after separation of the amylose. Recently Whistler and Hilbert<sup>2</sup> have shown that amylose is capable of yielding strong pliable acetate films of high quality whereas only weak

(1) T. J. Schoch, *THIS JOURNAL*, **64**, 2957 (1942).

(2) Roy L. Whistler and G. E. Hilbert, paper presented before the 105th meeting of the American Chemical Society, Detroit, Michigan, April, 1943, Division of Sugar Chemistry and Technology. Now in preparation for publication.

brittle films are produced from amylopectin acetate. The pliable amylose acetate films on stretching become birefringent and increase in tensile strength. This is evidence that a molecular orientation occurs within the film during stretching. The molecules constituting the film, therefore, most probably possess a highly anisodimensional or linear structure.

Confirmatory evidence that stretching produces orientation in these acetate films has been obtained by X-ray analysis. X-Ray diffraction patterns taken during the stretching of films clearly indicate a progressive change from an amorphous to a highly crystalline condition. Films which are stretched in hot water readily set to the new length when cooled and therefore can be easily handled in the elongated state. When elongated 400–600% the films produce a typical fiber pattern on exposure to X-rays (Fig. 1). The presence of discrete spots in the diffraction pattern is proof of the high degree of orientation attained by the film molecules and is further evidence that the molecules possess a linear nature. This is the first time a fiber pattern has been obtained with starch or its derivatives. Identical diffraction patterns are obtained from stretched acetate films prepared from either fractionated corn or potato starches. In each case a periodicity of 18.3 Å., occurs along the fiber (b) axis. While no experimental evidence was reported, French<sup>3</sup> claims that "oriented X-ray diffraction patterns from amylose films confirm previous unit cell determinations for granular (a-b) starches."

Although good patterns were obtained from un-plasticized films, best patterns were obtained from films containing about 20% plasticizer (tricresyl phosphate, dibutyl phthalate, dibutyl tartrate, pentaerythritol tetraacetate). The diffraction

(3) D. French, Doctor's Dissertation No. 696, Iowa State College, Ames, Iowa (1942); *Iowa State J. Science*, **17**, 62 (1942).




Fig. 1.—X-Ray pattern of stretched film of amylose triacetate (arrows indicate the direction of the fiber axis).

patterns seem to be independent of the type of plasticizer used. Patterns were obtained with flat film technique by allowing a beam of unfiltered Cu K $\alpha$  X-rays to strike the stretched film at an angle of 90° to the surface. The specimen was mounted on the pinhole and a specimen to photographic film distance of 5 cm. was used. All the stretched films showed a small birefringence which was negative with reference to the axis of elongation.

NORTHERN REGIONAL RESEARCH LABORATORY  
BUREAU OF AGRICULTURAL AND INDUSTRIAL CHEMISTRY  
AGRICULTURAL RESEARCH ADMINISTRATION  
U. S. DEPARTMENT OF AGRICULTURE ROY L. WHISTLER  
PEORIA, ILLINOIS N. CYRIL SCHIELTZ

RECEIVED MAY 27, 1943