

TABLE I
 POLYMETHYLENEDIAMINES

	R	B. p., °C. 1 mm.	Carbon, %		Hydrogen, %		Nitrogen, %	
			Calcd.	Found	Calcd.	Found	Calcd.	Found
H ₂ N(CH ₂) ₄ NR ₂	<i>n</i> -C ₂ H ₇	97-99	72.00	72.28	14.00	14.36
	<i>n</i> -C ₄ H ₁₁	137-138	75.00	75.03	14.06	14.35
H ₂ N(CH ₂) ₆ NR ₂	<i>n</i> -C ₂ H ₇	109-111	73.68	73.40	14.03	14.27
	<i>n</i> -C ₄ H ₉	131-132	10.83	10.62
	<i>n</i> -C ₆ H ₁₁	148-149	9.85	10.18
H ₂ N(CH ₂) ₁₀ NR ₂	<i>n</i> -C ₂ H ₇	136-137	75.00	74.93	14.06	14.25
	<i>n</i> -C ₄ H ₉	142-144	76.05	76.30	14.08	14.51	9.85	9.62

sium phthalimide by the conventional procedures.^{6,7} Hydrolysis of these compounds by hydrazine hydrate⁸ was more satisfactory than hydrolysis by concentrated hydrochloric acid.

The diamines are listed in Table I. They are colorless, viscous liquids, insoluble in water. They are hygroscopic and absorb carbon dioxide from the air. Efforts to prepare the usual solid derivatives yielded oils which would not crystallize. The hydrochlorides were too hygroscopic to be of value as derivatives.

(6) Muller and Kraus, *Monatsh.*, **61**, 219 (1932).

(7) Gabriel, *Ber.*, **22**, 1137 (1889).

(8) Ing and Manske, *J. Chem. Soc.*, 2350 (1926).

CHEMISTRY LABORATORY
UNIVERSITY OF MISSOURI
COLUMBIA, MISSOURI

H. E. FRENCH
H. E. UNGNADE
J. E. POE
L. H. EILERS

RECEIVED MARCH 14, 1945

Bis-(β -alkoxyethyl) Diselenides

Bis-(β -methoxyethyl) Diselenide.—A mixture of 28 g. of potassium sulfite and 9.5 g. of selenium in 300 ml. of 50% ethanol was digested under reflux for an hour. Four grams of anhydrous potassium carbonate and 11.3 ml. of methyl 2-bromoethyl ether in 12 ml. of ethanol were successively added and the refluxing was continued for two more hours. The mixture was acidified with excess 1 N

hydrochloric acid and then oxidized with 70 ml. of 3% hydrogen peroxide, added slowly with stirring. The diselenide separated out as an orange red oil with a strong alliaceous odor. The oil was washed with warm water, centrifuged and fractionated under reduced pressure; b. p. 124-126° (7 mm.), yield 10 g.

Properties.—Bis-(β -methoxyethyl) diselenide is a yellow, vesicant oil with alliaceous odor; i, H₂O, s. organic solvents.

Bis-(β -ethoxyethyl) diselenide and bis-(β -butoxyethyl) diselenide also have been prepared. Both are yellow oils of unpleasant odor and are skin irritants.

DATA ON BIS-(β -ALKOXYETHYL) DISELENIDES

Formula	C ₆ H ₁₄ O ₂ Se ₂			C ₈ H ₁₈ O ₂ Se ₂			C ₁₂ H ₂₆ O ₂ Se ₂			
	Calcd.	Found		Calcd.	Found		Calcd.	Found		
Analyses, %	Carbon	26.08	26.23	31.57	31.43	40.10	5.11	5.97	7.28	
	Hydrogen	5.11	5.11	5.97	5.89	7.39	124-126	135-137	175-177	
Boiling point	°C.	124-126	135-137	175-177	7	6	8	1.596	1.451	1.303
	Mm.	7	6	8	1.5554	1.5340	1.5150			
<i>d</i> ₄ ²⁵		1.596	1.451	1.303						
<i>n</i> _D ²⁰		1.5554	1.5340	1.5150						

CHEMICAL LABORATORY
CATHOLIC UNIVERSITY OF AMERICA
WASHINGTON, D. C.

H. P. WARD
I. LEO O'DONNELL¹

RECEIVED FEBRUARY 5, 1945

(1) Present address, St. Mary's College, Winona, Minnesota.

COMMUNICATIONS TO THE EDITOR

ORIENTED FIBERS OF SODIUM PECTATE

Sir:

Although brief mention has been made in several places concerning X-ray diffraction investigations of pectin or pectin derivatives, interpretation of the data obtained has not been attempted. This lack of information concerning the structure of this important natural polymer is partly due to the difficulty of getting X-ray diffraction photographs with sufficient reflections to enable a structural analysis to be made.

A short time ago it was noticed from X-ray powder photographs that sodium pectate is highly crystalline. Attempts were immediately made to produce sodium pectate fibers having molecular orientation. Best results were obtained by first making a pectic acid fiber and then converting it to the sodium salt. The pectic acid used was prepared in this Laboratory by the method of Baier

and Wilson¹ and had an intrinsic viscosity of 3.8.

A successful method for making well-oriented sodium pectate fibers consisted of titrating a 1% solution of pectic acid to a pH of 5.0 with sodium hydroxide. The resulting solution was then forced through a 1-mm. nozzle into a coagulating bath consisting of 85% ethyl alcohol in 1 N hydrochloric acid. The resulting wet fiber was strong enough to be handled readily. The hydrated fiber was held in a 60% alcoholic, 0.1 N sodium hydroxide solution overnight and then immersed for twenty-four hours in 60% ethyl alcohol. The fiber was then removed and slowly elongated 38% while drying. The dry fiber was strong, pliable, showed good molecular orientation, and was remarkably crystalline (Fig. 1).

The layer line spacings on the X-ray photograph can be measured readily. From these

(1) W. E. Baier and C. W. Wilson, *Ind. Eng. Chem.*, **33**, 287 (1941).