

## COMMUNICATIONS TO THE EDITOR

### METHIONINE AND ISOLEUCINE CONTENT OF MAMMALIAN HEMOGLOBINS. ITS SIGNIFICANCE FOR NUTRITIONAL AND METABOLIC STUDIES

Sirs:

Reports indicate that rats fail to grow when the only dietary protein is human or beef globin unless supplemented with isoleucine.<sup>1,2,3</sup>

Adult dogs receiving dog hemoglobin as virtually the sole source of protein maintain approximate nitrogen balance with slight loss of weight.<sup>4,5</sup> Dog hemoglobin supplemented with isoleucine is no better utilized.<sup>5</sup> However, supplementation with methionine results in positive nitrogen balance and maintenance of, or slight gain in, weight.<sup>5</sup> Further addition of isoleucine does not improve the effect of methionine.<sup>4,5</sup>

The nutritive value of "hemoglobin" seems, therefore, to depend on experimental conditions. It will be shown, however, that the results are exactly what could be expected from the difference in isoleucine and methionine content of hemoglobins.

From the data in the accompanying table,

#### ISOLEUCINE AND METHIONINE CONTENT OF MAMMALIAN HEMOGLOBINS

	Assumed Molecular Weight = 66,700					
	Isoleucine <sup>6</sup>			Methionine <sup>7</sup>		
	Protein, %	M/10 <sup>3</sup> g.	M/m	Protein, %	M/10 <sup>3</sup> g.	M/m
Human <sup>8</sup>	0	0	0	1.32	8.85	5.9
Dog <sup>7,9</sup>	1.36	10.4	6.9	0.42	2.81	1.9
Horse <sup>10</sup>	0 <sup>10</sup>	0	0			4 <sup>12</sup>
Bovine, adult <sup>13</sup>	0	0	0	1.76	11.8	7.9
Bovine, fetal <sup>14</sup>	0.63	4.80	3.2	0.97	6.50	4.3

it can be seen that the isoleucine content of dog hemoglobin is appreciable (1.36%) whereas human, horse and adult bovine hemoglobins lack this amino acid.<sup>14</sup>

Methionine is present in all hemoglobins studied, to the extent of 1% or higher, except in dog hemoglobin (only 0.4%).

The amino acid requirements for hemoglobin formation would seem to be different in different

(1) Devline and Zittle, *J. Biol. Chem.*, **156**, 393 (1944).

(2) Albanese, *ibid.*, **157**, 613 (1945).

(3) Orten, Bourque and Orten, *ibid.*, **160**, 435 (1945).

(4) Whipple, Robschelt-Robbins and Miller, *Ann. N. Y. Acad. Sci.*, in press.

(5) Miller, *Federation Proc.*, **5**, 73 (1946).

(6) Microbioassay with *L. arabinosus* 17-5 and with *L. mesenteroides* P-60, cf. Brand, *et al.*, *This Journal*, **67**, 1524 (1945).

(7) Microbioassay with *L. mesenteroides*.

(8) Prepared by R. K. Cannan and obtained from H. B. Vickery.

(9) Prepared by L. L. Miller.

(10) Prepared by G. L. Foster (*J. Biol. Chem.*, **159**, 431 (1945)).

(11) Microbioassay with *L. arabinosus*.

(12) Brand and Kassell, Abstracts, Am. Chem. Soc., Boston meeting, Sept. 1939.

(13) Prepared by Wyman, *et al.*, *J. Biol. Chem.*, **153**, 275 (1944), and obtained from H. B. Vickery, *ibid.*, **156**, 283 (1944).

(14) Devline and Zittle<sup>1</sup> and Albanese<sup>2</sup> report an isoleucine content of 0.29 and 0.8%, respectively, in human hemoglobin.

mammals. Whereas the human, the horse and cattle need for this purpose appreciable amounts of methionine but no isoleucine, the dog on the other hand needs isoleucine and only small amounts of methionine. The amino acid composition of hemoglobin is of metabolic significance since its breakdown contributes appreciably to the protein economy of an animal.

That fetal and adult bovine hemoglobins are two different proteins is again<sup>13</sup> illustrated by the difference in their methionine and isoleucine content.

This work was carried out in part under a contract recommended by the Committee on Medical Research, between the Office of Scientific Research and Development and Columbia University.

DEPARTMENT OF BIOCHEMISTRY  
COLLEGE OF PHYSICIANS AND SURGEONS  
COLUMBIA UNIVERSITY  
NEW YORK, N. Y.


ERWIN BRAND  
JEAN GRANTHAM

RECEIVED MARCH 16, 1946

### A NEW ALDEHYDE SYNTHESIS

Sir:

In connection with our previous study on the hydrogenolysis of thioacetals,<sup>1</sup> it was apparent that the reductive desulfurization reaction therein described might be extended to compounds such as thiol esters in which case the corresponding aldehyde would be obtained.



This would be of particular interest in the sugar series since it would constitute a new method for synthesizing aldehyde derivatives. We have established the above reaction by the hydrogenolysis of the thiol esters of benzoic, propionic and tetraacetyl-D-ribonic acids which gave benzaldehyde, propionaldehyde and *aldehydo-D-ribose* tetraacetate, respectively.

The thiol esters were prepared by the reaction of the acid chloride with an excess of ethyl mercaptan in pyridine or by reaction of the acid chloride with lead mercaptide in dry ether (as for ethyl thiolpropionate, *b. p.* 133-135°). The hydrogenolyses were carried out with Raney nickel in 70-80% ethanol under reflux essentially as described<sup>1</sup> for the hydrogenolysis of thioacetals. After catalyst removal, benzaldehyde and propionaldehyde were isolated in the distillate as their bisulfite compounds in 60-70% yield and identified by definitive derivatives. The residual sirup obtained on hydrogenolysis of ethyl thiol-

(1) M. L. Wolfrom and J. V. Karabinos, *This Journal*, **66**, 909 (1944).

D-ribonate tetraacetate (m. p. 87–87.5°,  $[\alpha]^{27}_D +17^\circ$  in chloroform), after removal of catalyst and solvent, was crystallized from acetone-methanol-(petroleum ether) and yielded *aldehydo-D-ribose tetraacetate*<sup>2</sup> in rather low yield.

Full details will be reported in a latter communication.

(2) R. Pasternack and E. V. Brown, U. S. Patent 2,237,263 (1941).

CHEMICAL LABORATORY  
THE OHIO STATE UNIVERSITY  
COLUMBUS, OHIO

M. L. WOLFROM  
J. V. KARABINOS


RECEIVED FEBRUARY 25, 1946

### PRESSURE AND SPECIFIC PRECIPITATION

Sir:

Although the reactions that lead to specific precipitation evidently involve a combination between a di- or polyvalent antigen or hapten and specific sites on the antibody protein, probably resulting in the formation of a framework comprising large aggregates of the two species of molecules,<sup>1</sup> convincing evidence is lacking as to whether the protein itself becomes significantly altered in the process. The following observations bear on this question and suggest an approach for further study.

The precipitation system consisted of serum from rabbits immunized against arsanic-azosheep serum and a simple trihaptenic dye antigen, *viz.*



To slow the precipitation for practical purposes, the serum was diluted 1:2, and the antigen 1:160,000. Visible precipitation occurred within a few minutes, and went to about three-fourths completion in an hour at room temperature. Small test-tubes were filled with corresponding specimens, stoppered with a rubber stopper, then placed in a water-filled steel pressure chamber, and hydrostatic pressure applied from a hydraulic pump within one to two minutes after mixing. At 10,000 pounds per square inch a scarcely visible precipitate formed during slightly more than an hour. After the pressure was released, precipitation continued in apparently the normal manner. Lower pressures were less effective, but quantitative analyses of protein nitrogen in precipitates centrifuged within less than five minutes after releasing pressure, showed decreases ranging from 55 to 77% under pressures between 1,500 and 8,000 pounds, in comparison with the amount under normal pressure. These

(1) Pauling, Campbell and Pressman, *Physiol. Rev.*, **23**, 203 (1943).

data indicate a molecular volume increase on the order of 50 cc. or more per mole, although an accurate value requires detailed data on the rate of the reaction at different pressures.

Pressures of this magnitude oppose the reversible and irreversible denaturations of certain enzyme systems<sup>2</sup> and greatly retard the denaturation of purified human serum globulin at 65°.<sup>3</sup> These pressure effects are considerably greater than those encountered with reactions of small molecules and indicate that a number of groups are involved, or perhaps the protein molecule as a whole. They suggest that changes as extensive as those which take place in denaturation occur also in specific precipitation. An extensive quantitative study of pressure effects on the rate of specific precipitation will perhaps yield cogent data regarding the mechanism of the reaction.

THE GATES AND CRELLIN DAN H. CAMPBELL  
LABORATORIES OF CHEMISTRY FRANK H. JOHNSON<sup>4</sup>  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
PASADENA, CALIF.

RECEIVED FEBRUARY 6, 1946

(2) Johnson, Eyring, Steblay, Chaplin, Huber and Gherardi, *J. Gen. Physiol.*, **28**, 463 (1945).

(3) Johnson and Campbell, *J. Cell. Comp. Physiol.*, **26**, 43 (1945).

(4) Guggenheim Fellow.

### METABOLIC PRODUCTS OF *ASPERGILLUS USTUS*

Sir:

Publication of preliminary results by Hogeboom and Craig<sup>1</sup> on the application of the latter's

countercurrent distribution method to the problem of isolating bacteriologically active components from the culture filtrate of *Aspergillus Ustus*<sup>2</sup> prompts us to report briefly the progress of our work carried out in collaboration with Mr. Joseph Kurung and Dr. Harry Bray, Superintendent of the New York State Hospital for Incipient Tuberculosis.

The crude active material,<sup>3</sup> obtained by ether extraction of the culture filtrate or mycelium, has been divided into a bicarbonate soluble, a carbonate soluble and a neutral fraction. By fractional crystallization three substances have been isolated from the neutral fraction in small, variable amounts, and never all three from a single sample of crude material: A.U.N.-1, m. p. 155–6°, (C, 38.24; H, 4.10; Cl, 23.87); A.U.N.-2, m. p. about 270° with severe decomposition, inadequately characterized at present; A.U.N.-3,

(1) Hogeboom and Craig, *J. Biol. Chem.*, **162**, 363 (1946).

(2) Kurung, *Science*, **102**, 11 (1945).

(3) Generously supplied to us by Mr. Joseph Kurung and the Wallerstein Company, Inc.

m. p. 225.5–226.5°, (C, 62.59; H, 4.81; Cl, 8.59).

The carbonate soluble fraction consistently has given the predominant active component which we are naming *ustin*. This material may be purified by fractional crystallization from toluene, acetic acid, ether, or ether-hexane, m. p. 184–186°. The probable empirical formula of *ustin*,  $C_{19}H_{16}O_5Cl_3$ , is consistent with the analytical results (C, 53.42; H, 3.69; Cl, 24.27), conductometric titration (N.E. 214), and the elementary composition of the monomethyl derivative, m. p. 174° (C, 54.09; H, 3.95; Cl, 23.54;  $CH_3O$ , 7.01), the dimethyl derivative, m. p. 147° (C, 55.00; H, 4.24; Cl, 23.19;  $CH_3O$ , 13.35), and the acetyl derivative, m. p. 212° (C, 53.68; H, 3.95; Cl, 22.76;  $CH_3O$ , 9.67). Compound I isolated by Hogeboom and Craig,<sup>1</sup> m. p. 185–187°, is probably identical with *ustin*<sup>4</sup> although the large discrepancy between our chlorine analysis and that obtained by Hogeboom and Craig (Cl, 22.63) suggests that their compound is impure even though isolated by the elegant counter-current distribution method.

The antimicrobial activity of *ustin*, A.U.N.-1, and A.U.N.-2 are indistinguishable by any of the *in vitro* tests used so far. In synthetic media, the compounds inhibit the growth of Gram positive cocci and mycobacteria (including pathogenic tubercle bacilli) to the same degree, but are inactive against Gram negative bacilli. Their anti-septic activity increases rapidly with the  $H^+$  concentration of the environment, but is sharply inhibited by the addition of serum albumin, and of certain other organic substances (lipids for example) to synthetic media. Thus any one of the compounds can exert an inhibitory effect on the growth of tubercle bacilli, streptococci or staphylococci in a dilution of 1/500,000 at pH 6.5 in a simple synthetic medium, but will fail to retard growth even in 1/50,000 in more complex media at slightly alkaline reactions.

We wish to express our sincere appreciation to Dr. Adalbert Elek for the excellence of his micro-analytical assistance.

The details of this work will be published shortly.

(4) Dr. Craig very kindly examined the melting points and found no depression (m. p. and mixed m. p. 186–188°).

DEPARTMENT OF CHEMISTRY  
COLUMBIA UNIVERSITY AND  
DEPARTMENT OF BACTERIOLOGY  
ROCKEFELLER INSTITUTE  
NEW YORK, NEW YORK

W. E. DORRING  
R. J. DUBOS  
D. S. NOYCE  
R. DREYFUS

RECEIVED MARCH 8, 1946

#### SURFACE FILM THICKNESSES OF POLYMERIC SUBSTANCES IN RELATION TO MOLECULAR STRUCTURE<sup>1</sup>

Sir:

It is generally well known that certain com-

(1) The work reported in this letter was done in connection with the Government research program on synthetic rubber under contract with the Office of Rubber Reserve, Reconstruction Finance Corporation (results first reported in November, 1945).

pounds, such as long chain fatty acids, can form monomolecular films on water and that the areas of such films can be related to the dimensions of the molecules.<sup>2</sup> Certain polymeric substances including cellulose derivatives,<sup>3</sup> proteins<sup>4</sup> and the polymer of  $\omega$ -hydroxydecanoic acid<sup>5</sup> likewise form stable films whose areas can be conveniently measured.<sup>6</sup> At this time we wish to report that we have prepared films of some rubber-like polymers as well and that the results appear to give useful information about their molecular structures, particularly in relation to branching.

Five ml. of a very dilute benzene solution of a polymer ( $c = 2 \times 10^{-6}$  g./ml.) is distributed over the water surface of a hydrophil balance. After the benzene has evaporated, the film area is measured in the usual manner.<sup>7</sup> Low conversion samples of GR-A (butadiene-acrylonitrile copolymer) gave films about 6 Å. thick whereas high conversion samples exhibited thicknesses as great as 30 Å. It is our belief that a large film thickness indicates a high degree of branching and that the corresponding polymer molecules can be better represented as "bottle brushes" than as straight chains. Other polymers containing hydrophilic groups have also been investigated; these include copolymers of butadiene with methacrylonitrile, 1-cyanobutadiene, *m*-fluorostyrene and some vinylpyridines.

Stable films for polymers such as natural rubber or GR-S could not be obtained directly. However, these substances can be investigated by the film technique if they are first rendered hydrophilic by the addition of thiocyanogen to some of the double bonds according to the methods of Pummerer and Stark and of Rehner.<sup>8</sup> After using their procedures, we found natural rubber and gutta percha to form relatively thin films (about 8 Å. thick) whereas the film thicknesses of GR-S exhibited wide variations with polymer conversion, indicating once more marked differences in branching.

On the whole, the method appears to show considerable promise as a tool for learning more about the structure of polymers.

NOYES CHEMICAL LABORATORY  
UNIVERSITY OF ILLINOIS  
URBANA, ILLINOIS

FREDERICK T. WALL  
MURRAY ZELIKOFF

RECEIVED FEBRUARY 18, 1946

(2) Harkins, Davis and Clark, *THIS JOURNAL*, **39**, 541 (1917); Langmuir, *ibid.*, **39**, 1848 (1917).

(3) Katz and Samwel, *Ann.*, **472**, 241; **474**, 296 (1929); Adam, *Trans. Faraday Soc.*, **29**, 90 (1933); Harding and Adam, *ibid.*, **29**, 837 (1933).

(4) Hughes and Rideal, *Proc. Roy. Soc. (London)*, **A137**, 62 (1932).

(5) Harkins, Ries and Carman, *J. Chem. Phys.*, **3**, 692 (1935).

(6) The authors are indebted to Dr. Anderson Pace, Rayon Department, E. I. du Pont de Nemours and Co., Buffalo 7, N. Y., for calling to their attention the possibility of such measurements.

(7) See for example, Adam, "The Physics and Chemistry of Surfaces," Oxford University Press, London, 1941, 3rd ed., p. 28.

(8) Pummerer and Stark, *Ber.*, **64**, 825 (1931); Rehner, *Ind. Eng. Chem.*, **36**, 118 (1944).