

to be of the same order of activity as the authentic material.

LABORATORY OF ORGANIC CHEMISTRY  
UNIVERSITY OF WISCONSIN  
MADISON, WISCONSIN

WILLIAM S. JOHNSON  
ROBERT P. GRABER



RECEIVED JUNE 14, 1948

### STREPTOMYCES ANTIBIOTICS. XIX. DIHYDRO-STREPTOSONIC ACID LACTONE AND CONFIGURATION OF STREPTOSE AND STREPTOBIOSAMINE

Sir:

2-Methyl pentaacetyldihydrostreptobiosaminide<sup>1</sup> was allowed to react stepwise with ethyl mercaptan-hydrogen chloride, acetic anhydride, and mercuric chloride for the preparation of amorphous pentaacetyldihydrostreptobiosamine.<sup>2</sup> Acetylation of this compound gave hexaacetyldihydrostreptobiosamine.<sup>2</sup> Oxidation by bromine and hydrolysis by hydrochloric acid of pentaacetyldihydrostreptobiosamine gave the known N-methyl-L-glucosamine and the new dihydrostreptosonic acid lactone, m. p. 143-144°,  $[\alpha]_D -32^\circ$  (c, 0.40 in water). Reaction of the lactone with hydrazine gave dihydrostreptosonic acid hydrazide, m. p. 137-139°,  $[\alpha]_D +23^\circ$  (c, 0.9 in water).

Application of Hudson's rules of rotation to streptosonic acid diamide<sup>3</sup> and dihydrostreptosonic acid hydrazide shows that the hydroxyl group at C<sub>2</sub> of streptose lies on the right. Since it has already been shown<sup>4</sup> that the hydroxyl groups at C<sub>2</sub> and C<sub>3</sub> of streptose are *cis*, and that the configuration about C<sub>4</sub> is *levo*,<sup>5</sup> the configuration of L-streptose is represented by structure I. On the basis of these data, and the calculations of the glycosidic linkage between streptose and N-



(1) Brink, Kuehl, Flynn and Folkers, *THIS JOURNAL*, **68**, 2557 (1946).

(2) Stavely, Wintersteiner, Fried, White and Moore, *ibid.*, **69**, 2742 (1947).

(3) Kuehl, Flynn, Brink and Folkers, *ibid.*, **68**, 2679 (1946).

(4) Brink, Kuehl, Flynn and Folkers, *ibid.*, **68**, 2405 (1945).

(5) Fried, Walz, and Wintersteiner, *ibid.*, **68**, 2746 (1946).

methyl-L-glucosamine to be  $\alpha$ -L,<sup>6</sup> the configuration of streptobiosamine is represented by structure II. The levorotations of streptosonic acid lactone<sup>3</sup> and dihydrostreptosonic acid lactone support the applicability of Hudson's rules to these streptose derivatives, since it is established conclusively that the configuration about C<sub>4</sub> of these lactones is L. That the lactone of dihydrostreptosonic acid lactone involves the secondary hydroxyl group at C<sub>4</sub> is shown by the liberation of formaldehyde when the lactone reacts with two equivalents of periodic acid.

(6) Lemieux, DeWalt and Wolfrom, *ibid.*, **69**, 1838 (1947).

RESEARCH DEPARTMENT  
MERCCK AND CO., INC.  
RAHWAY, N. J.

FREDERICK A. KUEHL, JR.  
MARY NEALE BISHOP  
EDWIN H. FLYNN  
KARL FOLKERS

RECEIVED MAY 27, 1948

### CHARACTERISTICS OF THE DROPPING MERCURY ELECTRODE IN FUSED SALTS

Sir:

In a preliminary investigation of the applicability of polarographic techniques to fused salt media we have obtained typical polarographic reduction waves for the cations of a number of salts dissolved in a fused salt solvent. The results indicate that the Ilkovic equation<sup>1,2</sup> is applicable to the melt employed, a ternary eutectic consisting of 66.65 mole % ammonium nitrate, 25.76% lithium nitrate, and 7.59% ammonium chloride (m. p. 86.2°).<sup>3</sup>

Mercury was used for the dropping electrode and the stationary unpolarized anode pool in a cell maintained at  $125 \pm 0.5^\circ$  in an oil-bath. Drops were collected in a Pyrex spoon, washed, dried, and weighed for tests of the Ilkovic equation.

Characteristic reduction waves were obtained with nickel(II), copper(II), and bismuth(III), the latter two exhibiting maxima. A trace of potassium iodide eliminated the maximum in the case of copper. Varying degrees of success have been had with other solute salts, prime difficulties being limited solubility in or reaction with the solvent electrolyte.

TABLE I  
TEST OF THE ILKOVIC EQUATION

No.	C, mmol./l.	$i_d$ , $\mu$ amp.	$m$ , mg./sec.	$i_{max.}$ , sec.	$\frac{i_d}{Cm^2/t^{1/2}}_{max}$
1	1.95	3.74	1.41	4.0	1.21
2	4.98	10.0	1.45	4.5	1.22
3	6.77	12.8	1.37	3.9	1.22
4	9.97	17.2	1.45	3.4	1.10
5	12.8	21.6	1.40	3.5	1.09
6	12.8	16.6	0.658	7.6	1.22

(1) D. Ilkovic, *Coll. Czechoslov. Chem. Commun.*, **6**, 498 (1934).

(2) I. M. Kolthoff and J. J. Lingane, "Polarography," Interscience Publishers, Inc., New York, N. Y., 1941, p. 88.

(3) E. P. Perman and R. E. Wilson, *J. Chem. Soc.*, **125**, 1700 (1924).