

For the case where the rate of exchange is measured by the growth of activity, y , in B

$$y(t) = y(\infty) [1 - \exp(-s)] \quad (4)$$

$$d \ln s/d \ln (y/y(\infty)) = (\exp(s) - 1)/s = F(s) \quad (5)$$

The figure contains a plot of this function, too. In this case, of course, the fractional error in y cannot possibly be constant as y approaches zero; an estimate of σ_y as a function of y is needed for selecting an optimum reaction time or calculating the error of a particular determination.

CONTRIBUTION NO. 1233 FROM THE
GATES AND CRELLIN LABORATORIES OF CHEMISTRY
CALIFORNIA INSTITUTE OF TECHNOLOGY
PASADENA 4, CALIFORNIA RECEIVED AUGUST 9, 1948

A Convenient Synthesis of *N,N*-Dimethyl-*p*-nitroaniline and *N,N*-Dimethyl-*o*-nitroaniline

BY TOD W. CAMPBELL

There are a number of methods in the literature for the preparation of *N,N*-dimethylnitroaniline¹; but most of them are troublesome and result in low yields of a product which is usually of not too high purity. The methods in the literature^{1,g,i} which give acceptable or good yields involve a reaction in which a nitrohalobenzene is allowed to react with dimethylamine.

The author has found that this reaction is most conveniently brought about by refluxing a pyridine solution of nitrohalobenzene with a mixture of dimethylamine hydrochloride and sodium bicarbonate. The desired product is obtained in virtually quantitative yield; in the case of the para isomer, the product can be crystallized directly from the reaction solvent in a high state of purity. This procedure is therefore recommended for the preparation of these two substances.

Experimental Part

***p*-Nitrodimethylaniline.**—A mixture of 42 g. of *p*-bromonitrobenzene, 300 cc. of pyridine, and 50 g. of sodium bicarbonate was placed in a 500-cc. round-bottom flask. To this mixture was added 30 g. of dimethylamine hydrochloride dissolved in about 10 cc. of warm water. The mixture was refluxed for ten hours. Mechanical stirring was not employed, since serious bumping did not occur. At the end of the reflux period, the hot solution was filtered free of inorganic salts, and the latter was extracted with 200 cc. of acetone, which was added to the pyridine solution. The mixed extracts were boiled, and water added to near the cloud point. On cooling, bright yellow needles of *p*-nitrodimethylaniline, 1–3 cm. in length, crystallized out. The melting point was observed to be 163.7–164.1° (lit. 163–166°) on a calibrated Anschütz thermometer in a Hershberg apparatus. The mother liquor on concentration to one third of its original volume gave an additional small yield of fine yellow needles, which had a melting point anywhere from 1–10 degrees low, for various experiments. One recrystallization from methanol raised the

(1) (a) Beilstein "Handbuch," Vol. XII, 690, 714, and first Supplement; (b) Le Fevre, *J. Chem. Soc.*, 147 (1930); (c) Davies, *Bull. soc. chim.*, [5] 2, 295 (1935); (d) Donald and Reade, *J. Chem. Soc.*, 53 (1935); (e) Marsden and Sutton, *ibid.*, 599 (1936); (f) Shorygin, Topchier and Anan'ina, *J. Gen. Chem. (U. S. S. R.)*, 8, 981 (1938); (g) Hodgson and Kershaw, *J. Chem. Soc.*, 280 (1930); (h) Evans and Williams, *ibid.*, 1199 (1939); (i) Senear, Rapport, Mead, Maynard and Koepfli, *J. Org. Chem.*, 11, 378 (1946).

melting point to 163.5–164°. The over-all yield of pure product was 32.4–33.6 g. (94–97%).

***o*-Nitrodimethylaniline.**—The above procedure was employed to prepare the ortho substituted derivative. Ten grams of *o*-nitrochlorobenzene gave 8.9 g. (85%) of product; b. p. 149 at 20 mm.; n_D^{25} 1.6080.

Anal. Calcd. for $C_8H_{10}N_2O_2$; C, 57.81; H, 6.06. Found: C, 57.53; H, 6.21.

UNIVERSITY OF CALIFORNIA
LOS ANGELES 24, CALIF. RECEIVED AUGUST 19, 1948

Vapor Density of Diborane¹

BY E. M. CARR, J. T. CLARKE AND H. L. JOHNSTON

In connection with the process of adding diborane to a calorimeter it was necessary to determine its density at 275.16°K. The diborane was obtained from the Naval Research Laboratory, subjected to a two-plate distillation and shown by the cryoscopic method to have a purity of 99.95 mole per cent. The diborane was introduced into an evacuated, weighed and calibrated 1-liter Pyrex bulb immersed in a constant temperature water-bath at 275.16°K. and constant to 0.01° C. during the measurement. The pressure was read on a 15 mm. i. d. manometer using a Gaertner cathetometer and a standard meter bar in an insulated case; the readings were converted to standard conditions and meniscus corrections made according to Cawood and Patterson.² The bulb was then reweighed (using a similar bulb as tare) on a Troemner 4-kg. balance. One bulb had 1/8 inch Pyrex helices with a surface area 16.0 times that of the interior part of the bulb added to it. Since all density measurements were made at approximately atmospheric pressure the amount of adsorption was assumed to be constant.

The results were used to calculate the density and the second virial coefficient B in the equation

$$n - A = \frac{PV}{RT + PV}$$

where A equals moles of diborane adsorbed on the surface of a 1-liter bulb and was found to have a value of 2.4×10^{-5} mole. A summary of the data is

Pressure, atm., P	Temperature, °K., T	Moles in gas phase, $n - A$	Volume of bulb, ml., V	Second virial coef. B (ml.), 275.16°K.	Density g./l. 1 atm. 275.16°K.
0.98832	275.20	0.048232	1090.2	-247	1.2398
.98201	275.14	.047296	1076.4	-233	1.2393
.99445	275.16	.048488	1090.2	-223	1.2386
.94553	275.16	.046104	1090.2	-234	1.2393
.96219	275.17	.046314	1076.4	-227	1.2388
.97098	275.15	.046690	1076.4	-199	1.2374
Average				-227	1.2389
Av. deviation				±11	±0.0006

(1) This work was carried out under contract between the Office of Naval Research and The Ohio State University Research Foundation.

(2) Cawood and Patterson, *Trans. Faraday Soc.*, 29, 514–523 (1933).