

for the gas observed with caproyl chloride but not with benzoyl or anisoyl chlorides.

Reaction of Trimethylacetyl Chloride with Diazomethane.—A solution of 24.1 g. (0.2 mole) of this acid chloride in 50 cc. of ether was added during eighty-five minutes to a cold solution of 20.4 g. (0.2 mole) of triethylamine and 8.5 g. of diazomethane in 275 cc. of ether. During the addition 0.022 mole (11%) of nitrogen was evolved, but on standing for twelve hours in the cold, no more nitrogen was liberated. The amine hydrochloride (96%) and crude diazoketone (98%) were isolated as described above. On treatment with acid, the crude diazoketone yielded 90% of nitrogen.

CHEMISTRY LABORATORY OF
THE OHIO STATE UNIVERSITY
COLUMBUS 10, OHIO

RECEIVED DECEMBER 13, 1948

Reduction of Anthracene and Phenanthrene with Lithium Aluminum Hydride

By JOHN R. SAMPEY AND JESSIE M. COX

Nystrom and Brown¹ have found lithium aluminum hydride a useful reagent for a number of organic reductions. No work has been reported on the reduction of aromatic or condensed ring nuclei with this reagent. Preliminary attempts to reduce naphthalene and anthracene in ether and dioxane solutions were unsuccessful, but when the solid lithium aluminum hydride was shaken with molten anthracene and phenanthrene, reduction did take place, the former giving as high as 60% yields of tetrahydroanthracene, and the latter 22% of dihydrophenanthrene. These reductions take place even though the decomposition point of lithium aluminum hydride has been reported at 125–150°. In all our reductions at high temperatures we mixed the powdered reagent thoroughly with the organic substances, and then employed vigorous shaking of the molten materials to ensure intimate mixing.

Method of Reduction.—A series of experiments is reported in which the temperature, the time and the ratio of reducing agent to anthracene and phenanthrene are varied. The experiments were run in a 200-ml. round-bottom flask, fitted with a calcium chloride tube, Bunsen valve and thermometer. The flask is heated with a Gyco heating jacket placed on a shaking machine. In each experiment 0.05 mole of the organic molecule was used. The molten material was shaken for the first two hours of each run, and then for fifteen minutes out of every thirty minutes; at the end of the time specified the hydride is decomposed by the slow addition of water. The reaction is vigorous and often flashes of fire were seen in the flask, but none of the reactions has gotten out of hand to date. After filtering and drying, the anthracene and its reduced form were extracted from the lithium and aluminum compounds with 200 ml. of hot chloroform and the phenanthrene with 200 ml. of ether or 95% alcohol. The unreduced hydrocarbons are less soluble than the hydrogenated products in the solvents used for the extractions, thus providing an effective method of separation. The tetrahydroanthracene was identified by a mixed melting point with a sample prepared by reduction of anthracene with Raney nickel (m. p. 103–104°). The dihydro-

(1) Nystrom and Brown, *THIS JOURNAL*, **69**, 1197, 2548 (1947); **70**, 441 (1948).

(2) Finholt, Bond and Schlesinger, *ibid.*, **69**, 1199–1203 (1947).

phenanthrene and its picrate were identical with those prepared by Schmidt.³

Results of Reduction.—A 60% yield of tetrahydroanthracene was obtained by heating 0.05 mole of anthracene at 220–230° for six hours with 8 g. of lithium aluminum hydride; a 45% yield was obtained by cutting the hydride to one-half. Increasing the temperature to 240–270° or lengthening the time of heating also reduced the yield. A 22% yield of dihydrophenanthrene was obtained by heating 0.05 mole of phenanthrene at 220–230° for six hours with 8 g. of lithium aluminum hydroxide. Reducing the temperature to 180–200° and the quantity of hydride, reduced the yield to 10–17%.

Acknowledgment.—The authors acknowledge the interest of Dr. E. Emmet Reid in this research and the assistance of a grant from the Office of Naval Research.

(3) Schmidt, *Ber.*, **40**, 4240–4257 (1907).

DEPARTMENT OF CHEMISTRY
FURMAN UNIVERSITY
GREENVILLE, S. C.

RECEIVED DECEMBER 8, 1948

Electroreparation of Technetium from Rhenium and Molybdenum

By L. B. ROGERS¹

Heretofore, technetium has been plated from acidic solutions^{2,3,4} under nearly the same conditions used for the deposition of rhenium. It seems quite likely, therefore, that the separation from rhenium would be poor. During the course of a polarographic study of technetium with G. E. Boyd⁵ it was found that distinct reduction waves were found for pertechnetate in 2 *M* sodium hydroxide whereas perrhenate and molybdate did not reduce, at the platinum electrode, up to the potential at which hydrogen was evolved rapidly.

The brown-black precipitate which appeared on the electrode was soluble in a mixture of ammonia and hydrogen peroxide indicating that a lower oxide of technetium was formed since technetium metal is insoluble.⁶ The height of the diffusion current confirmed the idea that the reduction step involved several electrons. Since the polarographic diffusion current reached a constant value at about -0.9 v. vs. the saturated calomel electrode (S.C.E.) and since the rapid evolution of hydrogen did not begin until about -1.2 v., the intermediate region of potential represented a promising one for the separation of technetium.

Experimental Results

A 100- μ l. sample of 2 *M* sodium hydroxide containing 0.2 mg. of technetium ($Tc^{99} - 9 \times 10^5$ years) as TcO_4^- plus a somewhat larger amount of molybdenum and a smaller amount of rhenium was placed upon a flat platinum foil used for the cathode. The anode was a large

(1) Present address: Massachusetts Institute of Technology, Cambridge 39, Mass.

(2) C. Perrier and E. Segrè, *J. Chem. Phys.*, **7**, 155 (1939).

(3) J. F. Flagg and W. E. Bleidner, *J. Chem. Phys.*, **13**, 269 (1945).

(4) E. E. Motta, Q. V. Larson and G. E. Boyd, *Plutonium Project Record*, MonC-99, April, 1947.

(5) G. E. Boyd and L. B. Rogers, reported by G. E. Boyd at the Symposium of the Division of Physical and Inorganic Chemistry at Syracuse, N. Y., June 28–30, 1948.

(6) G. E. Boyd, private communication.