

for the measurements using the No. 1554 Leeds and Northrup shielded capacitance and conductance bridge with a thousand cycle oscillator. The acid was purified by fractional crystallization, and the material had a melting point of  $-15.25 \pm 0.05^\circ$ . All manipulation of the pure sample was performed in a vacuum system and without contact of waxes and greases. The values obtained with a probable error of 2% are

Temp., °C.		$g \times 10^3 \text{ l./ohms}$
-11	26.2	1.28
-6.2	27.2	1.38
-2.3	28.7	1.49
0	29.5	1.55
12.8	36.1	1.68
14.2	36.0	1.72
21.5	40.2	2.03
27.7	43.4	2.33

The third column of the above table gives the determined conductances. The cell constant was approximately  $0.007 \text{ cm.}^{-1}$ . A slightly impure sample of the acid with a higher conductance was measured at  $0^\circ$ . The dielectric constant found was 26.9 and the conductance in the above units 1.69 showing that in the method of making the measurements an increase in conductivity resulted in a lower determined number for the dielectric constant. The same technique was employed using the same bridge and cell with nitrobenzene, and values were obtained which agreed with those recorded in the literature to better than 1%.

THE FLUORINE LABORATORIES  
THE PENNSYLVANIA STATE COLLEGE  
STATE COLLEGE, PENNSYLVANIA

J. H. SIMONS  
K. E. LORENTZEN

RECEIVED JANUARY 18, 1950

#### STUDIES ON LIGNIN AND RELATED PRODUCTS. II. 5-CARBOXYVANILIC ACID IN THE BISULFITE-SOLUBLE FRACTION FROM THE SILVER OXIDE OXIDATION OF FERMENTED SULFITE SPENT LIQUOR<sup>2</sup>

Sir:

In a recent paper on the silver oxide oxidation of basic calcium lignosulfonate<sup>1</sup> the isolation of substantial amounts of a 21% sodium bisulfite-soluble fraction which contained only small percentages of vanillin was noted. The bisulfite-soluble fraction from a similar silver oxide oxidation of a *Torulopsis utilis* fermented sulfite spent liquor has been found to contain at least 50% 5-carboxyvanillic acid.

The light brown crystalline bisulfite-soluble fraction (6.8 g.) was covered with 100 ml. of dry ether, heated to boiling, and filtered. The clear solution was treated with 50 ml. of dioxane and distilled slowly from a water-bath until all the ether was removed. After cooling and standing,

(1) For Part I of this series, see THIS JOURNAL, **71**, 2196 (1949).

(2) This paper represents a portion of the results obtained in the research program sponsored by the Sulphite Pulp Manufacturers' Research League and conducted for the League by The Institute of Paper Chemistry. Acknowledgment is made by the Institute for permission on the part of the League to publish these results.

the crystals were filtered, washed with cold dioxane, and dried. The yield of almost colorless crystals melting at  $276-277^\circ$  was 2.8 g. Concentration of the combined dioxane filtrate and washings yielded 0.6 g. of similar crystals. Repeated recrystallization from water raised the melting point to  $281^\circ$ . The product was identical with 5-carboxyvanillic acid reported by Freudenberg, Klink, Flickinger and Sobek,<sup>3</sup> who obtained it by partially hydrolyzing either 5-cyanoveratric acid or 5-carboxyveratric acid with hydrochloric acid.

*Anal.* Calcd. for  $\text{C}_9\text{H}_8\text{O}_6$ : C, 50.93; H, 3.80;  $\text{CH}_3\text{O}$ , 14.63. Found: C, 50.80; H, 3.88;  $\text{CH}_3\text{O}$ , 14.65.

The synthesis, degradation, derivatives, and significance to the structure of lignin of 5-carboxyvanillic acid will be discussed in forthcoming papers.

(3) Freudenberg, Klink, Flickinger and Sobek, *Ber.*, **72**, 224 (1939).

THE INSTITUTE OF PAPER CHEMISTRY  
APPLETON, WISCONSIN

IRWIN A. PEARL

RECEIVED JANUARY 16, 1950

#### ACID DEGRADATION OF AMYLOPECTIN TO ISOMALTOSE

Sir:

Considerable indirect evidence exists for C6 as the point of branching in amylopectin but it is desirable to place this on a more definitive basis through degradative experiments involving the isolation of crystalline products of known structure. Action of the  $\alpha$ - and  $\beta$ -amylases of malt on amylopectin does not yield isomaltose,<sup>1</sup> although levoglucosan is found in the hydrolyzate<sup>2</sup> (on subsequent treatment with the mold enzyme mentioned below). Isomaltose is isolable in crystalline form, and as the crystalline  $\beta$ -D-octaacetate, in the hydrolysis of amylopectin with an enzyme system of mold origin (a Takadiastase type from *Aspergillus oryzae*).<sup>3</sup> We report herein the isolation of isomaltose, as its crystalline  $\beta$ -D-octaacetate, from the acid hydrolyzate of amylopectin. Evidence is cited that this disaccharide is not an acid reversion product.

Following the procedure employed with glycogen,<sup>4</sup> amylopectin (waxy maize starch, 20.0 g.) in 0.4% concentration was hydrolyzed at  $100^\circ$  in 0.08 N hydrochloric acid for ten hours (degree of hydrolysis ca. 85%). After ion removal with exchange resins (Duolite C-3 and A-4), the amorphous solid obtained on solvent removal was acetylated with hot acetic anhydride and sodium acetate. A portion (5.9 g.) of the resultant sugar acetate mixture (39.0 g.) was chromatographed<sup>5</sup>

(1) M. L. Wolfrom, L. W. Georges, A. Thompson and I. L. Miller, THIS JOURNAL, **69**, 473 (1947); **71**, 2873 (1949).

(2) Edna M. Montgomery and G. E. Hilbert, *ibid.*, **68**, 916 (1946).

(3) Edna M. Montgomery, F. B. Weakley and G. E. Hilbert, *ibid.*, **69**, 2249 (1947); **71**, 1682 (1949).

(4) M. L. Wolfrom and A. N. O'Neill, *ibid.*, **71**, 3857 (1949).

(5) M. L. Wolfrom, L. W. Georges and I. L. Miller, *ibid.*, **71**, 125 (1949).