

TABLE I

Product, ^a methyl <i>p</i> -carbomethoxy-	Varying component	Re- fluxing time (hr.)	M.p., °C.	Yield, %	Sapon. equiv.		Analyses		Found, %	
					Calcd.	Found	Calcd., % C	% H	C	H
1 Phenoxyacetate	Methyl α -chloroacetate	15	93.6-94.5	63.9	112.1	112.4	58.92	5.39	58.75	5.38
2 Phenoxybutyrate	Methyl γ -chlorobutyrate	40	48.8-49.8	21.0	126.1	127.5	61.94	6.35
3 Phenoxyvalerate	Methyl δ -chlorovalerate	78	48.0-49.0	52.9	133.1	133.6	63.14	6.81	63.21	6.83
4 Phenoxycaproate	Methyl ϵ -chlorocaproate	82	44.6-45.4	52	140.2	141.0	64.27	7.19	64.35	7.10
5 Phenoxyheptanoate	Methyl ω -bromoheptano- ate	39	54.2-55.3	20	147.2	147.2	65.29	7.53	65.33	7.44
6 Phenoxyundecanoate	Methyl 11-bromoundecano- ate	24	69.0-70.0	91.5	175.2	175.6	68.57	8.57

^a All products were crystallized from methyl alcohol-water; however, where unreacted methyl *p*-hydroxybenzoate was present, it was removed by washing a benzene solution of the product with aqueous NaOH.

Freshly cut sodium (33.1 g., 1.44 moles) was gradually introduced into 600 ml. of dry methanol in a two-liter flask equipped with a stirrer, reflux condenser with a drying tube, and dropping funnel. Upon complete solution of the sodium, first, 219.1 g. (1.44 moles) of methyl *p*-hydroxybenzoate in 400 ml. of methanol and later 217.6 g. (1.44 moles) of methyl δ -chlorovalerate were added rapidly. The reaction mixture was refluxed for a total of 78 hours during which time the newly formed sodium chloride was filtered away several times as a means of determining the end of reaction. Upon pouring the cooled reaction mixture into ice-water, a white lustrous precipitate settled out. The product after recrystallization from methyl alcohol-water contained a small amount of a slightly higher melting impurity (assumed to be unreacted methyl *p*-hydroxybenzoate) which was removed by washing a benzene solution of the precipitate with aqueous sodium hydroxide. Physical data pertaining to this compound as well as to the other dimethyl esters are found in Table I.

Intermediates necessary to the preparation of these esters were made according to methods found in literature. The following references were employed: for methyl γ -chlorobutyrate,³ methyl ϵ -chlorocaproate,² methyl ω -bromoheptanoate² and methyl 11-bromoundecanoate.^{4,5}

(3) C. F. H. Allen, ref. 2, Coll. Vol. 1, 1941, p. 156.

(4) Conrad, *Ann.*, **188**, 218 (1877).

(5) R. Ashton and J. C. Smith, *J. Chem. Soc.*, 435 (1934).

RECEIVED SEPTEMBER 25, 1950

Ethyl 3 β -Acetoxy-5,17(20)-pregnadiene-21-thiolate

BY ROGER W. JEANLOZ¹

In a study of reductive desulfuration with Raney nickel, ethyl 3 β -acetoxy-5,17(20)-pregnadiene-21-thiolate has been prepared directly from 3 β ,17 β -diacetoxy-5-androstene-17 α -acetic acid² and from 3 β -acetoxy-5,17(20)-pregnadiene-21-ic acid.

3 β -Acetoxy-5,17(20)-pregnadiene-21-ic Acid.—By the use of acetic acid and pyridine, 150 mg. of 3 β -hydroxy-5,17(20)-pregnadiene-21-ic acid was acetylated. The product was crystallized from ether and gave 165 mg. (93%) of prisms, melting at 135-137°; $[\alpha]_D^{20}$ -74° (c, 1.04 in chloroform).³

Anal. Calcd. for C₂₃H₃₂O₄: C, 74.16; H, 8.66. Found: C, 74.10; H, 8.69.

Ethyl 3 β -Acetoxy-5,17(20)-pregnadiene-21-thiolate.—Thionyl chloride (1.5 ml.) was added at 0° to 270 mg. of dry 3 β ,17 β -diacetoxy-5-androstene-17 α -acetic acid (m.p. 200°)¹ and kept at 0° for 12 hours with exclusion of moisture. The excess thionyl chloride was evaporated and the last

(1) Worcester Foundation for Experimental Biology, Shrewsbury, Massachusetts.

(2) T. Reichstein, H. Müller, Ch. Meystre and M. Sutter, *Helv. Chim. Acta*, **22**, 741 (1939); Pl. A. Plattner and W. Schreck, *ibid.*, **22**, 1178 (1939).

(3) The melting points were determined on the Kofler micro melting point apparatus and the rotations observed in a 1-dm. tube of 0.8-ml. capacity.

traces removed by addition of absolute benzene and evaporation at reduced pressure. The crystalline acid chloride (not isolated) was dissolved in 2 ml. of absolute benzene and, after addition of 0.3 ml. of ethyl mercaptan and 0.1 ml. of absolute pyridine, was kept at room temperature for two days. The solution was then diluted with a small amount of water and extracted with ether. The extract was washed with dilute hydrochloric acid, sodium carbonate and water and dried over anhydrous sodium sulfate. The ethereal solution was decolorized with charcoal, evaporated to dryness and the residue, recrystallized from ether-pentane, yielded 255 mg. (95%) of colorless prismatic needles, m.p. 154-155°; $[\alpha]_D^{20}$ -65° (c, 1.38 in chloroform).

Anal. Calcd. for C₂₅H₃₀O₅S: C, 72.07; H, 8.71; S, 7.70. Found: C, 71.57; H, 9.15; S, 7.84.

One hundred and twenty-five milligrams of 3 β -acetoxy-5,17(20)-pregnadiene-21-ic acid was treated with thionyl chloride and then with ethyl mercaptan, as described above for 3 β ,17 β -diacetoxy-5-androstene-17 α -acetic acid. There was obtained 130 mg. (90%), melting at 150-152° (mixed m.p. with the specimen described above, 151-154°).

FRANK W. HORNER, LTD.

MONTREAL, QUE., CANADA RECEIVED NOVEMBER 20, 1950

Further Studies of the Ion-Exchange Separation of the Rare Earths¹

BY B. H. KETELLE AND G. E. BOYD

Recently Higgins and Street² have called attention to the mislabelling of the Tb activity peak in our Fig. 6c which reported on the ion-exchange column separation of all fourteen of the trivalent rare earths using citrate buffers at 100°.³ Partly reported^{4a,b} further work in the Tb-Gd-Eu region conducted at this Laboratory in 1948 and 1949 has revealed not only this mistake, in agreement with Higgins and Street, but has shown also that the peak marked Y in Fig. 6c was actually Dy.

Before discussing our later ion-exchange separations, the reasons for these misidentifications, which we take this occasion to correct, may be of interest: The earlier assignment of the activity peak in Fig. 6c occurring at 2400 minutes to Y was based partly upon the argument that after such a time dysprosium activity (assumed to be 145 m Dy¹⁶⁵) would not have been detectable, and partly

(1) This document is based on work performed under Contract Number W-7405 eng 26 for the Atomic Energy Project at Oak Ridge National Laboratory.

(2) G. H. Higgins and K. Street, *THIS JOURNAL*, **72**, 5321 (1950).

(3) B. H. Ketelle and G. E. Boyd, *ibid.*, **69**, 2800 (1947).

(4) (a) B. H. Ketelle, *Phys. Rev.*, **76**, 1256 (1949). (b) See private communication K(101) by B. H. Ketelle (Dec. 1948) to G. T. Seaborg concerning the decay sequence 3.6 m Gd¹⁶¹ → 7.0 d Tb¹⁶¹ reported in a Table of Isotopes by G. T. Seaborg and I. Periman, *Rev. Mod. Phys.*, **20**, 585 (1948).