

epinochrome; however, adrenochrome under non-reductive conditions as described above remains unchanged.

5,6-Dihydroxy-N-methylindole was first obtained by Duliere and Raper⁵ by intramolecular oxidation-reduction in the presence of sulfur dioxide of the red solution of enzymatically oxidized N-methyltyrosine (or from a red solution resulting from oxidation of epinine with silver oxide). They isolated it as the dimethyl ether which was subsequently identified by Burton⁶ who had observed the same rearrangement in the presence of alkali and who also prepared the diacetate. 5,6-Dihydroxy-N-methylindole itself was first isolated by partial reduction of adrenochrome,³ after Bergel and Morrison⁷ had demonstrated its formation by reduction of iodoadrenochrome, but isolated its diacetate only.

In a communication that has just come to our attention, Harley-Mason and Bu'Lock⁸ report that zinc ion catalyses the decolorization of the red solution from the oxidation of epinine or of its carboxy derivative. Our observations on the spontaneous rearrangement of pure epinochrome in the solid state and on the catalytic rearrangement of its solutions over palladium emphasize the facility with which this oxidation-reduction occurs under a variety of conditions.

(5) Duliere and Raper, *Biochem. J.*, **24**, 239 (1930).

(6) Burton, *J. Chem. Soc.*, 546 (1932).

(7) Bergel and Morrison, *ibid.*, 48 (1943).

(8) Harley-Mason and Bu'Lock, *Nature*, **166**, 1036 (1950).

DEPARTMENT OF CHEMISTRY
THE MOUNT SINAI HOSPITAL
NEW YORK 29, N. Y.

JOHN AUSTIN
J. D. CHANLEY
HARRY SOBOTKA

RECEIVED JANUARY 16, 1951


SYNTHESIS OF 11-KETO STEROIDS

Sir:

We wish to report a practical and general scheme for the synthesis of 11-keto steroids from $\Delta^{5,6}$ steroids, devoid of functional groups in ring C, which has been successfully applied to ergosterol, stigmasterol, diosgenin and to certain of their degradation products.

Starting with ergosterol acetate, the derived known $\Delta^{7,9(11),22}$ ergostatrien-3 β -ol acetate (ergosteryl-D-acetate)¹ is treated with one equivalent of perbenzoic acid to give a crystalline epoxide; m.p. 202–205°; $\alpha_D -35^\circ$ (CHCl_3); found: C, 79.01; H, 10.50. Hydrolytic rearrangement of the epoxide yielded $\Delta^{5,22}$ -ergostadien-3 $\beta,7,11$ -triol 3-acetate (I) m.p. 248–252°; $\alpha_D +85^\circ$ (CHCl_3); found: C, 76.04; H, 10.24; active hydrogens (Zerewitinoff determination) 2.1. Chromic acid oxidation of the triol monoacetate afforded $\Delta^{8,22}$ -ergostadien-3 β -ol-7,11-dione acetate, II; m.p. 135–136°; $\alpha_D +18.5^\circ$ (CHCl_3); λ_{max} 266 m μ , E_m 9360 (isoöctane); λ_{max} 270 m μ , E_m 8700 (alcohol); found: C, 76.91; H, 9.58. The dienedione on reduction with zinc and acetic acid was converted into Δ^{22} -ergostene-

3 β -ol-7,11-dione (III); m.p. 197–200°; $\alpha_D -30^\circ$ (CHCl_3); found: C, 76.68; H, 7.59. Modified Wolff-Kishner reduction² of the latter provided a monoketone, Δ^{22} -ergostene-3 β -ol-11-one (IV); m.p. 173–174°; $\alpha_D +26.6^\circ$ (CHCl_3); found: C, 81.72; H, 11.29. Ozonolysis of the acetylated ketoergostene, and esterification of the acidic side chain degradation product yielded an ester which proved to be methyl 3 β -hydroxy-11-keto-bisnorallocholanate acetate (V); m.p. 191–194°; $\alpha_D +24^\circ$ (CHCl_3); found: 71.89; H, 9.15. The structure of the degradation product was unequivocally established by relating it to methyl 3,11-diketo-bisnorallocholanate (VI) (m.p. 201–204°; $\alpha_D +63^\circ$; found: C, 74.18; H, 9.20) prepared by an independent synthesis from the known methyl 3-acetoxy-11-keto-bisnorcholelanate (VII).³



Methyl $\Delta^{5,7}$ -3-hydroxy-bisnorcholelanate acetate, obtained from either stigmasterol or cholesterol, was converted into methyl $\Delta^{5,7}$ -3-hydroxybisnorcholelanate acetate⁴ which was transformed into methyl 3-hydroxy-7,11-diketo-bisnorallocholanate acetate (m.p. 228–230°; $\alpha_D -16^\circ$ (CHCl_3); found: C, 69.30; H, 8.46) in the manner described for the ergosterol series. Wolff-Kishner reduction of the latter and subsequent esterification of the reduction product afforded V identical with that prepared from ergosterol.

Diosgenin acetate was converted to the Δ^7 -dehydrodiosgenin acetate⁵ which by the same sequence

(2) Huang-Minlon, *THIS JOURNAL*, **68**, 2487 (1946).

(3) Sarett, *J. Biol. Chem.*, **162**, 601 (1946).

(4) Prepared previously by Bergmann and Stevens, *J. Org. Chem.*, **13**, 10 (1948).

(5) Since completion of this work, the preparation of this compound was reported by Rosenkranz, Romo and Berlin, *J. Org. Chem.*, **16**, 290 (1951). The physical constants of the two products are essentially the same.

(1) Windaus and Brunken, *Ann.*, **460**, 225 (1928); Heilbron and Sexton, *J. Chem. Soc.*, 921 (1929); Windaus and Luttringhaus, *Ann.*, **481**, 119 (1930); Heilbron, Johnstone and Spring, *J. Chem. Soc.*, 2248 (1929).

of reactions yielded spirostan-3 β -ol-11-one acetate (11-ketotigogenin acetate), m.p. 224–229°; α_D -39.4° (CHCl₃); found: C, 73.60; H, 9.08. Pyrolysis of the ketogenin, oxidation of the pseudogenin and alkaline treatment of the latter provided Δ^{16} -allopregnene-3 β -ol-11,20-dione acetate, m.p. 183–185°; α_D +64.5° (CHCl₃); λ_{\max} . 234.5 m μ , E_m 9050 (alcohol); found: C, 73.93; H, 8.83. By hydrogenation of the latter, allopregnane-3 β -ol-11,20-dione acetate (m.p. 141–143° (VIII)); α_D +88° (CHCl₃); found: C, 73.79; H, 8.90) was obtained which was identical with the product obtained by the Barbier-Wieland degradation of VII.

E. M. CHAMBERLIN
W. V. RUYLE
A. E. ERICKSON
J. M. CHERMERDA
L. M. ALIMINOSA
R. L. ERICKSON
G. E. SITA
M. TISHLER

MERCK & CO., INC.
RAHWAY, NEW JERSEY

RECEIVED APRIL 26, 1951

11-KETOLITHOCHOLIC ACID AND 11-KETOCHOLESTANOL FROM THE $\Delta^{7,9(11)}$ -DIENES

Sir:

A practical route from cholic acid to methyl 3 α -7 α -diacetoxy-12-keto- $\Delta^{9(11)}$ -cholenate and methyl 12-keto- $\Delta^{7,9(11)}$ -lithocholadienate has been reported.¹ Dr. S. Rajagopalan found that both esters are converted in good yield by Wolff-Kishner reduction into $\Delta^{7,9(11)}$ -lithocholadienic acid (m.p. 201°, dec., $[\alpha]_D$ +121° Di; methyl ester, m.p. 120°, $[\alpha]_D$ +119° Di, λ_{\max}^{EtOH} 244.5 m μ (log ϵ 4.2)), which forms a methyl ester acetate m.p. 149–150°, $[\alpha]_D$ +123° Di, λ_{\max}^{EtOH} 244.5 (log ϵ 4.2) (C₂₇H₄₀O₄: C, 75.67; H, 9.64; found: C, 75.56; H, 9.63). We have now found that oxidation of this ester acetate with sodium dichromate dihydrate in acetic acid yields methyl 3 α -acetoxy-7,8-oxido- $\Delta^9(11)$ -cholenate (m.p. 185°, $[\alpha]_D$ +22° Di; C₂₇H₄₀O₅: C, 72.94; H, 9.07; found: C, 73.24; H, 9.29) and methyl 3 α -acetoxy-7,11-diketo- Δ^8 -cholenate (m.p. 115°, $[\alpha]_D$ +36° Di, λ_{\max}^{EtOH} 271 m μ , log ϵ 3.85; C₂₇H₃₈O₆: C, 70.72; H, 8.35; found: C, 70.65; H, 8.38). The unsaturated oxide, formed also by perbenzoic acid oxidation, was isomerized by zinc and acetic acid, aqueous dioxane at 160°, or methanolic potassium hydroxide (re-esterification) to methyl 3 α -acetoxy-7-keto- Δ^8 -cholenate (m.p. 182.5° $[\alpha]_D$ -15° Di, λ_{\max}^{EtOH} 254 m μ , log ϵ 4.04; found: C, 72.88; H, 9.13). Reduction of the unsaturated diketone with zinc and acetic acid afforded methyl 3 α -acetoxy-7,11-diketocholenate (m.p. 162°, $[\alpha]_D$ +25° Di, C₂₇H₄₀O₆: C, 70.40; H, 8.75; found: C, 70.28; H, 8.94), which on Wolff-Kishner reduction followed by esterification, acetylation, and chromatography gave both methyl lithocholate acetate (m.p. 134°, $[\alpha]_D$ +44° An, no depression with authentic sample²) and methyl 3 α -acetoxy-11-ketocholenate³


(1) L. F. Fieser, S. Rajagopalan, M. Tishler and E. Wilson, THIS JOURNAL, in press.

(2) Comparison sample: L. F. Fieser and S. Rajagopalan, THIS JOURNAL, **72**, 5530 (1950).

(3) Comparison sample: R. B. Turner, V. R. Mattox, L. L. Engel, B. F. McKenzie and E. C. Kendall, J. Biol. Chem., **166**, 345 (1946).

(m.p. 131°, $[\alpha]_D$ +67° An, C₂₇H₄₂O₅: C, 72.61; H 9.48; found: C, 72.60, H, 9.69, no depression with authentic sample.

Δ^7 -Cholestenol (m.p. 123–125°), available from cholesterol by Raney-nickel hydrogenation of the 7-dehydro derivative, was converted by dehydrogenation with mercuric acetate and benzooylation into $\Delta^{7,(911)}$ -cholestadienyl benzoate, m.p. 134°, $[\alpha]_D$ +32° Di, λ_{\max}^{EtOH} 243 m μ (log ϵ 4.0) (free alcohol), C₃₄H₄₈O₂: C, 83.55; H, 9.90; found: C, 83.81; H, 10.05). Oxidation of this diene with sodium dichromate in acetic acid-benzene at 25° yielded, after chromatography, Δ^8 -cholestene-3 β -ol-7,11-dione-3-benzoate (m.p. 150°, $[\alpha]_D$ about +6° Di, λ_{\max}^{EtOH} 268 m μ , log ϵ 3.8, λ_{\max}^{Chf} 5.95 μ , C₃₄H₄₆O₄: C, 78.72; H, 8.94; found: C, 78.77; H, 9.19); a second oxidation product is formed but



has not been fully characterized. Reduction of the enedione with zinc dust and acetic acid gave cholestan-3 β -ol-7,11-dione-3-benzoate, m.p. 200°, $[\alpha]_D$ +3° Di, C₃₄H₄₈O₄: C, 78.42; H, 9.29; found: C, 78.37; H, 9.71. Alternately, the total product from oxidation of the diene benzoate (8 g.) was reduced with zinc dust and acetic acid. Chromatography of the mixture yielded the saturated diketone benzoate (m.p. 200°), some of the steryl benzoate (m.p. 142°, $[\alpha]_D$ -10° Di), and 7-keto- Δ^8 -cholesteryl benzoate (m.p. 150°, $[\alpha]_D$ -13° Di, λ_{\max}^{EtOH} 252 m μ , log ϵ 3.8, C₃₄H₄₈O₃: C, 80.90; H, 9.58; $\lambda_{\max}^{CHCl_3}$ 5.95 μ ; found: C, 81.02; H, 9.87). Wolff-Kishner reduction of the saturated diketone gave a cholestanolone, m.p. 151°, $[\alpha]_D$ +49° Di, λ_{\max}^{Chf} 5.89 μ ; C₂₇H₄₆O₂: C, 80.54; H, 11.52; found: C, 80.66; H, 11.49. That cholesterol has been converted into the 11-keto stanol is evidenced by analogy to the synthesis in the bile acid series, by the presence of a carbonyl group (infrared) resistant to Wolff-Kishner reduction, and the M_D increment of +96° for the carbonyl group, as compared with known values (+79°, mean of 5 examples, +60 to +96°.⁴

CONVERSE MEMORIAL LABORATORY
HARVARD UNIVERSITY
CAMBRIDGE, MASSACHUSETTS

LOUIS F. FIESER
JOSEF E. HERZ
WEI-YUAN HUANG

RECEIVED APRIL 25, 1951

(4) D. H. R. Barton and W. Klyne, Chemistry and Industry, **26**, 757 (1948).