

described.<sup>3</sup> The molecular weight obtained was  $420,000,000 \pm 100,000,000$  with R, the root-mean-square radius about the center of mass, equal to  $14,000 \pm 4000 \text{ \AA}$ . The extreme angular dependence of the scattering caused the precision of the results to be lower than usual. This result is not in real disagreement with Potter and Hassid's molecular weight of 6,000,000, the latter being actually a lower limit in view of the well-known lack of precision of the osmotic method for this molecular weight range.

A sample of acetylated amylose also was measured. The weight-average molecular weight was about ten times the number-average value reported by Potter and Hassid, but the angular dependence of the scattered light indicated the presence of a few per cent. of high molecular weight impurity (possibly amylopectin).

(3) B. H. Zimm, *J. Chem. Phys.*, **16**, 1099 (1948); P. Outer, C. I. Carr and B. H. Zimm, *ibid.*, **18**, 830 (1950).

The angular dependence of the amylopectin scattering was of the normal type for a chain polymer. There was no evidence that the material was heterogeneous. Moreover, the measurements were repeated at  $70^\circ$  with almost identical results, indicating that the solution was actually a molecular dispersion without temperature-dependent aggregates. This is the first time, to our knowledge, that a molecular weight of this magnitude has been reported for an amylopectin.

We are indebted to Professor Hassid for his kind cooperation with this work.

(4) General Electric Research Laboratory, The Knolls, Schenectady, New York.

(5) Bell Telephone Laboratories, Murray Hill, New Jersey.

CHEMICAL DEPARTMENT  
UNIVERSITY OF CALIFORNIA  
BERKELEY 4, CALIFORNIA

B. H. ZIMM<sup>4</sup>

CARL D. THURMOND<sup>5</sup>

RECEIVED JANUARY 29, 1952

## BOOK REVIEWS

**Die biogenen Amine.** 4th edition. By M. GUGGENHEIM, Dr. phil., Dr. med. h. c., Basel. Interscience Publishers, Inc., 250 Fifth Avenue, New York 1, N. Y. (S. Karger, Holbeinstrasse 22, Basle, Switzerland.) 1951. xv + 619 pp.  $17 \times 23.5 \text{ cm}$ . \$19.50 (Swiss francs 75.-).

The new edition of Guggenheim's now classical monograph on the biogenic amines represents an extensive revision of the previous one. It follows the general plan of the earlier editions, presenting detailed information on the chemical, biochemical and pharmacological properties of these materials, their isolation, identification, quantitative determination, probable biogenesis and biochemical reactions. Some material has been omitted to make way for the great mass of information which has been added to our knowledge of these substances in the past decade. This is reflected in the fact that there has been approximately a forty per cent. increase in the number of substances covered with only a ten per cent. increase in the number of pages or references, now approximately 4600. Certain changes in the pattern of the chapters reflect the trend of recent discoveries. In particular, the discussions on biogenesis have been expanded and an introductory general chapter on the subject added.

A blemish in the chemical formulations throughout the book is the retention of uncharged structures for the amino acids. More serious is the perpetuation of uncharged forms for the guanidinium bases. While the author recognizes on page 396 that the guanidinium cation exists in solution, his formulation of guanidine as  $\text{CH}_5\text{N}_3$  helps to carry on the common misconception that non-resonating, anhydro-base forms of this and related materials are commonly isolated.

In the formulations of the reactions of biogenesis, it would have been desirable if the author had critically examined all equations proposing the assumption of formaldehyde as a biochemical intermediate. Some of these formulations should undoubtedly be replaced by transmethylation or formate ion reactions. Only those should be included in which investigators succeed in obtaining positive evidence that formaldehyde as such is utilized as an intermediate.

The new feature of this edition is the series of short but excellent reviews of enzymatic reactions, such as transmethylation, oxidative deamination and many others, in which biogenic amines participate as substrates, as coenzymes and as products. Such reactions help to bring the biochemical significance of these substances into clearer focus and to make their pharmacological manifestations more intelligible.

The work as a whole is a valuable and welcome review. The excellent quality of craftsmanship entering into the manufacture of the book is entirely in keeping with the high standards of scholarship displayed in the preparation of the manuscript.

DEPARTMENT OF CHEMISTRY  
THE JOHNS HOPKINS UNIVERSITY  
BALTIMORE 18, MARYLAND

ALSOPH H. CORWIN

**Organic Syntheses.** An Annual Publication of Satisfactory Methods for the Preparation of Organic Chemicals. Volume 31. By R. S. SCHREIBER, Editor-in-Chief. John Wiley and Sons, Inc., 440 Fourth Avenue, New York 16, N. Y. 1951. vi + 122 pp.  $15.5 \times 23.5 \text{ cm}$ . Price, \$2.75.

Ever since Volume 1 in this series appeared, organic chemists have looked forward with keen anticipation to the appearance of the next volume. The standards of publications throughout have been exceptionally high and the present volume is no exception.

Detailed directions are given for the preparation of forty organic compounds which include not only aliphatics and aromatics but some phosphorus compounds which are rapidly becoming important.

As usual there is a great deal of variety in the types of reactions and a mere study of the reactions involved is a good lesson in organic chemistry. No student of the subject, and no worker in the field, can afford to be without these volumes.

DOMINION RUBBER Co. LTD.  
RESEARCH LABORATORIES  
GUELPH, ONTARIO

R. H. MANSKE