

TABLE I
 MERCURIC NITRATE-PYRIDINE BASE ADDITION COMPOUNDS

(a) Desiccator dried; (b) vacuum dried; (c) nitric acid decomposition, reference 5; (d) hydrogen peroxide decomposition, reference 4

Pyridine base	M.p., °C.		Analyses (% Hg)				Calcd. for 2 base/1Hg(NO ₃) ₂
	(a)	(b)	(c)	(a) (d)	(c) (b)	(d)	
Pyridine	234-236 ^e	233-235	41.6	42.2	40.7	41.7	42.4
Quinoline	178 dec. ^f	179-181 dec.	34.8	35.1	35.0	35.2	34.6
β-Picoline	131-133 dec.	174-175 dec.	37.6	37.2	38.4	38.0	39.3
Lepidine	148 dec.	137 dec.	35.2	34.5	31.8	32.1	32.8
2,3-Lutidine	104-106 dec.	145	44.5	46.6	38.1	38.0	37.2 ^g
α-Picoline	122-125 dec.	109-115 dec.	51.0	48.7	50.4 ^h

^e Pyridine nitrate, m.p. 118, sublimes during determination. Reported to decompose without melting; reference 2.
^f Reported, m.p. 183°; reference 3. ^g 1/1 Ratio; Hg. calcd. 46.5%. ^h Value calcd. for 1/2 ratio.

curic nitrate except for the 2,3-lutidine complex and the α-picoline complex for which the analyses agreed with a one to one ratio and a one to two ratio, respectively. After vacuum drying the analyses were in at least fair agreement with the values prior to vacuum drying except again for the 2,3-lutidine complex which analyzed for a two to one base to salt ratio. This change in ratio, involving a loss of mercury, can be explained as a loss of the volatile nitrate from the less stable one to one complex. The change in melting point on vacuum drying observed with this compound is consistent with the possibility of such a change. The α-picoline complex, which sublimed on vacuum drying, was not analyzed as a vacuum-dried sample. These analytical data do not require that these complexes contain water of crystallization.

Experimental

The mercuric nitrate reagent was prepared by suspending 216.6 g. (1 mole) of yellow mercuric oxide in 91.4 g. of concd. nitric acid and stirring until solution was complete. The addition compounds were precipitated by adding 0.1 mole of the base to 26.3 g. (0.1 mole) of the reagent with stirring and cooling over a five-minute period. The precipitated solid was collected on a filter and recrystallized from water (pyridine, β-picoline, quinoline, lepidine) or from ethanol (α-picoline, 2,3-lutidine). The solids were dried in a desiccator over anhydrous calcium chloride and potassium hydroxide or in drying pistol over phosphorus pentoxide at 2-6 mm. Melting points are uncorrected. Mercury analysis were made by procedures previously described.^{4,5} Data are recorded in the Table.

DEPARTMENT OF CHEMISTRY
 COLLEGE OF ARTS AND SCIENCES
 UNIVERSITY OF LOUISVILLE
 LOUISVILLE 8, KENTUCKY

NEW COMPOUNDS

Tetraallylethylenediamine and Hexaallylethylenediammonium Dibromide

In the course of polymerization studies in this Laboratory,¹ tetraallylethylenediamine was prepared as an intermediate for the preparation of hexaallylethylenediammonium dibromide. The physical constants of these compounds have not been previously reported. Since polymerization studies of the quaternary ammonium salt as well as the properties of the polymer have been reported in a later paper,² it seems important that the preparation and properties of these compounds be recorded.

- (1) G. B. Butler and R. I. Bunch, *THIS JOURNAL*, **71**, 3120 (1949).
 (2) G. B. Butler, R. I. Bunch and F. L. Ingley, *ibid.*, **74**, 2543 (1952).

Tetraallylethylenediamine.—To a suspension of 67.2 g. of NaHCO₃ in 65 cc. of water was added 69 g. (0.71 mole) of diallylamine. With stirring, 66 g. (0.35 mole) of ethylene bromide was added dropwise. After the addition was complete, the mixture was heated under reflux for nine hr. After cooling, the sodium bromide was removed by filtration, and the filtrate saturated with NaOH. The amine layer was separated, dried over solid NaOH, and distilled; yield 32 g. (41%), b.p. 92° (3 mm.), *d*₂₅²⁵ 0.8517, *n*_D²⁵ 1.4702.

Anal. Calcd. for C₁₄H₂₄N₂: N, 12.71. Found: N, 12.62.

Hexaallylethylenediammonium Dibromide.—To 58 g. (0.26 mole) of tetraallylethylenediamine dissolved in 50 cc. of dry acetophenone, was added dropwise, with stirring, 63 g. (0.52 mole) of allyl bromide. The solution became warm on addition of the allyl bromide. Stirring was continued for four hr. after which time the salt began to precipitate. After an additional two hr., the salt was removed by filtration, washed several times with dry ether and dried; yield 108 g. (90%), m.p. 140°. After recrystallization from ethanol, the product had a melting point of 142°.

Anal. Calcd. for C₂₀H₃₄N₂Br₂: Br, 34.57. Found: Br, 34.28.

DEPARTMENT OF CHEMISTRY
 UNIVERSITY OF FLORIDA
 GAINESVILLE, FLORIDA

GEORGE B. BUTLER
 ROBERT L. BUNCH

RECEIVED FEBRUARY 29, 1952

Some Sulfonium and Selenonium Salts¹

Continuing the search for additional compounds which cause damage to tumor cells *in vivo*,² salts have been prepared by reaction of methyl sulfide, methyl ethyl sulfide, ethyl sulfide, bis-(2-hydroxyethyl) sulfide, and ethyl selenide with *p*-fluorophenacyl chloride, *p*-phenylphenacyl bromide and β-naphthacyl bromide. In some cases the reaction was carried out in refluxing alcohol solution according to the method of Bost and Schultzer,³ but in other cases best results were obtained by mixing the reactants and allowing them to stand at room temperature. Attempts to prepare sulfonium salts from bis-(2-cyanoethyl) sulfide were unsuccessful. The methods used are illustrated below and the properties of the products are listed in Table I. The salts were white or light buff solids which decomposed slowly on standing at room temperature and melted with decomposition on heating. The β-naphthacyl and *p*-phenylphenacyl salts were very slightly soluble in water (about 1 to 4 mg./ml.), while all the *p*-fluorophenacyl salts, except that of ethyl selenide, were more soluble.

Method I.—A solution of 7.25 g. of *p*-fluorophenacyl chloride in a slight excess of methyl sulfide was kept at room temperature for 1 day, then 6.8 g. (65%) of *p*-fluorophenacyl dimethyl sulfonium chloride was recovered by suction filtration and recrystallized from warm methanol by gradual

(1) This research was supported in part by a grant from the National Cancer Institute, of the National Institutes of Health, Public Health Service, and in part by a grant from the Damon Runyon Memorial Fund for Cancer Research.

(2) C. T. Bahner, *Texas Reports on Biology and Medicine*, **8**, 448 (1950); Henry A. Rutter, Jr., *THIS JOURNAL*, **78**, 5905 (1951).

(3) R. W. Bost and H. C. Schultzer, *ibid.*, **64**, 1165 (1942).

TABLE I

Salts from	Method	Reaction time	Yield, %	M.p., °C.	Empirical formula	Tonic halogen, %	
						Calcd.	Found
Methyl sulfide and <i>p</i> -Fluorophenacyl chloride	I	1 day	65 ^a	139	C ₁₀ H ₁₂ ClFOS	15.11	14.94, 15.20
Methyl ethyl sulfide and <i>p</i> -Fluorophenacyl chloride	I	1 day	67 ^a	130	C ₁₁ H ₁₄ ClFOS	14.25	14.22, 13.90
<i>p</i> -Fluorophenacyl bromide	I	10 min.	57 ^b	127	C ₁₁ H ₁₄ BrFOS	27.26	26.93, 27.25
Ethyl sulfide and β -Naphthacyl bromide	II	5 min.	58 ^b	127.5	C ₁₆ H ₁₉ BrOS	23.55	23.57
Bis-(2-hydroxyethyl) sulfide and <i>p</i> -Fluorophenacyl chloride	I ^c	2 days	99 ^a	115	C ₁₂ H ₁₆ ClFO ₃ S	12.03	12.18, 12.05
β -Naphthacyl bromide	II	5 min.	99 ^a	121-121.5	C ₁₆ H ₁₉ BrO ₃ S	21.52	21.65, 21.58
<i>p</i> -Phenylphenacyl bromide	II	20 min.		122.5-123	C ₁₈ H ₂₁ BrO ₃ S	20.11	20.03, 20.21
Ethyl selenide and <i>p</i> -Fluorophenacyl bromide	I	5 min.	85 ^a	96.5-99	C ₁₂ H ₁₆ BrFOSe	22.57	22.67, 22.74
β -Naphthacyl bromide	II	5 min.	58 ^a	106.5	C ₁₆ H ₁₉ BrOSE	20.70	20.86, 20.59

^a Crude yield. ^b Yield purified product. ^c Reacted at 40°.

addition of isopropyl ether (m.p. 139°). (Repeated recrystallization lowered the melting point, possibly as a result of reaction of the salt with the solvent.)

Method II.—A solution of 2.49 g. of β -naphthacyl bromide (0.01 mole) and 1.37 g. of ethyl selenide (0.01 mole) in 35 ml. of methanol was refluxed five minutes after all the solid β -naphthacyl bromide had dissolved. The β -naphthacyl diethyl selenonium bromide was precipitated by addition of isopropyl ether and recrystallized twice from methanol by addition of isopropyl ether; yield 2.2 g. (58%), m.p. 110-110.5°.

We wish to express our thanks to Mrs. Lydia Moore Rives and Miss Dorothy Marie Ellis for carrying out halogen analyses on these compounds.

CARL T. BAHNER

CARSON-NEWMAN COLLEGE
JEFFERSON CITY, TENNESSEE

PETER P. NEBLETT, JR.
HENRY A. RUTTER, JR.

RECEIVED FEBRUARY 1, 1952

to cause 100% inhibition at 1.5 p.p.m. These results were not unexpected in view of the work of Zentmeyer, Albert and others.³

We wish to express our thanks to C. J. Wessel and H. G. Shirk of the National Research Council Prevention of Deterioration Center for the screening tests carried out under Contract N7onr-291, Task Order 27, under sponsorship of the Departments of Air Force, Army and Navy and to Miss Dorothy M. Ellis for analyses.

(3) G. A. Zentmeyer, *Science*, **100**, 294 (1944); G. A. Zentmeyer, *Phytopathology*, **33**, 1121 (1943); A. Albert, *Med. J. Aust.*, **1**, 245 (1944); A. Albert, et al., *Brit. J. of Exp. Path.*, **28**, 69 (1947).

CARSON-NEWMAN COLLEGE
JEFFERSON CITY, TENNESSEE

CARL TABB BAHNER
BILLY LEE STUMP

RECEIVED FEBRUARY 21, 1952

8-Hydroxyquinolinium Salts¹

1-(*p*-Chlorophenacyl)-8-hydroxyquinolinium Bromide.—A mixture of 4.67 g. of *p*-chlorophenacyl bromide (0.02 mole) and 2.90 g. of 8-hydroxyquinoline (0.02 mole) was heated 30 min. at 100°. The resulting brown mass was dissolved in hot 95% ethanol and ethyl acetate was added to throw out 3.9 g. (52%) of crude quaternary salt. After recrystallization from warm ethanol by addition of ethyl ether the white crystals decomposed at 235-236°; solubility in water about 0.6% at 50°.

Anal. Calcd. for C₁₇H₁₃BrClNO₂: Br, 21.10. Found: Br, 21.08, 21.28.

1-Allyl-8-hydroxyquinolinium Bromide.—A solution of 4.35 g. (0.03 mole) of 8-hydroxyquinoline in 4.84 g. (0.04 mole) of allyl bromide was heated 6.5 hr. at 70° and the resulting yellow solid was recrystallized from hot absolute ethanol by addition of ethyl ether; crude yield 6.5 g. (81%). Repeated recrystallization yielded deep yellow crystals, decomposing at 148°; solubility in water about 10% at 26°.

Anal. Calcd. for C₁₂H₁₂BrNO: Br, 30.04. Found: Br, 30.26, 30.04.


When tested by the standard screening test of the Prevention of Deterioration Center these salts produced 18% and 11% growth inhibition, respectively, at a concentration of 250 p.p.m., whereas 8-hydroxyquinoline has been reported²

(1) This investigation was supported in part by a research grant from the National Cancer Institute, National Institutes of Health, Public Health Service, and in part by a grant from the Damon Runyon Memorial Fund for Cancer Research.

(2) R. E. Vicklund and M. Manowitz, *Technical Data Digest*, **15**, [5], 18 (1950).

Derivatives of Troger's Base

We have prepared the two quaternary salts of Troger's base listed below by the action of the appropriate halide on the crude base in ethanol. Purification was achieved by crystallization from ethanol-ether. The two compounds listed had no activity against Sarcoma 180 in mice.

R	X	M.p., °C. (uncor.)	Analyses, ^a % Calcd. Found
C ₆ H ₅ C ₆ H ₄ COCH ₂ —	Br	249.5-250	Br 15.21 15.03
	Cl	240 dec.	Cl 9.61 9.40

^a Mohr analyses.

Acknowledgment.—Grateful acknowledgment is made to Dr. C. Chester Stock of the Sloan-Kettering Institute for Cancer Research, for arranging the tests on these compounds.

DEPARTMENT OF CHEMISTRY
GEORGETOWN UNIVERSITY
WASHINGTON, D. C.

HENRY A. RUTTER, JR.

RECEIVED FEBRUARY 11, 1952