

Hill.⁷ The ozonide was reduced with zinc dust and acetic acid, and the acetaldehyde formed was collected in a solution⁸ of 2,4-dinitrophenylhydrazine. The phenylhydrazone weighed 2.5 g. (22%) and gave m.p. 164–166°. Admixture with authentic acetaldehyde 2,4-dinitrophenylhydrazone of m.p. 165–166° showed no depression in melting point. The recorded⁹ value for this derivative is 168°. The residual material resulting from the decomposition of the ozonide was purified in the previously described⁷ manner and 3.4 g. (54%) of 2-ethylcyclohexanone, n_D^{20} 1.4555, was obtained, the 2,4-dinitrophenylhydrazone of which gave a melting point of 133–134°.

Anal. Calcd. for $C_{14}H_{18}O_4N_4$: C, 54.89; H, 5.92; N, 18.29. Found: C, 55.21; H, 5.68; N, 18.60.

Since the melting point of the 2,4-dinitrophenylhydrazone differed from that reported in the literature¹⁰ (166°) this derivative was prepared from authentic 2-ethylcyclohexanone, derived from 2-chlorocyclohexanone, and this phenylhydrazone also gave a melting point of 133–134°. A mixed melting point with the derivative prepared from the ozonolysis product was unchanged. It would appear that 2-ethylcyclohexanone 2,4-dinitrophenylhydrazone exists in two forms.

The isolation of acetaldehyde and 2-ethylcyclohexanone, in fair yield, from the dehydration product of 1,2-diethylcyclohexanol of n_D^{20} 1.4665 indicates the structure of the olefin to be 1-ethyl-2-ethylidenecyclohexane rather than one of the expected cyclohexenes.

(7) A. L. Henne and P. Hill, *THIS JOURNAL*, **65**, 752 (1943).

(8) R. L. Shriner and R. C. Fuson, "Identification of Organic Compounds," Third Edition, John Wiley and Sons, Inc., New York, N. Y., 1948, p. 171.

(9) Reference 8, p. 229.

(10) F. E. King, J. A. Bartrop and R. J. Walley, *J. Chem. Soc.*, 277 (1945).

RESEARCH LABORATORIES
BLOCKSON CHEMICAL COMPANY
JOLIET, ILLINOIS

The Isolation of Reserpine from *Rauwolfia canescens* Linn

BY M. W. KLOHS, M. D. DRAPER, F. KELLER AND
F. J. PETRACEK

RECEIVED DECEMBER 18, 1953

The isolation of reserpine¹ from *Rauwolfia serpentina* Benth and the subsequent report of its sedative and hypotensive activity² has created widespread interest in this compound. In a search for other possible plant sources³ of this alkaloid we have recently isolated reserpine, by the method previously employed in our investigation of *Rauwolfia serpentina*,^{4,5} from the "oleoresin fraction" of *Rauwolfia canescens* Linn.

An earlier investigation of the alkaloidal fraction of *Rauwolfia canescens* by A. Chatterjee yielded one crystalline alkaloid, rauwolscline,⁶ to which the hypotensive activity of the crude drug has been ascribed.⁷ It now appears that there are at least two hypotensively active alkaloids present in this species.

(1) J. M. Müller, E. Schlittler and H. J. Bein, *Experientia*, **8**, 338 (1952).

(2) H. J. Bein, *ibid.*, **9**, 107 (1953).

(3) The isolation of reserpine from *Rauwolfia heterophylla* Roem. and Schult. has been reported in a recent communication by Carl Djerassi, Marvin Gorman, A. L. Nausbaum and J. Reynoso, *THIS JOURNAL*, **75**, 5446 (1953).

(4) M. W. Klohs, M. D. Draper, F. Keller and F. J. Petracek, *ibid.*, **75**, 4867 (1953).

(5) M. W. Klohs, M. D. Draper, F. Keller, W. Malesh and F. J. Petracek, *ibid.*, **76**, 1332 (1954).

(6) A. Chatterjee (née Mookerjee), *J. Indian Chem. Soc.*, **18**, 33 (1941).

(7) J. N. Mukherjee, *Science and Culture*, **18**, 338 (1953).

Experimental

The Isolation of Reserpine from the "Oleoresin Fraction."—The "oleoresin fraction" from the ground dried roots of *Rauwolfia canescens*⁸ was obtained in the same manner as previously described for fraction I in our investigation of *Rauwolfia serpentina*.⁵ Ten grams of this fraction was dissolved as much as possible in 2% methanol-chloroform (50 ml.). The solution was filtered and the filtrate was applied to a chromatographic column (5.5 × 25 cm.) containing silicic acid-celite 3:1. The column was developed with the same solvent system until the most rapid moving band had reached the bottom of the column. The adsorbent was then extruded and examined under ultraviolet light and with the aid of spot tests using Fröhdes reagent. A colorless zone in the lower section of the column, which exhibited blue fluorescence and the characteristic Fröhdes color reaction (yellow → yellow-green → light blue) for reserpine, was removed by sectioning and washed thoroughly with methanol (150 ml.). The methanol was concentrated *in vacuo* to approximately 10 ml. On the addition of several drops of ammonium hydroxide reserpine separated as flat needles (150 mg.). The material was recrystallized several times from methanol by dissolving in an excess and concentrating on the steam-bath; m.p. 254° dec., $[\alpha]_D^{25}$ -121.8° (*c* 1.03 in $CHCl_3$). A mixed melting point with an authentic sample gave no depression. The infrared and ultraviolet absorption spectra were identical. For analysis the sample was dried to constant weight at 110° (2 mm.).

Anal. Calcd. for $C_{23}H_{40}O_9N_2$: C, 65.11; H, 6.62; mol. wt., 608.67. Found: C, 65.13; H, 6.70; mol. wt., 611,¹² 614.¹³

(8) The plant material in this investigation was kindly identified by Dr. H. W. Youngken, Mass. College of Pharmacy, Boston, Mass.

(9) The empirical formula $C_{23}H_{40}O_9N_2$ originally proposed by us for reserpine has now been revised to $C_{22}H_{40}O_9N_2$ which is in agreement with that suggested by other investigators.^{3,10,11} Although the elementary analyses of reserpine and its derivatives reported by us fit both empirical formulas the results of more accurate equivalent weight determinations were incompatible with the original formula. The empirical formula for reserpinolic acid has likewise been changed to $C_{22}H_{38}O_8N_2$.

(10) A. Furlenmier, R. Lucas, H. B. MacPhillamy, J. M. Müller and E. Schlittler, *Experientia*, **9**, 331 (1953).

(11) N. Neuss, H. E. Boaz and J. W. Forbes, *THIS JOURNAL*, **75**, 4870 (1953).

(12) Potentiometric titration in glacial acetic acid with 0.01 *N* perchloric acid in dioxane.

(13) Potentiometric titration in 75% dimethylformamide-water with 0.01 *N* HCl.

RIKER LABORATORIES, INC.
LOS ANGELES 54, CALIFORNIA

1,1,1-Trichloro-2-methylpropyl-2-phosphoric Acid from 1,1,1-Trichloro-2-methyl-2-propanol

BY DONALD G. KUNDIGER AND THEODORE J. COLLIER

RECEIVED NOVEMBER 6, 1953

We have found that 1,1,1-trichloro-2-methyl-2-propanol (I) is converted to the intermediate $Cl_3C-C(CH_3)_2-O-PBr_4$ (III) by the action of phosphorus pentabromide upon I under anhydrous conditions. III can be hydrolyzed to 1,1,1-trichloro-2-methylpropyl-2-phosphoric acid (IV),¹ which can be obtained in 26% yield from (I).

The structure of IV was established by ultimate analyses, by neutralization equivalent and by cleavage with aqueous potassium hydroxide to potassium phosphate and the known cleavage products, *viz.*, carbon monoxide and acetone,² of I.

(1) *Cf.* the report by C. Willgerodt and F. Durr, *Ber.*, **20**, 539 (1887), wherein only replacement of the hydroxyl in I to give 1,1,1-trichloro-2-methyl-2-bromopropane (II) was indicated as the product of the reaction of phosphorus pentabromide upon I; no other product was indicated. This is the only previous report of this reaction.

(2) Bressanin and Segre, *Gass. chim. ital.*, **41**, [I] 673 (1911).