

ticular oxidation product that is similar to b". Allomerized in methanol or in barley leaves, b yields several pigments all more sorbed than b. One of these is spectrally similar to b (λ max. methanol, b, 652 m μ ; oxidized b, 653 m μ). This product, therefore, resembles b" both spectrally and chromatographically. Like b" it is not isomerized by heat or alkalis; and it does not give the phase test. Except for the method of preparation, this allomerized b is identical with b".

The a", for which few properties have been determined,³ has its counterpart among the oxidation products² of a. The increased sorbability of the oxidized chlorophylls² and of a" and b" relative to a and b may be attributed to additional oxygen (or decrease in hydrogen).⁴

Molecular structures proposed by Freed, *et al.*,³ for the chlorophylls and their isomers are unsuitable. If a, a', and oxidized a have the magnesium atom bound to different pairs of pyrrole nitrogen atoms,³ the three pigments should yield but one magnesium-free pheophytin. Yet, three have been reported.^{1,2} Moreover, oxidized chlorophylls, presumably with magnesium bound as in the chlorophylls, do not undergo isomerization.²

The strongly sorbed pigments prepared by Freed, *et al.*, and by Strain are oxidized chlorophylls, not isomeric chlorophylls. These products should not be called chlorophylls (unless found as natural constituents of plants).

(4) H. H. Strain, "Chromatographic Adsorption Analysis," Interscience Publishers, Inc., New York, N. Y., 1942, pp. 14-25.

CHEMISTRY DIVISION
ARGONNE NATIONAL LABORATORY HAROLD H. STRAIN
LEMONT, ILLINOIS

RECEIVED JULY 22, 1955

GLUTAMIC DEHYDROGENASE OF BEEF LIVER, A ZINC METALLOENZYME

Sir:

We recently have examined several crystalline preparations of beef liver glutamic dehydrogenase and find that zinc is an integral and functional component of the enzyme molecule.

The protein was crystallized from fresh beef liver.¹

Quantitative emission spectrography and microchemical determination of zinc were performed in duplicate according to methods previously described.² Protein weights were determined by trichloroacetic acid precipitation.²

Table I presents quantitative spectrochemical and microchemical data on three times crystallized preparations of glutamic dehydrogenase having high specific activity. Preparation 1 contained 333 μ g. Zn/g. of protein, preparation 2 contained 322 μ g. Zn/g. of protein, and a third preparation contained 260 μ g./g. The zinc content of these crystalline preparations was not lowered by prolonged dialysis. The molecular weight of glutamic dehydrogenase has been determined to be about one million.¹ Based on this molecular weight, the number of gram atoms of zinc per mole of enzyme protein vary between 4 and 5 for these preparations.

(1) J. A. Olson and C. B. Anfinsen, *J. Biol. Chem.*, **197**, 67 (1952).

(2) B. L. Vallee and F. L. Hoch, *Proc. Natl. Acad. Sci.*, **41**, 327 (1955).

TABLE I
ANALYSES OF THREE TIMES RECRYSTALLIZED GLUTAMIC DEHYDROGENASE OF BEEF LIVER

| Element | Preparation 1 (μ g./g. GDH) | Preparation 2 (μ g./g. GDH) |
|-----------|-------------------------------------|-------------------------------------|
| Zinc | 333 | 322 |
| Copper | " | ^b |
| Aluminum | 63.1 | ^b |
| Barium | 27.0 | 39.9 |
| Calcium | 1060 | ^b |
| Magnesium | 149 | 43.0 |

^a Not determined. ^b Not detected, and also beryllium, cadmium, cobalt, chromium, iron, lithium, molybdenum, manganese, nickel, potassium, silver, strontium, tin and lead.

Fractionation of beef liver shows an *aggregation* of zinc with purification of the enzyme, and as the activity:protein ratio reaches its maximum value. The metal:protein ratio of all *other elements* studied *decreases* with progressive enzyme purification. The activity:zinc ratio increases progressively with purification, reaching a high value in the third crystals.

The rate of conversion of DPN to DPNH at pH 7.7 in the presence of glutamate¹ is inhibited significantly when the enzyme is preincubated with a number of metal binding agents, including sodium sulfide, sodium diethyldithiocarbamate and 1,10-phenanthroline.

These data establish glutamic dehydrogenase as a zinc metalloenzyme according to the criteria published elsewhere.³

The oxidation of glutamic acid is the second DPN dependent dehydrogenation reaction known to be catalyzed by a zinc metalloenzyme.²

Acknowledgments.—The valuable suggestions of Dr. Eric G. Ball are gratefully acknowledged. This work was supported by grants-in-aid from the Office of Naval Research, Contract No. NR 119-277, to Harvard University, and by the Rockefeller Foundation, New York, N. Y.

BIOPHYSICS RESEARCH LABORATORY BERT L. VALLEE
OF THE DEPARTMENT OF MEDICINE S. JAMES ADELSTEIN⁴
HARVARD MEDICAL SCHOOL AND PETER
BENT BRIGHAM HOSPITAL, BOSTON, MASS.

DEPARTMENT OF CHEMISTRY JAMES A. OLSON
HARVARD UNIVERSITY
CAMBRIDGE, MASS.

(3) B. L. Vallee, "Zinc and Metalloenzymes" in "Advances in Protein Chemistry," Vol. X, Academic Press, New York, N. Y., 1955, in press.

(4) Fellow of the National Foundation for Infantile Paralysis.

STEROIDAL SAPOGENINS. XXXI. GENTROGENIN AND CORRELLOGENIN, NEW SAPOGENINS FROM *Dioscorea spiculiflora*¹

Sir:

Some years ago Marker announced the isolation of botogenin and neobotogenin from *D. mexicana*.^{2a,b} The structures assigned had both a 12-keto and a 5,6 double bond. Since substances with these groupings would be particularly desirable as cortisone precursors, there was an extensive but fruitless search made for them. We have now found in several collections of *D. spiculiflora*

(1) Paper XXX: M. E. Wall, submitted in *Experientia*.

(2) (a) R. E. Marker and J. Lopez, *THIS JOURNAL*, **69**, 2397 (1947); (b) R. E. Marker, *ibid.*, **71**, 2656 (1949).