

COMMUNICATIONS TO THE EDITOR

THE MELTING POINT OF NEO-C₅F₁₂

Sir:

In a recent publication¹ the preparation and detection of neo-C₅F₁₂ in a mixture of C₅F₁₂ isomers was reported. The detection of the neo-C₅F₁₂ was made with the aid of nuclear magnetic resonance analysis. When the mixture of C₅F₁₂ isomers was being removed from the sample tube in which the analysis was performed, it was noted that near the very end of the transfer some solid crystals remained in the tube. The crystals were sealed off in the tube and a melting point was determined. The melting range was observed to be 72–76°. When another aliquot of the mixture of C₅F₁₂ isomers was similarly treated, the crystalline residue was dissolved in pure CCl₄ and subjected to NMR analysis. The strong, single-peaked pattern obtained was interpreted as that which was characteristic of purified neo-C₅F₁₂.

More recently a larger sample of a mixture of C₅F₁₂ isomers was prepared in this laboratory. The mixture had a b.p. of 28.5–29.5° and a mol. wt. of 287–288 and was prepared from the reaction between (CF₃)₂SF₄ and CF₃CF=CF₂ at 520°. Neo-C₅F₁₂ enrichment was achieved by evaporation *in vacuo* at temperatures below 0°. Finally, a sample of material highly enriched in neo-C₅F₁₂ was isolated. This residual material was a crystalline solid, reminiscent of CO₂, and it exerted a vapor pressure of 650 ± 2 mm at 26°. The solid sample amounted to one gram.

Aliquots of the sample were successively transferred *in vacuo* to heavy wall glass melting point tubes 2 mm. in diameter and 4 cm. long. These aliquots amounted to about 0.2 g. each and were sealed off, air-free, in their respective tubes. The melting points of the five aliquots were determined in a water-bath with a thermometer and a stirrer. The temperature gradient over the depth of the bath did not exceed 0.2° and the rate of change of the temperature in the bath during the determinations did not exceed 0.25°/min. and was maintained at 0.1°/min. near and at the melting points. The following short table shows the melting range of the five successively condensed aliquots.

Aliquot Order	Melting range in °C.
1	72.0–75.8
2	73.3–76.1
3	75.3–77.6
4	76.4–78.3
5	76.3–78.2

Accordingly, with the highest purity that could be obtained by the procedure outlined, the m.p. of neo-C₅F₁₂, under its own vapor pressure, was at least 78.3°.

Grateful acknowledgment is offered to Pvts. N. Muller and George Svatos for performing the NMR analyses, which were done at the Army

(1) Richard Dreadner, *THIS JOURNAL*, **77**, 6633 (1955).

Chemical Center, Md. Appreciation is expressed to Drs. F. W. Hoffman and T. C. Simmons of the Chemical and Radiological Laboratories of A.C.C. for making the NMR analyses possible. The work in this laboratory was supported by the Chemistry Branch of the Office of Naval Research under contract Nonr 580(03); 333–356.

FLUORINE RESEARCH LABORATORIES
DEPARTMENT OF CHEMICAL ENGINEERING
UNIVERSITY OF FLORIDA
GAINESVILLE

R. D. DRESDNER

RECEIVED JANUARY 21, 1956

LIGHT CATALYZED ORGANIC REACTIONS. V.¹
THE ADDITION OF AROMATIC CARBONYL
COMPOUNDS TO A DISUBSTITUTED ACETYLENE

Sir:

In a previous paper of this series² we have described a method for the synthesis of substituted trimethylene oxides by light initiated addition of carbonyl compounds to polysubstituted olefins. With the view of exploring this reaction in greater detail, a series of experiments involving substituted acetylenes was carried out. The conversions outlined below constitute a new kind of combination leading to α,β -unsaturated ketones containing the system



A 3:1 mixture of benzaldehyde and 5-decyne was irradiated with a mercury resonance arc at 40° for 96 hours. Starting materials were removed by distillation, b.p. 40–60° (6 mm.) and V (13% yield) isolated by chromatography of the residue. 6-Benzylidene-5-decanone (V) had b.p. 108–111° (0.1 mm.), infrared max. 605, 6.24 μ ; 2,4-DNP m.p. 116–117°; *Anal.* Calcd. for C₂₃H₂₈N₄O₄: C, 65.07; H, 6.65; N, 13.20. Found: C, 64.87; H, 6.71; N, 13.26. Catalytic reduction of V over Pd/CaCO₃ catalyst in ethanol solution produced a homogeneous dihydro derivative (VII), infrared max. 5.85 μ ; *Anal.* Calcd. for C₁₇H₂₆O: C, 82.87; H, 10.64. Found: C, 82.69; H, 10.36. The structure of V was firmly established by unequivocal synthesis: α -Butylcinnamoyl chloride, b.p. 85° (0.35 mm.) secured from *trans*- α -butylcinnamic acid³ and thionyl chloride was condensed with dibutylcadmium to give V, b.p. 110° (0.06 mm.); λ_{max} 279 m μ , (log ϵ 4.2); *anal.* Calcd. for C₁₇H₂₄O: C, 83.55; H, 9.90. Found: C, 83.47; H, 9.72. The infrared spectra of this sample and the one obtained in the photochemical reaction were nearly identical. Further confirmation of identity was obtained by infrared and mixed melting point

(1) Part of a program of research supported by a grant from the Godfrey L. Cabot Fund, Publication No. 62, M.I.T. Solar Energy Conversion Project.

(2) G. Büchi, C. G. Inman and E. S. Lipinsky, *THIS JOURNAL*, **76**, 4327 (1954); E. Paterno and G. Chieffi, *Gazz. chim. ital.*, **39**, 341 (1909).

(3) A. Michael, *Chem. Ber.*, **34**, 918 (1901); M. Metayer, *Rec. trav. chim.*, **71**, 158 (1952).